

SCHEDULE

Regulation 6

Provisions to be substituted for Parts I and II of Schedule 5 to the Seed Potatoes Regulations 1984

“SCHEDULE 5 Regulations 2(1), 12(e), 18(1)(b) and 18(6)

PART I

SPECIFIED DISEASES OR PESTS, DAMAGE AND DEFECTS AND TOLERANCES—POTATOES (OTHER THAN VTSC CLASS) PRODUCED IN GREAT BRITAIN

<i>Column 1</i> <i>Specified diseases or pests, damage and defects</i>	<i>Column 2</i> <i>Individual tolerances</i>	<i>Column 3</i> <i>Group tolerances</i>	<i>Column 4</i> <i>Collective group tolerances</i>
GROUP I			
Wart Disease (<i>Synchytrium endobioticum</i> (Schilb.) Perc.)	NIL	—	—
Potato Tuber Eelworm (<i>Ditylenchus destructor</i> Thorne)	NIL	—	—
Potato Cyst Eelworm (<i>Globodera</i> species infesting potatoes)	NIL	—	—
Ring Rot (<i>Clavibacater michiganensis</i> (Smith) Davis <i>et al.</i> ssp. <i>sepedonicus</i> (Spieck. & Kotth.) Davis <i>et al.</i>)	NIL	—	—
Brown Rot (<i>Pseudomonas solanacearum</i> (E. F. Smith) Jensen)	NIL	—	—
Potato Tuber Moth (<i>Phthorimaea operculella</i> (Zell.))	NIL	—	—
Potato Spindle Tuber Viroid	NIL	—	—
Colorado Beetle (<i>Leptinotarsa decemlineata</i> (Say))	NIL	—	—
GROUP II			

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

<i>Column 1</i> <i>Specified diseases or pests, damage and defects</i>	<i>Column 2</i> <i>Individual tolerances</i>	<i>Column 3</i> <i>Group tolerances</i>	<i>Column 4</i> <i>Collective group tolerances</i>
Blight (<i>Phytophthora infestans</i> (Mont.) de Bary)	1.0%	1.0%	4.0%
Blackleg (<i>Erwinia carotovora</i> (Jones) Bergey <i>et al.</i> ssp. <i>atroseptica</i> (Hellmers & Dowson) Dye)	1.0%	1.0%	4.0%
Soft Rots including: Watery Wound Rot (<i>Pythium ultimum</i> Trow)	1.0%	1.0%	4.0%
Pink Rot (<i>Phytophthora erythroseptica</i> Pethybridge) and Pit Rot	1.0%	1.0%	4.0%
Dry Rot (<i>Fusarium</i> species)	1.0%	1.0%	4.0%
Gengrene (<i>Phoma</i> species)	1.0%	1.0%	4.0%
Frost damaged tubers	1.0%	1.0%	4.0%
GROUP III			
Skin Spot (<i>Polyscytalum</i> (Owen & Wakefield) M. B. Ellis)	2.0%	4.0%	4.0%
Powdery Scab (<i>Spongospora subterranea</i> (Wallr.) Lageth.)	3.0%	4.0%	4.0%
GROUP IV			
Black Scurf (<i>Rhizoctonia solant</i> Kühn)	3.0%	4.0%	4.0%
Common Scab (<i>Streptomyces</i> species)	4.0%	4.0%	4.0%
GROUP V			
External blemishes or tubers other than diseased tubers whose	2.0%	2.0%	4.0%

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

<i>Column 1</i> <i>Specified diseases or pests, damage and defects</i>	<i>Column 2</i> <i>Individual tolerances</i>	<i>Column 3</i> <i>Group tolerances</i>	<i>Column 4</i> <i>Collective group tolerances</i>
shape is atypical for the variety			
GROUP VI			
Dirt or other extraneous matter	1.0%	1.0%	

PART II

SPECIFIED DISEASES OR PESTS, DAMAGE AND DEFECTS AND TOLERANCES—VTSC CLASS

<i>Column 1</i> <i>Specified diseases or pests, damage and defects</i>	<i>Column 2</i> <i>Individual tolerances</i>	<i>Column 3</i> <i>Group tolerances</i>	<i>Column 4</i> <i>Collective group tolerances</i>
GROUP I			
Wart Disease (<i>Synchytrium endobioticum</i> (Schilb.) Perc.)	NIL	—	—
Potato Tuber Eelworm (<i>Ditylenchus destructor</i> Thorne)	NIL	—	—
Potato Cyst Eelworm (<i>Globodera</i> species infesting potatoes)	NIL	—	—
Ring Rot (<i>Clavibacater michiganensis</i> (Smith) Davis <i>et al.</i> ssp. <i>sepedonicus</i> (Spieck. & Kotth.) Davis <i>et al.</i>)	NIL	—	—
Brown Rot (<i>Pseudomonas solanacearum</i> (E. F. Smith) Jensen)	NIL	—	—
Potato Tuber Moth (<i>Phthorimaea operculella</i> (Zell.))	NIL	—	—
Potato Spindle Tuber Viroid	NIL	—	—

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

<i>Column 1</i> <i>Specified diseases or pests, damage and defects</i>	<i>Column 2</i> <i>Individual tolerances</i>	<i>Column 3</i> <i>Group tolerances</i>	<i>Column 4</i> <i>Collective group tolerances</i>
Colorado Beetle (<i>Leptinotarsa decemlineata</i> (Say))	NIL	—	—
GROUP II			
Blight (<i>Phytophthora infestans</i> (Mont.) de Bary)	0.5%	1.0%	4.0%
Blackleg (<i>Erwinia carotovora</i> (Jones) Bergey <i>et al.</i> ssp. <i>atroseptica</i> (Hellmers & Dowson) Dye)	NIL	1.0%	4.0%
Soft Rots including: Watery Wound Rot (<i>Pythium ultimum</i> Trow)	0.5%	1.0%	4.0%
Pink Rot (<i>Phytophthora erythroseptica</i> Pethybridge) and Pit Rot	0.5%	1.0%	4.0%
Dry Rot (<i>Fusarium</i> species)	0.5%	1.0%	4.0%
Gengrene (<i>Phoma</i> species)	0.5%	1.0%	4.0%
Frost damaged tubers	0.5%	1.0%	4.0%
GROUP III			
Skin Spot (<i>Polyscytalum</i> (Owen & Wakefield) M. B. Ellis)	0.5%	3.0%	4.0%
GROUP IV			
Black Scurf (<i>Rhizoctonia solant</i> Kühn)	1.5%	3.0%	4.0%
Common Scab (<i>Streptomyces</i> species)	3.0%	3.0%	4.0%
GROUP V			
Powdery Scab (<i>Spongospora subterranea</i> (Wallr.) Lageth.)	1.5%	3.0%	4.0%

Status: This is the original version (as it was originally made). This item of legislation is currently only available in its original format.

<i>Column 1</i> <i>Specified diseases or pests, damage and defects</i>	<i>Column 2</i> <i>Individual tolerances</i>	<i>Column 3</i> <i>Group tolerances</i>	<i>Column 4</i> <i>Collective group tolerances</i>
GROUP VI			
External blemishes or tubers other than diseased tubers whose shape is atypical for the variety	2.0%	2.0%	4.0%
GROUP VII			
Dirt or other extraneous matter	1.0%	1.0%”	