Document Generated: 2023-08-18

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

# [F1ANNEX II

## CONDITIONS TO BE SATISFIED BY THE SEED

### **Textual Amendments**

Substituted by Commission Directive 2009/74/EC of 26 June 2009 amending Council Directives 66/401/ EEC, 66/402/EEC, 2002/55/EC and 2002/57/EC as regards the botanical names of plants, the scientific names of other organisms and certain Annexes to Directives 66/401/EEC, 66/402/EEC and 2002/57/EC in the light of developments of scientific and technical knowledge (Text with EEA relevance).

## I.CERTIFIED SEED

1. The seed shall have sufficient varietal identity and purity.

In particular, the seeds of the species listed below shall conform to the following standards or other conditions. The minimum varietal purity shall be:

Poa pratensis, varieties referred to in the second part of the third sentence of paragraph 4 of Annex I, Brassica napus var. napobrassica and Brassica oleracea convar. acephala: 98 %, Pisum sativum, Vicia faba: certified seed, 1st generation: 99 %, certified seed, 2nd generation: 98 %, I<sup>F2</sup>Trifolium subterraneum, Medicago spp., except M. lupulina, M. sativa, M. x varia: for the production of basic seed: 99,5 %, for the production of certified seed for further multiplication: 98 %,

for the production of certified seed: 95 %.]

# **Textual Amendments**

Inserted by Commission Implementing Directive (EU) 2016/2109 of 1 December 2016 amending Council Directive 66/401/EEC as regards the inclusion of new species and the botanical name of the species Lolium x boucheanum Kunth (Text with EEA relevance).

The minimum varietal purity shall be examined mainly in field inspections carried out in accordance with the conditions laid down in Annex I.

The seed shall conform to the following standards or other conditions as regards 2. germination, analytical purity and content of seeds of other plant species, including the presence of bitter seed in sweet varieties of *Lupinus* spp.:

#### Table: A.

[F3Speciesminationalytical purity	Maximum content of seeds of other plant species in a sample of the weight specified in column 4 of	Conditions as regards content of seeds of Lupinus spp.
-----------------------------------	--	--

											(tot	nex II al per imn)		of another colour
	1		io <b>m</b> ú2		dr pla	ant sp	ecies(	of see % by	weig		Ave fatu	naCus aspp	spp	n <i>e</i> xnd
	pur	e har l) seed of pur seed	d by l(‰ei e		sing	gl <i>erep</i>	enmy	os <b>a pq</b> i	idexip	h anis	erisia <sup>r</sup>	ilis	than Run acen and Run	n lupin n <i>e</i> seeds osella
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15
	ceae amine	(a)												
Agro canii	s <b>TiS</b>	ue)	90	2,0	1,0	0,3	0,3				0	0 (j) (k)	2 (n)	
Agro capi	s <b>Ti5</b> ll( <b>a</b> c)s		90	2,0	1,0	0,3	0,3				0	0 (j) (k)	2 (n)	
Agro giga			90	2,0	1,0	0,3	0,3				0	0 (j) (k)	2(n)	
Agro stolo	sTis Mistera		90	2,0	1,0	0,3	0,3				0	0 (j) (k)	2 (n)	
Alop prate	etOru. en(su)s	S	75	2,5	1,0 (f)	0,3	0,3				0	0 (j) (k)	5 (n)	
Arrh elati	eīhāthe usa)	erum	90	3,0	1,0 (f)	0,5	0,3				0 (g)	0 (j) (k)	5 (n)	
Bron cath	ทนี5 a <b>(น</b> ัวเม	5	97	1,5	1,0	0,5	0,3				0 (g)	0 (j) (k)	10 (n)	
Bron sitch	ทนี§ e(ห)is		97	1,5	1,0	0,5	0,3				0 (g)	0 (j) (k)	10 (n)	
Cyno dacty			90	2,0	1,0	0,3	0,3				0	0 (j) (k)	2	
Dacı glom	ty <b>80</b> ne(ra))ta		90	1,5	1,0	0,3	0,3				0	0 (j) (k)	5 (n)	

Festu <b>&amp;0</b> arund <b>(n)</b> acea	95	1,5	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Festu <b>do</b> filifor <b>(rai)</b> s	85	2,0	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Festud <b>o</b> ovina(a)	85	2,0	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Festu&d praten(si)s	95	1,5	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Festud <b>ō</b> rubra(a)	90	1,5	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Festu <b>do</b> trach <b>yph</b> ylla	85	2,0	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
× 75 Festu <b>lal</b> jum	96	1,5	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Lolium5 multif( <b>a)</b> rum	96	1,5	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Lolium peren(m)	96	1,5	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Lolium5 × (a) hybridum	96	1,5	1,0	0,5	0,3	0	0 (j) (k)	5 (n)	
Phala <b>7fs</b> aquat <b>(e)</b>	96	1,5	1,0	0,3	0,3	0	0 (j) (k)	5	
Phleu80 nodos(a)n	96	1,5	1,0	0,3	0,3	0	0 (k)	5	
Phleu80 prater(sæ	96	1,5	1,0	0,3	0,3	0	0 (k)	5	
Poa 75 annu(a)	85	2,0 (c)	1,0 (c)	0,3	0,3	0	0 (j) (k)	5 (n)	
Poa 75 nemo <b>(a)</b> is	85	2,0 (c)	1,0 (c)	0,3	0,3	0	0 (j) (k)	2 (n)	

Council Directive of 14 June 1966 on the marketing of fodder plant seed...

ANNEX II

Document Generated: 2023-08-18

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After

IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

Poa palus	75 (vai)		85	2,0 (c)	1,0 (c)	0,3	0,3		0	0 (j) (k)	2 (n)	
Poa prate	75 (8.1)s		85	2,0 (c)	1,0 (c)	0,3	0,3		0	0 (j) (k)	2 (n)	
Poa trivia	75 ul(ias)		85	2,0 (c)	1,0 (c)	0,3	0,3		0	0 (j) (k)	2 (n)	
Trise flave			75	3,0	1,0 (f)	0,3	0,3		0 (h)	0 (j) (k)	2 (n)	
	aceae gumin		,			,						
Biser pelec	r <b>10</b> a		98	0,5					0 (i)	0 (j) (k)	10	
Gale		40	97	2,0	1,5			0,3	0	0 (l) (m)	10 (n)	
	<i>slt</i> brun n <b>(æ)</b> jun (b)		95	2,5	1,0			0,3	0	0 (k)	5	
Lathy			95	1	0,5			0,3	0 (i)	0 (j) (k)	20	
Lotus	s75 i <b>¢a]</b> ata (b)	40 us	95	1,8 (d)	1,0 (d)			0,3	0	0 (l) (m)	10	
Lupii	n&0 s(a) (b)	20	98	0,5 (e)	0,3 (e)			0,3	0 (i)	0 (j)	5 (n)	(o) (p)
Lupii	ก <b>ฟิร์</b> s( <b>tā</b> f)oli (b)	20 us	98	0,5 (e)	0,3 (e)			0,3	0 (i)	0 (j)	5 (n)	(o) (p)
Lupii luteu		20	98	0,5 (e)	0,3 (e)			0,3	0 (i)	0 (j)	5 (n)	(o) (p)
Medi dolia			98	2					0 (i)	0 (j) (k)	10	
Medi italic	cago (b)	20	98	2					0 (i)	0 (j) (k)	10	

Medic <b>tig</b> o littoralis		98	2			0 (i)	0 (j) (k)	10
Media (go lupuli(na) (b)	20	97	1,5	1,0	0,3	0	0 (l) (m)	10
Medic <b>u</b> go murex(b)	30	98	2			0 (i)	0 (j) (k)	10
Medi <b>čkig</b> o polym <b>(br)</b> ph		98	2			0 (i)	0 (j) (k)	10
Medi <b>člig</b> o rugos <b>(</b> b)	20	98	2			0 (i)	0 (j) (k)	10
Media go sativa(a) (b)	40	97	1,5	1,0	0,3	0	0 (l) (m)	10
Medi <b>ck(g</b> o scutellata		98	2			0 (i)	0 (j) (k)	10
Medi <b>čk</b> go trunc <b>áh)</b> la	20	98	2			0 (i)	0 (j) (k)	10
Medialgo × (a) varia (b)	40	97	1,5	1,0	0,3	0	0 (l) (m)	10
Onobīķāchi viciifa <b>ki</b> g (b)	is20	95	2,5	1,0	0,3	0	0 (j)	5
Ornit <b>H5</b> pu compressu	1	90	1			0 (i)	0 (j) (k)	10
Ornit <b>П5</b> ри sativus	S	90	1			0 (i)	0 (j) (k)	10
Pisum80 sativu( <b>n</b> )		98	0,5	0,3	0,3	0	0 (j)	5 (n)
Trifol <b>80</b> n alexa <b>nd</b> rin	20 ium	97	1,5	1,0	0,3	0	0 (l) (m)	10
Trifol <b>ilû</b> n fragiferum	,	98	1			0 (i)	0 (j) (k)	10

Trifo	1770m	30	98	1			0	0	10
gland	d <b>(lbi)</b> fei			1			(i)	(j) (k)	
Trifo hirtu	l <b>īl0</b> m m		98	1			0 (i)	0 (j) (k)	10
Trifo hybri		20	97	1,5	1,0	0,3	0	0 (l) (m)	10
Trifo incar	l <b>îl5</b> m <b>'1(a)</b> un (b)	20 n	97	1,5	1,0	0,3	0	0 (l) (m)	10
Trifo isthn	l <b>ilû</b> n 10carj	oum	98	1			0 (i)	0 (j) (k)	10
Trifo mich	l <b>îl5</b> m e <b>(li</b> c)nı	30 ım	98	1			0 (i)	0 (j) (k)	10
Trifo prate		20	97	1,5	1,0	0,3	0	0 (l) (m)	10
Trifo repei		40	97	1,5	1,0	0,3	0	0 (l) (m)	10
Trifo resup	l <b>80</b> m i(nac)tu (b)	20 m	97	1,5	1,0	0,3	0	0 (l) (m)	10
Trifo squa	lī <b>l5</b> m r <b>(b)</b> :ui	20 n	97	1,5		0,3	0	0 (l) (m)	10
Trifo subte	l <b>8</b> 0m en <b>lo</b> gne	40 rum	97	0,5			0 (i)	0 (j) (k)	10
Trifo vesic	l <b>īlû</b> n ulosu	m	98	1			0 (i)	0 (j) (k)	10
Trigo foeni graeo			95	1,0	0,5	0,3	0	0 (j)	5
Vicia beng	80 h(dd)en.	20 sis	97 (e)	1			0 (i)	0 (j) (k)	10
Vicia faba	1	5	98	0,5	0,3	0,3	0	0 (j)	5 (n)

Document Generated: 2023-08-18

Vicia 85 panna(ni)ca (b)	20	98	1,0 (e)	0,5 (e)		0,3			0 (i)	0 (j)	5 (n)	
Vicia 85 sativa(a) (b)	20	98	1,0 (e)	0,5 (e)		0,3			0 (i)	0 (j)	5 (n)	
Vicia 85 villos <b>ú</b> a) (b)	20	98	1,0 (e)	0,5 (e)		0,3			0 (i)	0 (j)	5 (n)	
Other species			·		1			'			'	
Brass Ran napus(a) var. napobrassi	ica	98	1,0	0,5			0,3	0,3	0	0 (j) (k)	5	
Brassika oleradad convar. acephala (acephala var. medullosa viridis)	+ var	98	1,0	0,5			0,3	0,3	0	0 (j) (k)	10	
Phacesus tanacesusfol	ia	96	1,0	0,5					0	0 (j) (k)		
Plantā <b>§</b> o lanceolata		85	1,5						0 (i)	0 (j) (k)	10	
Raph&Ous sativu(xa) var. oleiformis		97	1,0	0,5			0,3	0,3	0	(j)	5	]

- B. Other standards or conditions applicable where reference is made to them in the table under Section I (2) (A) of this Annex:
  - all fresh and healthy seeds which do not germinate after pre-treatment shall (a) be regarded as seeds which have germinated;
  - up to the maximum quantity indicated, hard seed present shall be regarded (b) as seed capable of germination;
  - a maximum total of 0,8 % by weight of seeds of other *Poa* species shall not (c) be regarded as an impurity;
  - a maximum of 1 % by weight of seeds of Trifolium pratense shall not be (d) regarded as an impurity;

- (e) [F3 a maximum total of 0,5 % by weight of seeds of *Lupinus albus*, *Lupinus angustifolius*, *Lupinus luteus*, *Pisum sativum*, *Vicia faba*, *Vicia* spp. in another relevant species shall not be regarded as an impurity;]
- (f) the prescribed maximum percentage by weight of seeds of a single species shall not apply to seeds of *Poa* spp;
- (g) a maximum total of two seeds of *Avena fatua*, and *Avena sterilis* in a sample of the prescribed weight shall not be regarded as an impurity where a second sample of the same weight is free from any seeds of these species;
- (h) the presence of one seed of *Avena fatua*, and *Avena sterilis* in a sample of the prescribed weight shall not be regarded as an impurity where a second sample of twice the prescribed weight is free from any seeds of these species;
- (i) the determination of seeds of *Avena fatua*, and *Avena sterilis* by number need not be carried out unless there is doubt whether the conditions laid down in column 12 have been satisfied;
- (j) the determination of seeds of *Cuscuta* spp. by number need not be carried out unless there is doubt whether the conditions laid down in column 13 have been satisfied;
- (k) the presence of one seed of *Cuscuta* spp. in a sample of the prescribed weight shall not be regarded as an impurity where a second sample of the same weight is free from any seeds of *Cuscuta* spp;
- (l) the weight of the sample for the determination of seeds of *Cuscuta* spp. by number shall be twice the weight specified in column 4 of Annex III for the relevant species;
- (m) the presence of one seed of *Cuscuta* spp. in a sample of the prescribed weight shall not be regarded as an impurity where a second sample of twice the prescribed weight is free from any seeds of *Cuscuta* spp;
- (n) the determination of seeds of *Rumex* spp. other than *Rumex acetosella* and *Rumex maritimus* by number need not be carried out unless there is doubt whether the conditions laid down in column 14 have been satisfied;
- (o) the percentage by number of seeds of *Lupinus* spp. of another colour shall not exceed:

—in 2 % bitter lupin, —in 1 % Lupinus spp. other than bitter lupin,

(p) the percentage by number of bitter seeds in varieties of *Lupinus* spp. shall not exceed 2,5 %.

Document Generated: 2023-08-18

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

### **Textual Amendments**

- Substituted by Commission Implementing Directive (EU) 2016/2109 of 1 December 2016 amending Council Directive 66/401/EEC as regards the inclusion of new species and the botanical name of the species Lolium x boucheanum Kunth (Text with EEA relevance).
- 3. I<sup>F4</sup>The seed shall be practically free from any pests which reduce the usefulness and quality of the seed.

### **Textual Amendments**

Substituted by Commission Implementing Directive (EU) 2020/177 of 11 February 2020 amending Council Directives 66/401/EEC, 66/402/EEC, 68/193/EEC, 2002/55/EC, 2002/56/ EC and 2002/57/EC, Commission Directives 93/49/EEC and 93/61/EEC and Implementing Directives 2014/21/EU and 2014/98/EU as regards pests of plants on seeds and other plant reproductive material (Text with EEA relevance).

The seed shall also comply with the requirements concerning Union quarantine pests, protected zone quarantine pests and RNQPs provided for in implementing acts adopted pursuant to Regulation (EU) 2016/2031, as well as the measures adopted pursuant to Article 30(1) of that Regulation.

The presence of RNQPs on the seeds and the respective categories shall comply with the following requirements as set out in the table:

RNQPs or symptoms caused by RNQPs	Plants for planting (genus or species)	Thresholds for pre-basic seed	Thresholds for basic seed	Thresholds certified seed
Clavibacter michiganensis ssp. Insidiosus (McCulloch 1925) Davis et al. [CORBIN]	Medicago sativa L.	0 %	0 %	0 %
Ditylenchus dipsaci (Kuehn) Filipjev [DITYDI]	Medicago sativa L.	0 %	0 %	0 %]

#### II. **BASIC SEED**

Subject to the provisions below, the conditions laid down in Section I of this Annex shall apply to basic seed:

1. The seed of Pisum sativum, Brassica napus var. napobrassica, Brassica oleracea convar. acephala, Vicia faba and of varieties of Poa pratensis referred to in the second part of the third sentence of paragraph 4 of Annex I shall conform to the following standards or other conditions: the minimum varietal purity shall be 99,7 %.

The minimum varietal purity shall be examined mainly in field inspections carried out in accordance with the conditions laid down in Annex I.

2. The seed shall satisfy the following other standards or conditions:

# A. Table:

[F3Speci	Other						
1	Total(% by weight)		standards or conditions				
		A single species	Rumex spp. other than Rumex acetose and Rumex maritim	Elytrigi repens Ila	a Alopeci myosur		us
1	2	3	4	5	6	7	8
Poaceae (Gramir							
Agrostis canina	0,3	20	1	1	1		(j)
Agrostis capillari		20	1	1	1		(j)
Agrostis gigantea		20	1	1	1		(j)
Agrostis stolonife		20	1	1	1		(j)
Alopecun	-	20 (a)	2	5	5		(j)
Arrhenai elatius	t <b>l</b> QgBum	20 (a)	2	5	5		(i) (j)
Bromus cathartic	0,4 us	20	5	5	5		(j)
Bromus sitchensi	0,4 s	20	5	5	5		(j)
Cynodor dactylon		20 (a)	1	1	1		(j)
Dactylis glomera		20 (a)	2	5	5		(j)
Festuca arundina		20 (a)	2	5	5		(j)

Festuca filiformis	0,3	20 (a)	2	5	5	(j)
Festuca ovina	0,3	20 (a)	2	5	5	(j)
Festuca pratensis	0,3	20 (a)	2	5	5	(j)
Festuca rubra	0,3	20 (a)	2	5	5	(j)
Festuca trachyph		20 (a)	2	5	5	(j)
× Festuloli	0,3 ium	20 (a)	2	5	5	(j)
Lolium multiflor	0,3 um	20 (a)	2	5	5	(j)
Lolium perenne	0,3	20 (a)	2	5	5	(j)
Lolium × hybridum	0,3	20 (a)	2	5	5	(j)
Phalaris aquatica		20	2	5	5	(j)
Phleum nodosum	0,3	20	2	1	1	(j)
Phleum pratense	0,3	20	2	1	1	(j)
Poa annua	0,3	20 (b)	1	1	1	(f) (j)
Poa nemorali	0,3 is	20 (b)	1	1	1	(f) (j)
Poa palustris	0,3	20 (b)	1	1	1	(f) (j)
Poa pratensis	0,3	20 (b)	1	1	1	(f) (j)
Poa trivialis	0,3	20 (b)	1	1	1	(f) (j)
Trisetum flavescer		20 (c)	1	1	1	(i) (j)
Fabaced (Legum)					,	
Biserrule pelecinu.	a0,3	20	5			

	1					
Galega 0,3 orientalis	20	2			0 (e)	(j)
Hedysarum3 coronarium	20	2			0 (e)	(j)
Lathyrus 0,3 cicera	20	5	_	_	0 (d)	
Lotus 0,3 corniculatus	20	3			0 (e)	(g) (j)
Lupinus 0,3 albus	20	2			0 (d)	(h) (k)
Lupinus 0,3 angustifolius	20	2			0 (d)	(h) (k)
Lupinus 0,3 luteus	20	2			0 (d)	(h) (k)
Medicag <sub>0</sub> ,3 doliata	20	5	_	_	0 (e)	
Medicag <sub>0</sub> ,3 italica	20	5	_	_	0 (e)	
Medicaga,3 littoralis	20	5	_	_	0 (e)	
Medicaga,3 lupulina	20	5	_	_	0 (e)	(j)
Medicag <sub>0,3</sub>	20	5	_	_	0 (e)	
Medicag <sub>0</sub> ,3 polymorpha	20	5	_	_		
Medicag <sub>0</sub> ,3 rugosa	20	5	_	_		
Medicag <sub>0</sub> ,3 sativa	20	3			0 (e)	(j)
Medicag <sub>0</sub> ,3 scutellata	20	5				
Medicag <sub>0</sub> ,3 truncatula	20	5				
Medicag <sub>0</sub> ,3 × varia	20	3			0 (e)	(j)
Onobryc <b>hi</b> ş3 viciifolia	20	2			0 (d)	
Ornithop <b>u</b> s3 compressus	20	5				

(j)
(j)
(j)
(j)
_
(j)
(j)
(j)
_
(j)
(j)
_
(h)

			,			
Vicia sativa	0,3	20	2		0 (d)	(h)
Vicia villosa	0,3	20	2		0 (d)	(h)
Other species			I			
Brassica napus var. napobra		20	2			(j)
Brassica oleracea convar. acephal (acephal var. medullos viridis)	a Ia	20	3			(j)
Phacelia tanacetij		20				
Plantago lanceola		20	3			
Raphani sativus var. oleiform		20	2			]

- B. Other standards or conditions applicable where reference is made to them in the table under Section II (2) (A) of this Annex:
  - (a) a maximum total of 80 seeds of *Poa* spp. shall not be regarded as an impurity;
  - (b) the condition laid down in column 3 is not applicable to the seeds of *Poa* spp. The maximum total content of seeds of *Poa* spp other than the species to be examined shall not exceed one in a sample of 500 seeds;
  - (c) a maximum total of 20 seeds of *Poa* spp. shall not be regarded as an impurity;
  - (d) the determination of seeds of *Melilotus* spp. by number need not be carried out unless there is doubt whether the conditions laid down in column 7 have been satisfied;
  - (e) the presence of one seed of *Melilotus* spp. in a sample of the prescribed weight shall not be regarded as an impurity where a second sample of twice the prescribed weight is free from any seeds of *Melilotus* spp;

- (f) the condition (c) laid down in Section I (2) of this Annex is not applicable;
- (g) the condition (d) laid down in Section I (2) of this Annex is not applicable;
- (h) the condition (e) laid down in Section I (2) of this Annex is not applicable;
- (i) the condition (f) laid down in Section I (2) of this Annex is not applicable;
- (j) the conditions (k) and (m) laid down in Section I (2) of this Annex are not applicable;
- (k) the percentage by number of bitter seeds in varieties of *Lupinus* spp. shall not exceed 1 %.

## III. COMMERCIAL SEED

Subject to the provisions below, the conditions laid down in Section I (2) and (3) of this Annex shall apply to commercial seed:

- 1. The percentages by weight laid down in columns 5 and 6 of the table under Section I (2) (A) of this Annex are increased by 1 %.
- 2. In *Poa annua* a maximum total of 10 % by weight of seeds of other *Poa* species shall not be regarded as an impurity.
- 3. In *Poa* spp. other than *Poa annua* a maximum total of 3 % by weight of seeds of other *Poa* species shall not be regarded as an impurity.
- 4. In *Hedysarum coronarium* a maximum total of 1 % by weight of seeds of *Melilotus* spp. shall not be regarded as an impurity.
- 5. The condition (d) laid down in Section I (2) of this Annex is not applicable to *Lotus corniculatus*.
- 6. In *Lupinus* spp.:
  - (a) the minimum analytical purity shall be 97 % by weight;
  - (b) the percentage by number of seeds of *Lupinus* spp. of another colour shall not exceed:

—ın	4 %
bitter	
lupin	
—in	2 %
Lupinus	
spp.	
other	
than	
bitter	
lupin	

- 7. [F3 In *Vicia* spp. a maximum total of 6 % by weight of seeds of *Vicia pannonica, Vicia villosa, Vicia benghalensis* or related cultivated species in another relevant species shall not be regarded as an impurity.]
- 8. [F3In *Vicia pannonica, Vicia sativa, Vicia villosa, Vicia benghalensis* the minimum analytical purity shall be 97,0 % by the weight.]
- 9. [F2For *Lathyrus cicera* the minimum analytical purity shall be 90 % by the weight. A maximum total of 5 % by the weight of seeds of similar cultivated species shall not be regarded as an impurity.]]