

This text is meant purely as a documentation tool and has no legal effect. The Union's institutions do not assume any liability for its contents. The authentic versions of the relevant acts, including their preambles, are those published in the Official Journal of the European Union and available in EUR-Lex. Those official texts are directly accessible through the links embedded in this document

► **B****COUNCIL DIRECTIVE****of 27 November 1990****on the fixing of maximum levels for pesticide residues in and on certain products of plant origin, including fruit and vegetables**

(90/642/EEC)

(OJ L 350, 14.12.1990, p. 71)

Amended by:

		Official Journal		
		No	page	date
► <u>M1</u>	Council Directive 93/58/EEC of 29 June 1993	L 211	6	23.8.1993
► <u>M2</u>	Council Directive 94/30/EC of 23 June 1994	L 189	70	23.7.1994
► <u>M3</u>	Council Directive 95/38/EC of 17 July 1995	L 197	14	22.8.1995
► <u>M4</u>	Council Directive 95/61/EC of 29 November 1995	L 292	27	7.12.1995
► <u>M5</u>	Council Directive 96/32/EC of 21 May 1996	L 144	12	18.6.1996
► <u>M6</u>	Council Directive 97/41/EC of 25 June 1997	L 184	33	12.7.1997
► <u>M7</u>	Commission Directive 97/71/EC of 15 December 1997	L 347	42	18.12.1997
► <u>M8</u>	Commission Directive 98/82/EC of 27 October 1998	L 290	25	29.10.1998
► <u>M9</u>	Commission Directive 1999/65/EC of 24 June 1999	L 172	40	8.7.1999
► <u>M10</u>	Commission Directive 1999/71/EC of 14 July 1999	L 194	36	27.7.1999
► <u>M11</u>	Commission Directive 2000/24/EC of 28 April 2000	L 107	28	4.5.2000
► <u>M12</u>	Commission Directive 2000/42/EC of 22 June 2000	L 158	51	30.6.2000
► <u>M13</u>	Commission Directive 2000/48/EC of 25 July 2000	L 197	26	3.8.2000
► <u>M14</u>	Commission Directive 2000/57/EC of 22 September 2000	L 244	76	29.9.2000
► <u>M15</u>	Commission Directive 2000/58/EC of 22 September 2000	L 244	78	29.9.2000
► <u>M16</u>	Commission Directive 2000/81/EC of 18 December 2000	L 326	56	22.12.2000
► <u>M17</u>	Commission Directive 2000/82/EC of 20 December 2000	L 3	18	6.1.2001
► <u>M18</u>	Commission Directive 2001/35/EC of 11 May 2001	L 136	42	18.5.2001
► <u>M19</u>	Commission Directive 2001/39/EC of 23 May 2001	L 148	70	1.6.2001
► <u>M20</u>	Commission Directive 2001/48/EC of 28 June 2001	L 180	26	3.7.2001
► <u>M21</u>	Commission Directive 2001/57/EC of 25 July 2001	L 208	36	1.8.2001
► <u>M22</u>	Commission Directive 2002/5/EC of 30 January 2002	L 34	7	5.2.2002
► <u>M23</u>	Commission Directive 2002/23/EC of 26 February 2002	L 64	13	7.3.2002
► <u>M24</u>	Commission Directive 2002/42/EC of 17 May 2002	L 134	29	22.5.2002
► <u>M25</u>	Commission Directive 2002/66/EC of 16 July 2002	L 192	47	20.7.2002
► <u>M26</u>	Commission Directive 2002/71/EC of 19 August 2002	L 225	21	22.8.2002
► <u>M27</u>	Commission Directive 2002/76/EC of 6 September 2002	L 240	45	7.9.2002
► <u>M28</u>	Commission Directive 2002/79/EC of 2 October 2002	L 291	1	28.10.2002

► <u>M29</u>	Commission Directive 2002/97/EC of 16 December 2002	L 343	23	18.12.2002
► <u>M30</u>	Commission Directive 2002/100/EC of 20 December 2002	L 2	33	7.1.2003
► <u>M31</u>	Council Regulation (EC) No 806/2003 of 14 April 2003	L 122	1	16.5.2003
► <u>M32</u>	Commission Directive 2003/62/EC of 20 June 2003	L 154	70	21.6.2003
► <u>M33</u>	Commission Directive 2003/60/EC of 18 June 2003	L 155	15	24.6.2003
► <u>M34</u>	Commission Directive 2003/69/EC of 11 July 2003	L 175	37	15.7.2003
► <u>M35</u>	Commission Directive 2003/113/EC of 3 December 2003	L 324	24	11.12.2003
► <u>M36</u>	Commission Directive 2003/118/EC of 5 December 2003	L 327	25	16.12.2003
► <u>M37</u>	Commission Directive 2004/2/EC of 9 January 2004	L 14	10	21.1.2004
► <u>M38</u>	Commission Directive 2004/59/EC of 23 April 2004	L 120	30	24.4.2004
► <u>M39</u>	Commission Directive 2004/61/EC of 26 April 2004	L 127	81	29.4.2004
► <u>M40</u>	Commission Directive 2004/95/EC of 24 September 2004	L 301	42	28.9.2004
► <u>M41</u>	Commission Directive 2004/115/EC of 15 December 2004	L 374	64	22.12.2004
► <u>M42</u>	Commission Directive 2005/37/EC of 3 June 2005	L 141	10	4.6.2005
► <u>M43</u>	Commission Directive 2005/46/EC of 8 July 2005	L 177	35	9.7.2005
► <u>M44</u>	Commission Directive 2005/48/EC of 23 August 2005	L 219	29	24.8.2005
► <u>M45</u>	Commission Directive 2005/70/EC of 20 October 2005	L 276	35	21.10.2005
► <u>M46</u>	Commission Directive 2005/74/EC of 25 October 2005	L 282	9	26.10.2005
► <u>M47</u>	Commission Directive 2005/76/EC of 8 November 2005	L 293	14	9.11.2005
► <u>M48</u>	Commission Directive 2006/9/EC of 23 January 2006	L 22	24	26.1.2006
► <u>M49</u>	Commission Directive 2006/4/EC of 26 January 2006	L 23	69	27.1.2006
► <u>M50</u>	Commission Directive 2006/30/EC of 13 March 2006	L 75	7	14.3.2006
► <u>M51</u>	Commission Directive 2006/53/EC of 7 June 2006	L 154	11	8.6.2006
► <u>M52</u>	Commission Directive 2006/59/EC of 28 June 2006	L 175	61	29.6.2006
► <u>M53</u>	Commission Directive 2006/60/CE of 7 July 2006	L 206	1	27.7.2006
► <u>M54</u>	Commission Directive 2006/61/EC of 7 July 2006	L 206	12	27.7.2006
► <u>M55</u>	Commission Directive 2006/62/EC of 12 July 2006	L 206	27	27.7.2006
► <u>M56</u>	Commission Directive 2006/92/EC of 9 November 2006	L 311	31	10.11.2006
► <u>M57</u>	Commission Directive 2007/7/EC of 14 February 2007	L 43	19	15.2.2007
► <u>M58</u>	Commission Directive 2007/12/EC of 26 February 2007	L 59	75	27.2.2007
► <u>M59</u>	Commission Directive 2007/8/EC of 20 February 2007	L 63	9	1.3.2007
► <u>M60</u>	Commission Directive 2007/9/EC of 20 February 2007	L 63	17	1.3.2007
► <u>M61</u>	Commission Directive 2007/11/EC of 21 February 2007	L 63	26	1.3.2007
► <u>M62</u>	Commission Directive 2007/27/EC of 15 May 2007	L 128	31	16.5.2007
► <u>M63</u>	Commission Directive 2007/28/EC of 25 May 2007	L 135	6	26.5.2007
► <u>M64</u>	Commission Directive 2007/39/EC of 26 June 2007	L 165	25	27.6.2007
► <u>M65</u>	Commission Directive 2007/55/EC of 17 September 2007	L 243	41	18.9.2007
► <u>M66</u>	Commission Directive 2007/56/EC of 17 September 2007	L 243	50	18.9.2007
► <u>M67</u>	Commission Directive 2007/57/EC of 17 September 2007	L 243	61	18.9.2007
► <u>M68</u>	Commission Directive 2007/62/EC of 4 October 2007	L 260	4	5.10.2007
► <u>M69</u>	Commission Directive 2007/73/EC of 13 December 2007	L 329	40	14.12.2007
► <u>M70</u>	Commission Directive 2008/17/EC of 19 February 2008	L 50	17	23.2.2008

Corrected by:

- **C1** Corrigendum, OJ L 219, 24.8.1994, p. 26 (93/58/EEC)
- **C2** Corrigendum, OJ L 155, 28.6.1996, p. 62 (95/38/EC)
- **C3** Corrigendum, OJ L 175, 10.7.1999, p. 83 (98/82/EC)
- **C4** Corrigendum, OJ L 262, 17.10.2000, p. 46 (2000/42/EC)
- **C5** Corrigendum, OJ L 140, 30.5.2002, p. 39 (2002/42/EC)

- ▶ C6 Corrigendum, OJ L 342, 30.12.2003, p. 58 (2002/79/EC)
- ▶ C7 Corrigendum, OJ L 14, 21.1.2004, p. 55 (2003/60/EC)
- ▶ C8 Corrigendum, OJ L 28, 31.1.2004, p. 30 (2004/2/EC)
- ▶ C9 Corrigendum, OJ L 98, 2.4.2004, p. 61 (2003/113/EC)
- ▶ C10 Corrigendum, OJ L 104, 8.4.2004, p. 135 (2003/113/EC)
- ▶ C11 Corrigendum, OJ L 5, 7.1.2005, p. 26 (2004/115/EC)
- ▶ C12 Corrigendum, OJ L 72, 18.3.2005, p. 50 (2004/115/EC)
- ▶ C13 Corrigendum, OJ L 140, 1.6.2007, p. 58 (2007/27/EC)

▼B**COUNCIL DIRECTIVE****of 27 November 1990****on the fixing of maximum levels for pesticide residues in and on certain products of plant origin, including fruit and vegetables**

(90/642/EEC)

*Article 1***▼M6**

1. This Directive shall apply to products within the groups specified in column 1 of Annex I, examples of which are given in column 2, in so far as products in those groups, or the parts of product described in column 3, may contain certain pesticide residues.

The Directive shall also apply to the same products after drying or processing or after inclusion in a composite food in so far as they may contain certain pesticide residues.

▼B

2. This Directive shall apply without prejudice to:

- (a) the provisions of Council Directive 64/54/EEC of 5 November 1963 on the approximation of the laws of the Member States concerning the preservatives authorized for use in foodstuffs intended for human consumption⁽¹⁾, as last amended by Directive 85/585/EEC⁽²⁾, relating to biphenyl (diphenyl), orthophenylphenol, sodium orthophenyl phenate and 2-(4-thiazolyl)-benzimidazole (thiabendazole), which shall continue to regulate the use of those substances until they and their maximum levels are included in the list referred to in paragraph 1;
- (b) the provisions of Council Directive 74/63/EEC of 17 December 1973 on the fixing of maximum permitted levels for undesirable substances and products in feedingstuffs⁽³⁾, as last amended by Directive 87/519/EEC⁽⁴⁾;
- (c) the provisions of Directive 76/895/EEC;
- (d) the provisions of Council Directive 86/362/EEC of 24 July 1986 on the fixing of maximum levels for pesticide residues in and on cereals⁽⁵⁾, as last amended by Directive 88/298/EEC⁽⁶⁾;

▼M6

- (e) the provisions of Commission Directive 91/321/EEC of 14 May 1991 on infant formulae and follow-on formulae⁽⁷⁾ and Commission Directive 96/5/EC of 16 February 1996 on processed cereal-based foods and baby foods for infants and young children⁽⁸⁾. However, until maximum levels have been established in accordance with Article 6 of Directive 91/321/EEC or Article 6 of Directive 96/5/EC the provisions of Article 5a (1) and (3) to (6) of this Directive shall apply for the products concerned.

⁽¹⁾ OJ No 12, 27. 1. 1964, p. 161/64.

⁽²⁾ OJ No L 372, 31. 12. 1985, p. 43.

⁽³⁾ OJ No L 38, 11. 2. 1974, p. 31.

⁽⁴⁾ OJ No L 304, 27. 10. 1987, p. 38.

⁽⁵⁾ OJ No L 221, 7. 8. 1986, p. 37.

⁽⁶⁾ OJ No L 126, 20. 5. 1988, p. 53.

⁽⁷⁾ OJ No L 175, 4. 7. 1991, p. 35. Directive as last amended by Directive 96/4/EC (OJ No L 49, 28. 2. 1996, p. 12).

⁽⁸⁾ OJ No L 49, 28. 2. 1996, p. 17.

▼B

3. This Directive shall also apply to products referred to in paragraph 1 intended for export to third countries. However, maximum pesticide residue levels set in accordance with this Directive shall not apply in the case of products treated before export where it can be satisfactorily proved that:

- (a) the third country of destination requires that particular treatment in order to prevent the introduction of harmful organisms into its territory; or
- (b) the treatment is necessary in order to protect the products against harmful organisms during transport to the third country of destination and storage there.

4. This Directive shall not apply to the products referred to in paragraph 1 where it can be established by appropriate evidence that they are intended for:

- (a) the manufacture of products other than foodstuffs and animal feed; or
- (b) sowing or planting.

Article 2

For the purposes of this Directive:

▼M6

- (a) ‘pesticide residues’ shall mean residues of pesticides and of their metabolites, and breakdown or reaction products, which are present in or on the products referred to in Article 1.

▼B

- (b) ‘putting into circulation’ shall mean any post-harvest handing over, whether or not for a consideration, of the products referred to in Article 1.

▼M6*Article 3*

1. The products in the groups or, where applicable, the parts of products referred to in Article 1 shall not contain, from the time they are put into circulation, pesticide residue levels higher than those specified in the list referred to in Annex II.

The list of pesticide residues concerned and their maximum levels shall be established in Annex II in accordance with the procedure laid down in Article 10a having regard to current scientific and technical knowledge. A pesticide residue will be included on the list for as long as Directive 76/895/EEC fixes a maximum level for that residue.

2. In the case of dried and processed products for which maximum levels are not explicitly fixed in Annex II, the maximum residue level applicable shall be that laid down in Annex II, taking into account, respectively, the concentration caused by the drying process or the concentration or dilution caused by processing. A concentration or dilution factor covering the concentration and/or dilution caused by certain drying or processing operations may be determined for certain dried or processed products in accordance with the procedure laid down in Article 10a.

▼ **M6**

3. In the case of compound foods which contain a mixture of ingredients and for which maximum residue levels are not fixed, the maximum residue levels applied may not exceed the levels laid down in Annex II, taking into account the relative concentrations of the ingredients in the mixture and also the provisions of paragraph 2.

4. Member States shall ensure, at least by check sampling, compliance with the maximum levels referred to in paragraph 1. The necessary inspections and monitoring shall be carried out in accordance with Council Directive 89/397/EEC of 14 June 1989 on the official control of foodstuffs⁽¹⁾, except for Article 14 thereof, and Directive 93/99/EEC of 29 October 1993 on the subject of additional measures concerning the official control of foodstuffs⁽²⁾ except for Articles 5, 6 and 8 thereof.

Article 4

1. Member States shall designate an authority to ensure that the monitoring specified in Article 3 (4) is carried out.

2. (a) By ► **M9** 30 September ◀ each year, Member States shall send to the Commission their forward national monitoring programmes for the following calendar year. These forward programmes shall specify at least:

- the products to be inspected and the number of inspections to be carried out,
- the pesticide residues to be inspected,
- the criteria applied in drawing up these programmes.

(b) By ► **M9** 31 December ◀ each year, the Commission shall submit to the Standing Committee on Plant Health a draft recommendation setting out a coordinated Community monitoring programme identifying the taking of specific samples to be included in the national monitoring programmes. The recommendation shall be adopted in accordance with the procedure laid down in Article 10. The basic objective of the Community monitoring programme shall be to make optimum use at Community level of the sampling of plant products included in the groups listed in Annex I, produced in the Community or imported into it, when problems have been identified, in order to ensure compliance with the maximum levels for pesticide residues set out in Annex II.

3. By 31 August each year, Member States shall send to the Commission and the other Member States the results of the analyses of the samples taken during the previous year under their national monitoring programmes and under the coordinated Community monitoring programme. The Commission shall collate and combine this information together with the results of the checks carried out in accordance with Directives 86/362/EEC and 86/363/EEC and analyse:

- infringements of the maximum residue levels, and
- the average actual levels of residues and their relative values with respect to the maximum residue levels established.

⁽¹⁾ OJ No L 186, 30. 6. 1989, p. 23.

⁽²⁾ OJ No L 290, 24. 11. 1993, p. 14.

▼M6

The Commission should progressively work towards a system, when preparing the coordinated monitoring programme, which could permit the estimation of actual pesticide dietary exposure.

The Commission shall forward this information to the Member States in the framework of the Standing Committee on Plant Health before ►**M9** 31 December ◀ for each year, for review and adoption of any necessary measures such as:

- any action to be taken at Community level in the case of reported infringements of the maximum levels,
- the desirability of publication of the collated and compiled information.

4. The following may be adopted in accordance with the procedure laid down in Article 9:

- (a) amendments to paragraphs 2 and 3 of this Article in so far as these amendments concern the dates for notification;
- (b) detailed implementing rules necessary for proper functioning of the provisions of paragraphs 2 and 3.

5. Not later than 31 December 1999 the Commission shall forward to the Council a report on the application of this Article, accompanied, if necessary, by any appropriate proposals.

▼B*Article 5*

Member States may not prohibit or impede the putting into circulation within their territories of the products referred to in Article 1 on the grounds that they contain pesticide residues, if the quantity of such residues in and on the products or parts of products concerned does not exceed the maximum levels specified in the list referred to in Article 1.

▼M6*Article 5a*

Where for a product belonging to a group referred to in Annex I, a provisional maximum residue level applicable throughout the Community is set by the Commission in accordance with the provisions of Article 4 (1) (f) of Council Directive 91/414/EEC of 15 July 1991 concerning the placing of plant protection products on the market⁽¹⁾, this level will be indicated in Annex II with a reference to that procedure.

Article 5b

1. For the purposes of this Article a Member State of origin shall be defined as the Member State in whose territory a product specified in Article 1 (1) is either legally produced and marketed or put into free circulation and a Member State of destination as the Member State into whose territory such product is introduced and put into circulation for operations other than transit to another Member State or third country.

⁽¹⁾ OJ No L 230, 19. 8. 1991, p. 1. Directive as last amended by Directive 96/32/EC (OJ No L 144, 18. 6. 1996, p. 12).

▼M6

2. Member States shall introduce arrangements for establishing maximum residue levels, whether permanent or temporary, for products referred to in Article 1 (1), brought into their territories from a Member State of origin, taking into account good agricultural practice in the Member State of origin, and without prejudice to conditions necessary to protect the health of consumers, in cases where no maximum residue levels have been established for these products in accordance with the provisions of Articles 3 (1) or 5a.

3. Where

— no maximum residue level has been established for a product referred to in Article 1 (1) in accordance with Articles 3 (1) or 5a, and

— that product, which satisfies the maximum residue levels applied by its Member State of origin, has been subjected in the Member State of destination to measures whose effect is to prohibit or restrict its putting into circulation, on the grounds that the product contains pesticide residue levels in excess of the maximum residue level accepted in the Member State of destination, and

— either the Member State of destination has introduced new maximum residue levels or has altered the levels laid down in its legislation, or it has made changes to its controls which are disproportionate and/or discriminatory compared with those for its domestic production, or the maximum residue level applied by the Member State of destination differs substantially from the corresponding levels established by other Member States, or the maximum residue level applied by the Member State of destination represents a disproportionate level of protection compared with the level of protection applied by the Member State to pesticides carrying a similar risk or to similar agricultural products or foodstuffs,

the following exceptional provisions shall apply:

(a) the Member State of destination shall communicate the measures adopted to the other Member State concerned and the Commission within 20 days of their application. The notification shall document the facts involved;

(b) on the basis of the notification referred to in (a), the two Member States concerned shall contact each other without delay in order to remove, whenever possible, the prohibitive or restrictive effect of the measures adopted by the Member State of destination by means of measures agreed between them; the Member States shall submit all the requisite information to each other.

Within a period of three months of the notification referred to in (a), the Member States concerned shall inform the Commission of the result of such contacts and in particular the measures they intend to apply, if any, including the maximum residue level they have agreed. The Member State of origin shall inform the other Member States of the result of such contacts;

▼M6

- (c) the Commission shall immediately refer the matter to the Standing Committee on Plant Health and, if possible, submit a proposal aimed at establishing in Annex II a temporary maximum residue level, which shall be adopted in accordance with the procedure laid down in Article 10a.

In its proposal, the Commission shall take account of existing technical and scientific knowledge on the matter and in particular data submitted by the Member States with an interest, especially the toxicological assessment and estimated ADI, good agricultural practice and the trial data which the Member State of origin used to establish the maximum residue level, together with the reasons given by the Member State of destination for deciding on the measures in question.

The period of validity of the temporary maximum level shall be laid down in the legal act adopted and may not exceed four years. That period may be linked to the supply, by the Member State of origin and/or other Member States with an interest, of the trial data required by the Commission in order to set the maximum residue level in accordance with Article 3 (1). At their request, the Commission and the Member States shall be kept informed regarding the programme of trials established.

4. Any measure provided for in paragraphs 2 or 3 shall be taken by a Member State with due regard for its obligations under the Treaty, in particular Articles 30 to 36 thereof.

5. Council Directive 83/189/EEC of 28 March 1983 laying down a procedure for the provision of information in the field of technical standards and regulations⁽¹⁾ shall not apply to measures adopted and notified by Member States in accordance with paragraph 3 of this Article.

6. Detailed measures for the implementation of the procedure set out in this Article may be adopted in accordance with the procedure laid down in Article 9.

▼B*Article 6*

1. The sampling methods necessary for carrying out the checks provided for in Article 3 on fruit and vegetables shall be those laid down by Commission Directive 79/700/EEC⁽²⁾. The sampling methods necessary for carrying out such checks on products other than fruit and vegetables, and the methods of analysis for all products, shall be determined in accordance with the procedure laid down in Article 9.

⁽¹⁾ OJ No L 109, 26. 4. 1983, p. 8. Directive as last amended by Decision 96/139/EC (OJ No L 32, 10. 2. 1996, p. 31).

⁽²⁾ OJ No L 207, 15. 8. 1979, p. 26.

▼B

The existence of Community methods of analysis shall not preclude Member States from using other tested and scientifically valid methods provided that this does not hinder the free movement of products recognized by virtue of Community methods as complying with the rules provided for in this Directive. In the event of differences in the interpretation of results, those obtained by the use of Community methods shall prevail.

2. The methods of analysis determined under paragraph 1 shall comply with the criteria set out in the Annex to Directive 85/591/EEC.

3. Member States shall inform the other Member States and the Commission of the other methods used pursuant to paragraph 1.

▼M6*Article 7*

Without prejudice to the amendments made to the Annexes in accordance with Articles 5a, 5b (3) and 8, amendments to Annexes I and II as a result of developments in scientific or technical knowledge shall be adopted in accordance with the procedure laid down in Article 10a. In particular, when establishing maximum residue levels, account shall be taken of a relevant dietary intake risk assessment and of the number and quality of the data available.

▼B*Article 8*

1. Where a Member State, as a result of new information or of a reassessment of existing information, considers that a maximum level fixed in the list referred to in Article 1 endangers human or animal health, and therefore requires swift action to be taken, that Member State may temporarily reduce the level in its own territory. In that case, it shall immediately notify the other Member States and the Commission of the measures, attaching a statement of the reasons therefor.

2. The Commission shall quickly examine the grounds given by the Member State referred to in the first subparagraph and shall consult the Member States within the Standing Committee on Plant Health, hereinafter referred to as 'the Standing Committee'; it shall then deliver its opinion forthwith and take the appropriate measures. The Commission shall immediately notify the Council and the Member States of any measures taken. Any Member State may refer the Commission's measures to the Council within 15 days of such notification. The Council acting by a qualified majority may take a different decision within 15 days of the date on which the matter was referred to it.

3. If the Commission considers that the maximum levels laid down in the list referred to in Article 1 should be amended to resolve the difficulties mentioned in paragraph 1 and to ensure the protection of human health, it shall initiate the procedure laid down in Article 10, with a view to adopting those amendments. In this case, the Member State which has taken measures under paragraph 1 may in that event maintain them until the Council or the Commission has taken a decision in accordance with the said procedure.

▼B*Article 9*

1. Where the procedure laid down in this Article is to be followed, the matter shall be referred without delay to the Standing Committee by its chairman, either on his own initiative or at the request of a Member State.
2. The representative of the Commission shall submit to the Standing Committee a draft of the measures to be taken. The Standing Committee shall deliver its opinion on the draft within a time limit which the chairman may lay down according to the urgency of the matter. The opinion shall be delivered by the majority laid down in Article 148 (2) of the Treaty in the case of decisions which the Council is called upon to adopt on a proposal from the Commission. The votes of the representatives of the Member States within the Standing Committee shall be weighted in the manner set out in that Article. The chairman shall not vote.
3. The Commission shall adopt the measures envisaged if they are in accordance with the opinion of the Standing Committee.
4. If the measures are not in accordance with the opinion of the Standing Committee, or if no opinion is delivered, the Commission shall without delay submit to the Council a proposal relating to the measures to be taken. The Council shall act by a qualified majority.
5. If, within three months following the date on which the matter was referred to the Council, the Council has not acted, the proposed measures shall be adopted by the Commission.

Article 10

1. Where the procedure laid down in this Article is to be followed, the matter shall be referred without delay to the Standing Committee by its chairman, either on his own initiative or at the request of a Member State.
2. The representative of the Commission shall submit to the Standing Committee a draft of the measures to be taken. The Standing Committee shall deliver its opinion on the draft within a time limit which the chairman may lay down according to the urgency of the matter. The opinion shall be delivered by the majority laid down in Article 148 (2) of the Treaty in the case of decisions which the Council is called upon to adopt on a proposal from the Commission. The votes of the representatives of the Member States within the Standing Committee shall be weighted in the manner set out in that Article. The chairman shall not vote.
3. The Commission shall adopt the measures envisaged if they are in accordance with the opinion of the Standing Committee.
4. If the measures are not in accordance with the opinion of the Standing Committee, or if no opinion is delivered, the Commission shall without delay submit to the Council a proposal relating to the measures to be taken. The Council shall act by a qualified majority.

▼ B

5. If, within 15 days following the date on which the matter was referred to the Council, the Council has not acted, the proposed measures shall be adopted by the Commission.

▼ M31*Article 10a*

1. The Commission shall be assisted by a committee.
2. Where reference is made to this Article, Articles 5 and 7 of Decision 1999/468/EC ⁽¹⁾ shall apply.
The period laid down in Article 5(6) of Decision 1999/468/EC shall be set at three months.
3. The Committee shall adopt its Rules of Procedure.

▼ M6*Article 10b*

Member States shall bring into force the laws, regulations and administrative provisions necessary to ensure that the amendments in Annex II resulting from decisions referred to in Articles 3 (1) and (2), 5a, 5b (3), 7 and 8 (3) can be implemented in their territory within a maximum period of eight months from their adoption, and within a shorter implementation period when required for urgent reasons of human health protection.

In order to safeguard legitimate expectations, Community legal implementing acts may provide for transitional periods for the implementation of certain maximum residue levels allowing the normal marketing of the harvested products.

▼ B*Article 11*

1. Member States shall take the measures necessary to comply with this Directive not later than 31 December 1992.
2. ‘When Member States adopt the measures referred to in paragraph 1, they shall contain a reference to this Directive or shall be accompanied by such reference on the occasion of their official publication. The methods of making such a reference shall be laid down by the Member States.’

Article 12

This Directive is addressed to the Member States.

⁽¹⁾ OJ L 184, 17.7.1999, p. 23.

▼BANNEX ►M1 I ◀**List of products referred to in Article 1 and the portion of the products to which the maximum residue levels apply**

Note: The word 'fresh' is taken to extend to products which have been chilled or frozen ►M1 and in the case of dried fruit and vegetables, attention is drawn to Article 3 (1) of the Directive ◀

Groups of products	Products included in the groups	Part of product to which maximum residue levels apply
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts		
(i) CITRUS FRUIT	Grapefruit	} Whole product
	Lemons	
	Limes	
	Mandarins (including clem-entines and similar hybrids)	
	Oranges	
	Pomelos	
(ii) TREE (SHELLED OR UNSHELLED) NUTS OR	Almonds	} Whole product after removal of shell
	Brazil nuts	
	Cashew nuts	
	Chestnuts	
	Coconuts	
	Hazelnuts	
	Macadamia nuts	
	Pecans	
	Pine nuts	
	Pistachios	
	Walnuts	
(iii) POME FRUIT	Apples	} Whole product after removal of stems
	Pears	
	Quinces	
(iv) STONE FRUIT	Apricots	} Whole product after removal of stems
	Cherries	
	Peaches (including nectarines and similar hybrids)	
	Plums	

▼ B

Groups of products	Products included in the groups	Part of product to which maximum residue levels apply
(v) BERRIES AND SMALL FRUIT	(a) <i>Table and wine grapes</i> (b) <i>Strawberries</i> (other than wild) (c) <i>Cane fruit</i> (other than wild): Blackberries Loganberries Raspberries (d) <i>Other small fruit and berries</i> (other than wild): Bilberries Cranberries Currants (red, black and white) Gooseberries (e) <i>Wild berries and wild fruit</i>	Whole product after removal of caps and stems (if any) and, in the case of currants, fruits with stems
(vi) MISCELLANEOUS FRUIT	Avocados Bananas Dates Figs Kiwi fruit Kumquats Litchis Mangoes ► M1 ————— ◀	Whole fruit after removal of stems (if any) and in the case of pineapple after removal of the crown
▼ M46	Papaya	
▼ B	Passion fruit Pineapples Pomegranates	
▼ M1	Olives	Whole fruit after removal of stems (if any) after removal of soil (if any) by rinsing in running water

▼ B

Groups of products	Products included in the groups	Part of product to which maximum residue levels apply
2. Vegetables, fresh or uncooked, frozen or dry		
(i) ROOT AND TUBER VEGETABLES	Beetroot Carrots Cassava Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips Yams	Whole product after removal of tops and adhering soil (if any) (removal of soil by rinsing in running water or by gentle brushing of the dry product)
▼ <u>M46</u>		
▼ <u>B</u>		
(ii) BULB VEGETABLES	Garlic Onions Shallots Spring onions	Onions (dry), shallots (dry), garlic (dry): whole product after removal of easily detachable skin and soil (if any). Onions, shallots and garlic other than dry, spring onions: whole product after removal of roots and soil (if any)
(iii) FRUITING VEGETABLES	(a) <i>Solanacea</i> Tomatoes Peppers ► <u>M50</u> Okra ◀ (b) <i>Cucurbits — edible peel</i> Cucumbers Gherkins Courgettes (c) <i>Cucurbits — inedible peel</i> Melons Squashes Watermelons (d) <i>Sweet corn</i>	Whole product after removal of stems Kernels or cobs without husks

▼ B

Groups of products	Products included in the groups	Part of product to which maximum residue levels apply
(iv) BRASSICA VEGETABLES	(a) <i>Flowering brassicas</i> Broccoli Cauliflower	} Cauliflower and broccoli: curd only
	(b) <i>Head brassicas</i> Brussels sprouts Head cabbage	}
	(c) <i>Leafy brassicas</i> Chinese cabbage Kale	} Product after removal of decayed leaves (if any)
	(d) <i>Kohlrabi</i>	Whole product after removal of tops and adhering soil (if any) (removal of soil by rinsing in running water or by gentle brushing of the dry product)
(v) LEAF VEGETABLES AND FRESH HERBS	(a) <i>Lettuce and similar</i> Cress Lamb's lettuce Lettuce Broad-leaf endive ▶ M51 ▶ M62 Leaves and stems of brassica, including turnip greens ◀ Rucola ◀	}
	(b) <i>Spinach and similar</i> Beet leaves (chard)	} Whole product after removal of decayed outer leaves, root and soil (if any)
	(c) <i>Watercress</i>	}
	(d) <i>Witloof</i>	}
	(e) <i>Herbs</i> Chervil Chives Parsley	}
(vi) LEGUME VEGETABLES (FRESH)	Beans	}
	Peas	} Whole product after removal of pods or with pods if they are intended to be eaten

▼ B

Groups of products	Products included in the groups	Part of product to which maximum residue levels apply
(vii) STEM VEGETABLES	Asparagus	} Whole product after removal of decayed tissue and soil (if any); leeks and fennel: whole product after removal of roots and soil (if any)
	Cardoons	
	Celery	
	Fennel	
	Globe artichokes	
	Leeks	
	Rhubarb	
(viii) FUNGI	Mushrooms (other than wild)	} Whole product after removal of soil or growing medium
	Wild mushrooms	

3. Pulses

Beans	} Whole product
Lentils	
Peas	
Lupines	

▼ M53**▼ B****4. Oil seeds**

Linseed	} Whole seed or kernel after removal of shell or husk, when possible
Peanuts	
Poppy seed	
Rape seed	
Sesame seed	
► M1 ◀	
Colza seed	
Soya bean	

▼ M1

Sunflower seed	Whole seed including shell, when present, and whole seed without shell, when shell is absent'
----------------	---

▼ M48

Hemp seed

▼ M66

Pumpkin seed

▼ M1

ANNEX II

▼ M3

PART A

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts	► <u>M70</u> 0,02 (*) ◀										► <u>M25</u> 0,05 (*) ◀
I) CITRUS FRUIT		► <u>M70</u> 0,01 (*) ◀			► <u>M8</u> 2 ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀		► <u>M69</u> 5 ◀		► <u>M25</u> — ◀
Grapefruit											
Lemons			► <u>M70</u> 0,2 ◀	► <u>M8</u> 0,3 ◀						► <u>M63</u> 5 (P) ◀	
Limes											
Mandarines (including clementines and similar hybrids)			► <u>M70</u> 2 ◀	► <u>M8</u> 1 ◀				► <u>M53</u> 0,5 (P) ◀		► <u>M63</u> 1 (P) ◀	
Oranges				► <u>M8</u> 0,5 ◀				► <u>M53</u> 0,5 (P) ◀			
Pomelos											
Others			► <u>M70</u> 0,3 ◀	► <u>M8</u> 0,05 (*) ◀				► <u>M53</u> 0,1 (*) (P) ◀		► <u>M63</u> 0,02 (*) (P) ◀	
II) TREE NUTS (shelled or unshelled)		► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M53</u> 0,1 (*) (P) ◀	► <u>M69</u> 0,05 (*) ◀		► <u>C1</u> — ◀
Almonds											► <u>M25</u> — ◀
Brazil nuts											

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► M69 Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Cashew nuts											
Chestnuts											
Coconuts											
Hazelnuts										► M63 0,2 (P) ◀	
Macadamia											
Pecans											
Pine nuts											
Pistachios											
Walnuts											
Others										► M63 0,02 (*) (P) ◀	► M25 — ◀
III) POME FRUIT		► M70 1 ◀	► M70 0,5 ◀	► M8 0,5 ◀	► M8 1 ◀		► M8 1 (*) ◀	► M53 0,1 (*) (P) ◀	► M69 2 ◀	► M63 5 (P) ◀	► M25 — ◀
Apples						► M69 0,2 ◀					
Pears											
Quinces											
Others						► M69 0,1 ◀					
IV) STONE FRUIT							► M8 0,05 (*) ◀	► M53 0,1 (*) (P) ◀	► M69 0,05 (*) ◀	► M63 3 (P) ◀	► M25 — ◀
Apricots		► M70 1 ◀			► M8 2 ◀		► M8 — ◀				

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Cherries			► <u>M70</u> 0,3 ◀		► <u>M8</u> 1 ◀	► <u>M69</u> 0,2 ◀					
Peaches (including nectarines and similar hybrids)		► <u>M70</u> 1 ◀	► <u>M70</u> 0,2 ◀	► <u>M8</u> 0,5 ◀	► <u>M8</u> 2 ◀		► <u>M8</u> — ◀				
Plums			► <u>M70</u> 0,2 ◀		► <u>M8</u> 1 ◀						
Others		► <u>M70</u> 0,01 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,1 ◀	► <u>M8</u> — ◀				
V) BERRIES AND SMALL FRUIT									► <u>M69</u> 0,05 (*) ◀		
Table and wine grapes			► <u>M70</u> 0,5 ◀	► <u>M8</u> 0,2 ◀	► <u>M8</u> 0,5 ◀	► <u>M69</u> 0,2 ◀	► <u>M8</u> 1 (*) ◀	► <u>M53</u> 0,5 (P) ◀		► <u>M63</u> 10 (P) ◀	► <u>M25</u> — ◀
Table grapes		► <u>M70</u> 1 ◀									
Wine grapes		► <u>M70</u> 3 ◀									
Strawberries (other than wild)		► <u>M70</u> 3 ◀	► <u>M70</u> 0,2 ◀	► <u>M8</u> 0,5 ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,2 ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M53</u> 0,1 (*) (P) ◀		► <u>M63</u> 15 (P) ◀	► <u>M25</u> — ◀
Cane fruit (other than wild)		► <u>M70</u> 0,01 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,5 ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M53</u> 0,1 (*) (P) ◀		► <u>M63</u> 10 (P) ◀	► <u>M25</u> — ◀
Blackberries			► <u>M70</u> 0,5 ◀			► <u>M69</u> 0,5 ◀					
Dewberries											
Loganberries											

▼ M8

▼ M1

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◄	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Raspberries			► <u>M70</u> 0,5 ◄			► <u>M69</u> 0,5 ◄					
Others			► <u>M70</u> 0,05 (*) ◄			► <u>M69</u> 0,05 (*) ◄					
Other small fruit and berries (other than wild)				► <u>M8</u> 0,05 (*) ◄	► <u>M8</u> 0,05 (*) ◄		► <u>M8</u> 0,05 (*) ◄	► <u>M53</u> 0,1 (*) (P) ◄		► <u>M63</u> 10 (P) ◄	► <u>M25</u> — ◄
Bilberries					► <u>M4</u> — ◄						
Bilberries (fruit of species <i>Vaccinium myrtillus</i>)											
Cranberries		► <u>M70</u> 2 ◄									
Currants (red, black and white)		► <u>M70</u> 10 ◄	► <u>M70</u> 1 ◄		► <u>M4</u> — ◄	► <u>M69</u> 0,5 ◄					► <u>M4</u> — ◄
Gooseberries		► <u>M70</u> 10 ◄	► <u>M70</u> 1 ◄		► <u>M4</u> — ◄	► <u>M69</u> 0,2 ◄					► <u>M4</u> — ◄
Others		► <u>M70</u> 0,01 (*) ◄	► <u>M70</u> 0,05 (*) ◄		► <u>M4</u> — ◄	► <u>M69</u> 0,05 (*) ◄					► <u>M4</u> — ◄
Wild berries and wild fruit		► <u>M70</u> 0,01 (*) ◄	► <u>M70</u> 0,05 (*) ◄	► <u>M8</u> 0,05 (*) ◄	► <u>M8</u> 2 ◄	► <u>M69</u> 0,05 (*) ◄	► <u>M8</u> 0,05 (*) ◄	► <u>M53</u> 0,1 (*) (P) ◄		► <u>M63</u> 0,02 (*) (P) ◄	► <u>M25</u> — ◄
VI) MISCELLANEOUS FRUIT				► <u>M8</u> 0,05 (*) ◄	► <u>M8</u> 0,05 (*) ◄		► <u>M8</u> 0,05 (*) ◄				
Avocados											
Bananas		► <u>M70</u> 0,2 ◄	► <u>M70</u> 3 ◄						► <u>M69</u> 2 ◄		

▼ M8▼ M1

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Dates											
Figs											
Kiwi			► <u>M70</u> 2 ◀			► <u>M69</u> 0,2 ◀				► <u>M63</u> 5 (P) ◀	► <u>M25</u> — ◀
Kumquats											
Litchis											
Mangoes											
Olives											► <u>M4</u> — ◀
▼ <u>M8</u>											
Olives (table consumption)						► <u>M69</u> 1 ◀					
Olives (oil extraction)						► <u>M69</u> 1 ◀		► <u>M53</u> 1 (P) ◀			
▼ <u>M66</u>											
Papaya		20									
▼ <u>M1</u>											
Passion fruit											
Pineapples											
Pomegranates											
Other		► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀			► <u>M69</u> 0,05 (*) ◀		► <u>M53</u> 0,1 (*) (P) ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
2. Vegetables, fresh or uncooked, frozen or dry	► <u>M70</u> 0,02 (*) ◀										► <u>M25</u> 0,05 (*) ◀
1) ROOT AND TUBER VEGETABLES				► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M53</u> 0,1 (*) (P) ◀	► <u>M69</u> 0,05 (*) ◀		

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► M69 Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Beetroot											
Carrots		► M70 1 ◀	► M70 0,1 ◀							► M63 0,5 (P) ◀	
Celeriac		► M70 1 ◀									► M25 — ◀
Horseradish										► M63 0,5 (P) ◀	
Jerusalem artichokes											
Parsnips										► M63 0,5 (P) ◀	
Parsley root										► M63 0,5 (P) ◀	
Radishes			► M70 0,2 ◀		► M4 — ◀					► M63 0,3 (P) ◀	► M25 — ◀
Salsify											
Sweet potatoes											
Swedes					► M4 — ◀						
Turnips					► M4 — ◀						► M4 — ◀
Yams											
Others		► M70 0,01 (*) ◀	► M70 0,05 (*) ◀		► M4 — ◀					► M63 0,02 (*) (P) ◀	► M25 — ◀
II) BULB VEGETABLES				► M8 0,05 (*) ◀			► M8 0,05 (*) ◀	► M53 0,1 (*) (P) ◀	► M69 0,05 (*) ◀		► M25 — ◀
Garlic		► M70 0,5 ◀			► M8 0,1 ◀	► M69 0,1 ◀				► M63 0,2 (P) ◀	

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Onions		► <u>M70</u> 0,5 ◀	► <u>M70</u> 0,2 ◀		► <u>M8</u> 0,1 ◀	► <u>M69</u> 0,1 ◀				► <u>M63</u> 0,2 (P) ◀	► <u>M8</u> — ◀
Shallots		► <u>M70</u> 0,5 ◀			► <u>M8</u> 0,1 ◀	► <u>M69</u> 0,1 ◀				► <u>M63</u> 0,2 (P) ◀	► <u>M8</u> — ◀
Springonions		► <u>M70</u> 10 ◀				► <u>M69</u> 0,1 ◀				► <u>M63</u> 3 (P) ◀	► <u>M8</u> — ◀
Others		► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀				► <u>M63</u> 0,02 (*) (P) ◀	► <u>M8</u> — ◀
III) FRUITING VEGETABLES							► <u>M53</u> 0,1 (*) (P) ◀				
Solanacea		► <u>M70</u> 2 ◀	► <u>M70</u> 0,5 ◀	► <u>M8</u> 0,5 ◀	► <u>M8</u> 0,5 ◀					► <u>M63</u> 5 (P) ◀	► <u>M25</u> — ◀
Tomatoes						► <u>M69</u> 0,3 ◀	► <u>M8</u> 1 (*) ◀	► <u>M69</u> 0,5 ◀			
Peppers							► <u>M8</u> 0,2 (*) ◀	► <u>M4</u> — ◀			
Aubergines						► <u>M69</u> 0,3 ◀					
Okra						0,3					
Others						► <u>M69</u> 0,2 ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀			
Cucurbits — edible peel			► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,2 ◀	► <u>M69</u> 0,2 ◀		► <u>M69</u> 0,2 ◀	► <u>M63</u> 2 (P) ◀	► <u>M25</u> — ◀	
Cucumbers		► <u>M70</u> 1 ◀					► <u>M8</u> 0,2 (*) ◀				

▼ M69▼ M1

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Gherkins		► <u>M70</u> 5 ◀									
Courgettes											
Others		► <u>M70</u> 0,01 (*) ◀									
Cucurbits — inedible peel		► <u>M70</u> 1 ◀	► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,2 ◀	► <u>M69</u> 0,2 ◀				► <u>M63</u> 1 (P) ◀	► <u>M25</u> — ◀
Melons											
Squashes											
Watermelons											
Others											
Sweet corn		► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀				
IV) BRASSICA VEGETABLES											
Flowering brassica		► <u>M70</u> 3 ◀	► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,5 ◀	► <u>M69</u> 0,1 ◀	► <u>M8</u> 1 (S) ◀				
Broccoli											► <u>M4</u> — ◀
Cauliflower											► <u>M25</u> — ◀
Others											► <u>M25</u> — ◀

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Head brassica				► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,5 ◀	► <u>M69</u> 0,1 ◀	► <u>M8</u> 0,05 (*) ◀				► <u>M8</u> — ◀
Brussels sprouts		► <u>M70</u> 3 ◀								► <u>M63</u> 0,5 (P) ◀	
Head cabbage		► <u>M70</u> 3 ◀	► <u>M70</u> 1 ◀							► <u>M63</u> 5 (P) ◀	► <u>M25</u> — ◀
Others		► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀							► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
Leafy brassica		► <u>M70</u> 0,01 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 1 ◀	► <u>M69</u> 0,5 ◀					► <u>M25</u> — ◀
Chinese cabbage			► <u>M70</u> 0,5 ◀				► <u>M8</u> 1 (*) ◀			► <u>M63</u> 5 (P) ◀	
Kale							► <u>M8</u> — ◀				
Others			► <u>M70</u> 0,05 (*) ◀				► <u>M8</u> 0,05 (*) ◀			► <u>M63</u> 0,02 (*) (P) ◀	
Kohlrabi		► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,2 ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀			► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
V) LEAF VEGETABLES AND FRESH HERBS			► <u>M70</u> 0,05 (*) ◀					► <u>M53</u> 0,1 (*) (P) ◀	► <u>M69</u> 0,05 (*) ◀		
Lettuce and similar		► <u>M70</u> 0,01 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 2 ◀	► <u>M69</u> 0,5 ◀	► <u>M8</u> 0,05 (*) ◀			► <u>M63</u> 10 (P) ◀	► <u>M25</u> — ◀
Cress											
Lamb's lettuce											

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Lettuce											
Scarole											
Others											
Spinach and similar		► <u>M70</u> 0,01 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,5 ◀	► <u>M69</u> 0,5 ◀	► <u>M8</u> 0,05 (*) ◀			► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
▼ <u>M8</u>											
Spinach											
▼ <u>M1</u>											
Beet leaves (chord)											
▼ <u>M8</u>											
Others											
▼ <u>M1</u>											
Water cress		► <u>M70</u> 0,01 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀			► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
Witloof		► <u>M70</u> 0,01 (*) ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀			► <u>M63</u> 2 (P) ◀	► <u>M25</u> — ◀
Herbs		► <u>M70</u> 5 ◀		► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 2 ◀	► <u>M69</u> 0,5 ◀	► <u>M8</u> 0,05 (*) ◀			► <u>M63</u> 10 (P) ◀	► <u>M25</u> — ◀
Chervil											
Chives											
Parsley											
Celery leaves											
Others											

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
VI) LEGUME VEGETABLES (fresh)			► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀		► <u>M69</u> 0,2 ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M53</u> 0,1 (*) (P) ◀	► <u>M69</u> 0,05 (*) ◀		
Beans (with pods)		► <u>M70</u> 5 ◀			► <u>M8</u> 0,5 ◀					► <u>M63</u> 5 (P) ◀	► <u>M25</u> — ◀
Beans (without pods)		► <u>M70</u> 2 ◀			► <u>M8</u> — ◀						► <u>M8</u> — ◀
Peas (with pods)		► <u>M70</u> 2 ◀			► <u>M8</u> 0,5 ◀					► <u>M63</u> 2 (P) ◀	► <u>M25</u> — ◀
Peas (without pods)		► <u>M70</u> 0,3 ◀			► <u>M8</u> — ◀					► <u>M63</u> 0,3 (P) ◀	► <u>M8</u> — ◀
Others		► <u>M70</u> 0,01 (*) ◀			► <u>M8</u> 0,05 (*) ◀					► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
VII) STEM VEGETABLES				► <u>M8</u> 0,05 (*) ◀			► <u>M8</u> 0,05 (*) ◀	► <u>M53</u> 0,1 (*) (P) ◀	► <u>M69</u> 0,05 (*) ◀		
Asparagus					► <u>M22</u> 0,1 (1) ◀						
Cardoons											
Celery		► <u>M70</u> 10 ◀			► <u>M8</u> — ◀						► <u>M25</u> — ◀
Fennel					► <u>M4</u> — ◀						► <u>M4</u> — ◀
Globe artichokes			► <u>M70</u> 1 ◀	► <u>M8</u> — ◀	► <u>M8</u> 2 ◀	► <u>M69</u> 0,1 ◀					
Leek		► <u>M70</u> 10 ◀	► <u>M70</u> 0,5 ◀		► <u>M8</u> 0,5 ◀	► <u>M69</u> 0,2 ◀					► <u>M25</u> — ◀

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► <u>M69</u> Deltamethrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
Rhubarb										► <u>M63</u> 0,2 (P) ◀	► <u>M25</u> — ◀
Others		► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> — ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 0,05 (*) ◀				► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
VIII) FUNGI			► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀		► <u>M69</u> 0,05 ◀	► <u>M8</u> 0,05 (*) ◀		► <u>M69</u> 0,05 (*) ◀	► <u>M63</u> 0,02 (*) (P) ◀	► <u>M25</u> — ◀
Cultivated mushrooms		► <u>M70</u> 2 ◀			► <u>M8</u> 0,05 (*) ◀			► <u>M53</u> 0,1 (*) (P) ◀			
Wild mushrooms		► <u>M70</u> 0,01 (*) ◀			► <u>M8</u> 1 ◀			► <u>M53</u> 50 (P) ◀			
3. Pulses	► <u>M70</u> 0,02 (*) ◀	► <u>M70</u> 0,01 (*) ◀	► <u>M70</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M8</u> 0,05 (*) ◀	► <u>M69</u> 1 ◀	► <u>M8</u> 0,05 (*) ◀		► <u>M69</u> 0,05 (*) ◀	► <u>M63</u> 0,2 (P) ◀	► <u>M25</u> 0,05 (*) ◀
Beans					► <u>M8</u> — ◀			► <u>M53</u> 2 (P) ◀			
Lentils					► <u>M8</u> — ◀						
Peas					► <u>M8</u> — ◀			► <u>M53</u> 10 (P) ◀			
▼ <u>M53</u>								10 (P)			
Lupins											
▼ <u>M1</u>					► <u>M8</u> — ◀			► <u>M53</u> 0,1 (*) (P) ◀			
Others											

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► M69 Deltamethrin (cis-deltamethrin) (*) ◄	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
4. Oil seeds			► M70 0,05 (*) ◄	► M8 0,05 (*) ◄			► M8 0,1 (*) ◄		► M69 0,05 (*) ◄		► M25 0,05 ◄
Linseed					► M8 0,2 ◄			► M53 10 (P) ◄		► M63 0,5 (P) ◄	
Peanuts		► M70 0,05 ◄									► M25 — ◄
Poppy seed					► M8 0,2 ◄						
Sesame seed					► M8 0,2 ◄						
Sunflower seed					► M8 0,2 ◄			► M53 20 (P) ◄		► M63 0,5 (P) ◄	► M4 — ◄
Rape seed				► M8 — ◄	► M8 0,2 ◄	► M69 0,1 ◄		► M53 10 (P) ◄		► M63 0,5 (P) ◄	► M25 — ◄
Soya bean	► M70 0,3 ◄							► M53 20 (P) ◄			
Mustard						► M69 0,1 ◄		► M53 10 (P) ◄			► M25 — ◄
Cotton seed					► M8 0,2 ◄			► M53 10 (P) ◄			► M25 — ◄
Others	► M70 0,05 (*) ◄	► M70 0,01 (*) ◄		► M8 — ◄	► M8 0,05 (*) ◄	► M69 0,05 (*) ◄		► M53 0,1 (*) (P) ◄		► M63 0,02 (*) (P) ◄	► M25 — ◄
5. Potatoes	► M70 0,02 (*) ◄	► M70 0,01 (*) ◄	► M70 0,05 (*) ◄	► M8 0,05 (*) ◄	► M8 0,05 (*) ◄	► M69 0,05 (*) ◄	► M8 0,05 (*) ◄	► M53 0,5 (P) ◄	► M69 3 ◄	► M63 0,02 (*) (P) ◄	► M25 0,05 (*) ◄
Early potatoes											
Ware potatoes											

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)										
	Acephate	Chlorothalonil	Chlorpyrifos	Chlorpyrifos-methyl	Cypermethrin, including other mixtures of constituent isomers (sum of isomers)	► M69 Delta-methrin (cis-deltamethrin) (*) ◀	Fenvalerate, including other mixtures of constituent isomers (sum of isomers)	Glyphosate	Imazalil	Iprodione	Permethrin (sum of isomers)
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	► M70 0,05 (*) ◀	► M70 0,1 (*) ◀	► M70 0,10 (*) ◀	► M8 0,1 (*) ◀	► M8 0,5 ◀	► M69 5 ◀	► M8 10 (*) ◀	► M53 2 (P) ◀	► M69 0,1 (*) ◀	► M63 0,1 (*) (P) ◀	► M25 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► M70 0,05 (*) ◀	► M70 50 ◀	► M70 0,10 (*) ◀	► M8 0,1 (*) ◀	► M8 30 ◀	► M69 5 ◀	► M8 5 (*) ◀	► M53 0,1 (*) (P) ◀	► M69 0,1 (*) ◀	► M63 0,1 (*) (P) ◀	► M25 0,1 (*) ◀

(*) Indicates lower limit of analytical determination.

► **M66** (*) Temporary MRLs valid until 1 November 2008, pending review of the Annex III dossier under Directive 91/414/EEC and re-registration of deltamethrin formulations at Member State level. ◀

(P) ► **M53** Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC. ◀

(a) (b) (c) (d) Should levels not be adopted by ► **M7** 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*)

► **M8** ► **C3** x Should this level not be confirmed or amended by a directive, with effect from 1 July 2000, the appropriate lower limit of analytical determination shall apply. ◀ ◀

► **M22** (1) Pending trials for the second season, results to be submitted before 1 September 2002. ◀

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► <u>M70</u> Carben-dazim and benomyl (sum of carben-dazim expressed as carben-dazim) ◀	► <u>M70</u> Dithioc-arbamates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram (1), (2) ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5-dicloroaniline moiety, expressed as vinclozolin)	► <u>M70</u> Thioph-anate-methyl ◀
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts						
(I) CITRUS FRUIT	► <u>M70</u> 0,5 ◀	► <u>M70</u> 5 (mz) ◀	► <u>M54</u> 0,01 (*) ◀	► <u>M8</u> 0,02 (*) ◀	0,05 (*)	► <u>M70</u> 0,1 (*) ◀
Grapefruit						
Lemons						
Limes						
Mandarines (including clem-entines and similar hybrids)						
Oranges						
Pomelos						
Others						
(II) TREE NUTS (shelled or unshelled)	► <u>M70</u> 0,1 (*) ◀		► <u>M54</u> 0,01 (*) ◀	► <u>M8</u> 0,05 (*) ◀	0,05 (*)	► <u>M70</u> 0,2 ◀
Almonds						
Brazil nuts						
Cashew nuts						
Chestnuts						
Coconuts						
Hazelnuts						
Macadamia						
Pecans						
Pine nuts						
Pistachios						

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► M70 Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► M70 Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram (1), (2) ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► M70 Thioph-anate-methyl ◀
Walnuts		► M70 0,1 (mz) ◀				
Others		► M70 0,05 (*) ◀				
(III) POME FRUIT	► M70 0,2 ◀	► M70 5 (ma, mz, me, pr, t, z) ◀	► M54 0,01 (*) ◀		1	► M70 0,5 ◀
Apples				► M8 ◀		
Pears				► M8 1 ◀		
Quinces				► M8 0,02 (*) ◀		
Others				► M8 ◀		
(IV) STONE FRUIT				► M8 ◀		
Apricots	► M70 0,2 ◀	► M70 2 (mz, t) ◀	► M54 0,1 ◀		2	► M70 2 ◀
Cherries	► M70 0,5 ◀	► M70 2 (mz, me, pr, t, z) ◀		► M8 0,02 (*) ◀	0,5	► M70 0,3 ◀
Peaches (including nectarines and similar hybrids)	► M70 0,2 ◀	► M70 2 (mz, t) ◀	► M54 0,05 ◀		► M14 0,05 (*) ◀	► M70 2 ◀
Plums	► M70 0,5 ◀	► M70 2 (mz, me, t, z) ◀			(c)	► M70 0,3 ◀
Others	► M70 0,1 (*) ◀	► M70 0,05 (*) ◀	► M54 0,01 (*) ◀	► M8 2 ◀	0,05 (*)	► M70 0,1 (*) ◀

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► M70 Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► M70 Dithiocarbamates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram (1), (2) ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dicloroaniline moiety, expressed as vinclozolin)	► M70 Thiophanate-methyl ◀
(V) BERRIES AND SMALL FRUIT			► M54 0,01 (*) ◀			
Table and wine grapes		► M70 5 (ma, mz, me, pr, t) ◀		► M8 5 ◀	5	
Table grapes	► M70 0,3 ◀					► M70 0,1 (*) ◀
Wine grapes	► M70 0,5 ◀					► M70 3 ◀
Strawberries (other than wild)	► M70 0,1 (*) ◀	► M70 10 (t) ◀		► M8 5 ◀	5	► M70 0,1 (*) ◀
Cane fruit (other than wild)	► M70 0,1 (*) ◀	► M70 0,05 (*) ◀			5	► M70 0,1 (*) ◀
Blackberries						
Dewberries						
Loganberries						
Raspberries				► M8 10 ◀		
Others				► M8 0,02 (*) ◀		
Other small fruit and berries (other than wild)	► M70 0,1 (*) ◀	► M70 5 (mz) ◀		► M8 0,02 (*) ◀		► M70 0,1 (*) ◀
Bilberries (fruit of species <i>vaccinium myrtilus</i>)						
Cranberries						
Currants (red, black and white)		► M70 5 (mz) ◀			(c)	

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► <u>M70</u> Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► <u>M70</u> Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram ⁽¹⁾ , ⁽²⁾ ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► <u>M70</u> Thioph-anate-methyl ◀
Gooseberries					0,05 (*)	
Others		► <u>M70</u> 0,05 (*) ◀			0,05 (*)	
Wild berries and wild fruit	► <u>M70</u> 0,1 (*) ◀	► <u>M70</u> 0,05 (*) ◀		► <u>M8</u> 0,02 (*) ◀	0,05 (*)	► <u>M70</u> 0,1 (*) ◀
(VI) MISCELLANEOUS FRUIT			► <u>M54</u> 0,01 (*) ◀		0,05 (*) (except kiwi)	
Avocados						
Bananas		► <u>M70</u> 2 (mz, me) ◀				
Dates						
Figs						
Kiwi				► <u>M8</u> 5 ◀		
Kumquats						
Litchis						
Mangoes	► <u>M70</u> 0,5 ◀	► <u>M70</u> 2 (mz) ◀				► <u>M70</u> 1 ◀
Olives						
Olives (table consumption)		► <u>M70</u> 5 (mz, pr) ◀				
Olives (oil extraction)		► <u>M70</u> 5 (mz, pr) ◀				

▼ M8▼ M1

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► <u>M70</u> Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► <u>M70</u> Dithiocarbamates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram ⁽¹⁾ , ⁽²⁾ ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► <u>M70</u> Thiophanate-methyl ◀
Papaya	► <u>M70</u> 0,2 ◀	► <u>M70</u> 7 (mz) ◀				► <u>M70</u> 1 ◀
Passion fruit						
Pineapples						
Pomegranates						
Others	► <u>M70</u> 0,1 (*) ◀	► <u>M70</u> 0,05 (*) ◀		► <u>M8</u> 0,02 (*) ◀		► <u>M70</u> 0,1 (*) ◀
2. Vegetables, fresh or uncooked, frozen or dry						
(I) ROOT AND TUBER VEGETABLES	► <u>M70</u> 0,1 (*) ◀		► <u>M54</u> 0,01 (*) ◀	► <u>M8</u> 0,02 (*) ◀		► <u>M70</u> 0,1 (*) ◀
Beetroot		► <u>M70</u> 0,5 (mz) ◀				
Carrots		► <u>M70</u> 0,2 (mz) ◀			(c)	
Celeriac		► <u>M70</u> 0,3 (ma, me, pr, t) ◀				
Horseradish		► <u>M70</u> 0,2 (mz) ◀			(c)	
Jerusalem artichokes						
Parsnips		► <u>M70</u> 0,2 (mz) ◀				
Parsley root		► <u>M70</u> 0,2 (mz) ◀				

▼ M67▼ M1

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► <u>M70</u> Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► <u>M70</u> Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram ⁽¹⁾ , ⁽²⁾ ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► <u>M70</u> Thioph-anate-methyl ◀
Radishes					(c)	
Salsify		► <u>M70</u> 0,2 (mz) ◀				
Sweet potatoes						
Swedes					(c)	
Turnips						
Yams						
Others		► <u>M70</u> 0,05 (*) ◀			0,05 (*)	
(II) BULB VEGETABLES	► <u>M70</u> 0,1 (*) ◀		► <u>M54</u> 0,01 (*) ◀		1	► <u>M70</u> 0,1 (*) ◀
Garlic		► <u>M70</u> 0,1 (mz) ◀		► <u>M8</u> 0,2 ◀		
Onions		► <u>M70</u> 1 (ma, mz) ◀		► <u>M8</u> 0,2 ◀		
Shallots		► <u>M70</u> 1 (ma, mz) ◀		► <u>M8</u> 0,2 ◀		
Spring onions		► <u>M70</u> 1 (mz) ◀		► <u>M4</u> — ◀		
Others		► <u>M70</u> 0,05 (*) ◀		► <u>M8</u> 0,02 (*) ◀		
(III) FRUITING VEGETABLES			► <u>M54</u> 0,01 (*) ◀			
Solanacea				► <u>M8</u> 2 ◀	3	
Tomatoes	► <u>M70</u> 0,5 ◀	► <u>M70</u> 3 (mz, me, pr) ◀			► <u>M14</u> 0,05 (*) ◀	► <u>M70</u> 2 ◀

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► <u>M70</u> Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► <u>M70</u> Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram ⁽¹⁾ , ⁽²⁾ ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► <u>M70</u> Thioph-anate-methyl ◀
Peppers		► <u>M70</u> 5 (mz, pr) ◀				
Aubergines	► <u>M70</u> 0,5 ◀	► <u>M70</u> 3 (mz, me) ◀				► <u>M70</u> 2 ◀
Okra	► <u>M70</u> 2 ◀	► <u>M70</u> 0,5 (mz) ◀				► <u>M70</u> 1 ◀
Others	► <u>M70</u> 0,1 (*) ◀	► <u>M70</u> 0,05 (*) ◀				► <u>M70</u> 0,1 (*) ◀
Cucurbits — edible peel	► <u>M70</u> 0,1 (*) ◀	► <u>M70</u> 2 (mz, pr) ◀		► <u>M8</u> 1 ◀	1	► <u>M70</u> 0,1 (*) ◀
Cucumbers						
Gherkins						
Courgettes						
Others						
Cucurbits — inedible peel	► <u>M70</u> 0,1 (*) ◀	► <u>M70</u> 1 (mz, pr) ◀		► <u>M8</u> 1 ◀	1	► <u>M70</u> 0,3 ◀
Melons						
Squashes						
Watermelons						
Others						
Sweet corn	► <u>M70</u> 0,1 (*) ◀	► <u>M70</u> 0,05 (*) ◀		► <u>M8</u> 0,02 (*) ◀	0,05 (*)	► <u>M70</u> 0,1 (*) ◀

▼ M67▼ M1

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► <u>M70</u> Carben-dazim and benomyl (sum of carben-dazim and benomyl expressed as carben-dazim) ◄	► <u>M70</u> Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram (1), (2) ◄	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5-dichloroaniline moiety, expressed as vinclozolin)	► <u>M70</u> Thioph-anate-methyl ◄
(IV) BRASSICA VEGETABLES						
Flowering brassica	► <u>M70</u> 0,1 (*) ◄	► <u>M70</u> 1 (mz) ◄	► <u>M54</u> 0,02 ◄	► <u>M8</u> 0,02 (*) ◄	0,05 (*)	► <u>M70</u> 0,1 (*) ◄
Broccoli						
Cauliflower						
Others						
Head brassica			► <u>M54</u> 0,01 (*) ◄	► <u>M8</u> 0,02 (*) ◄	0,05 (*)	
Brussels sprouts	► <u>M70</u> 0,5 ◄	► <u>M70</u> 2 (mz) ◄				► <u>M70</u> 1 ◄
Head cabbage		► <u>M70</u> 3 (mz) ◄				
Others	► <u>M70</u> 0,1 (*) ◄	► <u>M70</u> 0,05 (*) ◄				► <u>M70</u> 0,1 (*) ◄
Leafy brassica	► <u>M70</u> 0,1 (*) ◄	► <u>M70</u> 0,5 (mz) ◄	► <u>M54</u> 0,01 (*) ◄	► <u>M8</u> 0,02 (*) ◄		► <u>M70</u> 0,1 (*) ◄
Chinese cabbage					2	
Kale						
Others					0,05 (*)	
Kohlrabi	► <u>M70</u> 0,1 (*) ◄	► <u>M70</u> 1 (mz) ◄	► <u>M54</u> 0,01 (*) ◄	► <u>M8</u> 0,02 (*) ◄	0,05 (*)	► <u>M70</u> 0,1 (*) ◄
(V) LEAF VEGETABLES AND FRESH HERBS	► <u>M70</u> 0,1 (*) ◄		► <u>M54</u> 0,01 (*) ◄			► <u>M70</u> 0,1 (*) ◄
Lettuce and similar		► <u>M70</u> 5 (mz, me, t) ◄		► <u>M8</u> 5 ◄	5	
Cress						

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► <u>M70</u> Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► <u>M70</u> Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram ⁽¹⁾ , ⁽²⁾ ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► <u>M70</u> Thioph-anate-methyl ◀
Lamb's lettuce						
Lettuce						
Scarole						
Others						
Spinach and similar		► <u>M70</u> 0,05 (*) ◀		► <u>M8</u> 0,02 (*) ◀	0,05 (*)	
Spinach						
Beet leaves (chord)						
Others						
Water cress		► <u>M70</u> 0,3 (mz) ◀		► <u>M8</u> 0,02 (*) ◀	0,05 (*)	
Witloof		► <u>M70</u> 0,5 (mz) ◀		► <u>M8</u> 2 ◀	2	
Herbs		► <u>M70</u> 5 (mz, me) ◀		► <u>M8</u> 0,02 (*) ◀	0,05 (*)	
Chervil						
Chives						
Parsley						
Celery leaves						
Others						

▼ M8▼ M1▼ M8▼ M1

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► M70 Carben-dazim and benomyl and carbendazim expressed as carbendazim ◀	► M70 Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram ⁽¹⁾ , ⁽²⁾ ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► M70 Thioph-anate-methyl ◀
(VI) LEGUME VEGETABLES (fresh)						► M70 0,1 (*) ◀
Beans (with pods)	► M70 0,2 ◀	► M70 1 (mz) ◀	► M54 0,5 ◀	► M8 2 ◀	2	
Beans (without pods)		► M70 0,1 (mz) ◀		► M8 ◀	(c)	
Peas (with pods)	► M70 0,2 ◀	► M70 1 (ma, mz) ◀	► M54 0,5 ◀	► M8 1 ◀	2	
Peas (without pods)		► M70 0,1 (mz) ◀		► M8 0,3 ◀	(c)	
Others	► M70 0,1 (*) ◀	► M70 0,05 (*) ◀	► M54 0,01 (*) ◀	► M8 0,02 (*) ◀	0,05 (*)	
(VII) STEM VEGETABLES	► M70 0,1 (*) ◀			► M8 0,02 (*) ◀		► M70 0,1 (*) ◀
Asparagus		► M70 0,5 (mz) ◀				
Cardoons						
Celery					(c)	
Fennel						
Globe artichokes			► M54 0,1 ◀			
Leek		► M70 3 (ma, mz) ◀				
Rhubarb		► M70 0,5 (mz) ◀				
Others		► M70 0,05 (*) ◀	► M54 0,01 (*) ◀		0,05 (*)	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► M70 Carben-dazim and benomyl (sum of carben-dazim and benomyl expressed as carben-dazim) ◀	► M70 Dithioc-aramates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram (1), (2) ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolins and all metabolites containing the 3,5 dicloroaniline moiety, expressed as vinclozolin)	► M70 Thioph-anate-methyl ◀
(VIII) FUNGI		► M70 0,05 (*) ◀	► M54 0,01 (*) ◀	► M8 0,02 (*) ◀	0,05 (*)	► M70 0,1 (*) ◀
Cultivated mushrooms	► M70 1 ◀					
Wild mushrooms	► M70 0,1 (*) ◀					
3. Pulses	► M70 0,1 (*) ◀		► M54 0,01 (*) ◀	► M8 — ◀	(c)	► M70 0,1 (*) ◀
Beans		► M70 0,1 (mz) ◀				
Lentils						
Peas		► M70 0,1 (mz) ◀		► M8 0,2 ◀		
Others		► M70 0,05 (*) ◀		► M8 0,02 (*) ◀		
4. Oil seeds						
Linseed						
Peanuts						
Poppy seed						
Sesame seed						
Sunflower seed (with shell)				► M8 1 ◀		
Sunflower seed (without shell)						
Rape seed		► M70 0,5 (ma, mz) ◀		► M8 1 ◀	1	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)					
	► M70 Carben-dazim and benomyl (sum of benomyl and carbendazim expressed as carbendazim) ◀	► M70 Dithiocarbamates, expressed as CS ₂ , including maneb, mancozeb, metiram, propineb, thiram and ziram ⁽¹⁾ , ⁽²⁾ ◀	Methamidophos	Procymidone	Vinclozolin (sum of vinclozolin and all metabolites containing the 3,5 dichloroaniline moiety, expressed as vinclozolin)	► M70 Thiophanate-methyl ◀
Soya bean	► M70 0,2 ◀		► M54 0,2 ◀	► M8 1 ◀		► M70 0,3 ◀
Mustard						
Cotton seed			► M54 0,2 ◀			
Others	► M70 0,1 (*) ◀	► M70 0,1 (*) ◀	► M54 0,01 (*) ◀	► M8 0,05 (*) ◀	0,05 (*)	► M70 0,1 (*) ◀
5. Potatoes	► M70 0,1 (*) ◀	► M70 0,3 (ma, mz, me, pr) ◀	► M54 0,01 (*) ◀	► M8 0,02 (*) ◀	0,05 (*)	► M70 0,1 (*) ◀
Early potatoes						
Ware potatoes						
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	► M70 0,1 (*) ◀	► M70 0,1 (*) ◀	► M54 0,02 (*) ◀	► M8 0,1 (*) ◀	0,1 (*)	► M70 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► M70 0,1 (*) ◀	► M70 25 (pr) ◀	► M54 0,02 (*) ◀	► M8 0,1 (*) ◀	40	► M70 0,1 (*) ◀

► **M70** ⁽¹⁾ The MRLs expressed as CS₂ can arise from different dithiocarbamates and therefore they do not reflect a single Good Agricultural Practice (GAP). It is therefore not appropriate to use these MRLs to check compliance with a GAP.

⁽²⁾ In brackets the origin of the residue (ma: maneb; me: metiram; mz: mancozeb; pr: propineb; t: thiram; z: ziram). ◀

(*) Indicates lower limit of analytical determination.

(a) (b) (c) (d) Should levels not be adopted by ► **M7** 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*)

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	DDT (sum of p,p'-DDT, o,p'-DDT, p,p'-DDE and p,p'-TDE (DDD) expressed as DDT)
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts	
(I) CITRUS FRUIT	0,05 (*)
Grapefruit	
Lemons	
Limes	
Mandarines (including clementines and similar hybrids)	
Oranges	
Pomelos	
Others	
(II) TREE NUTS (shelled or unshelled)	0,05 (*)
Almonds	
Brazil nuts	
Cashew nuts	
Chestnuts	
Coconuts	
Hazelnuts	
Macadamia	
Pecans	
Pine nuts	
Pistachios	
Walnuts	
Others	
(III) POME FRUIT	0,05 (*)
Apples	
Pears	
Quinces	
Others	
(IV) STONE FRUIT	0,05 (*)
Apricots	
Cherries	
Peaches (including nectarines and similar hybrids)	
Plums	
Others	
(V) BERRIES AND SMALL FRUIT	0,05 (*)
Table and wine grapes	
Table grapes	
Wine grapes	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	DDT (sum of p,p'-DDT, o,p'-DDT, p,p'-DDE and p,p'-TDE (DDD) expressed as DDT)
Strawberries (other than wild)	
Cane fruit (other than wild)	
Blackberries	
Dewberries	
Loganberries	
Raspberries	
Others	
Other small fruit and berries (other than wild)	
Bilberries (fruit of species <i>vaccinium myrtyllus</i>)	
Cranberries	
Currants (red, black and white)	
Gooseberries	
Others	
Wild berries and wild fruit	
(VI) MISCELLANEOUS FRUIT	0,05 (*)
Avocados	
Bananas	
Dates	
Figs	
Kiwi	
Kumquats	
Litchis	
Mangoes	
Olives (table consumption)	
Olives (oil extraction)	
Passion fruit	
Pineapples	
Pomegranates	
Others	
2. Vegetables, fresh or uncooked, frozen or dry	
(I) ROOT AND TUBER VEGETABLES	0,05 (*)
Beetroot	
Carrots	
Celeriac	
Horseradish	
Jerusalem artichokes	
Parsnips	
Parsley root	
Radishes	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	DDT (sum of p,p'-DDT, o,p'-DDT, p,p'-DDE and p,p'-TDE (DDD) expressed as DDT)
Salsify	
Sweet potatoes	
Swedes	
Turnips	
Yams	
Others	
(II) BULB VEGETABLES	0,05 (*)
Garlic	
Onions	
Shallots	
Spring onions	
Others	
(III) FRUITING VEGETABLES	0,05 (*)
Solanacea	
Tomatoes	
Peppers	
Aubergines	
Others	
Cucurbits — edible peel	
Cucumbers	
Gherkins	
Courgettes	
Others	
Cucurbits — inedible peel	
Melons	
Squashes	
Watermelons	
Others	
Sweet corn	
(IV) BRASSICA VEGETABLES	0,05 (*)
Flowering brassica	
Broccoli	
Cauliflower	
Others	
Head brassica	
Brussels sprouts	
Head cabbage	
Others	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	DDT (sum of p,p'-DDT, o,p'-DDT, p,p'-DDE and p,p'-TDE (DDD) expressed as DDT)
Leafy brassica	
Chinese cabbage	
Kale	
Others	
Kohlrabi	
(V) LEAF VEGETABLES AND FRESH HERBS	0,05 (*)
Lettuce and similar	
Cress	
Lamb's lettuce	
Lettuce	
Scarole	
Others	
Spinach and similar	
Beet leaves (chord)	
Water cress	
Witloof	
Herbs	
Chervil	
Chives	
Parsley	
Celery leaves	
Others	
(VI) LEGUME VEGETABLES (fresh)	0,05 (*)
Beans (with pods)	
Beans (without pods)	
Peas (with pods)	
Peas (without pods)	
Others	
(VII) STEM VEGETABLES	0,05 (*)
Asparagus	
Cardoons	
Celery	
Fennel	
Globe artichokes	
Leek	
Rhubarb	
Others	
(VIII) FUNGI	0,05 (*)
Cultivated mushrooms	
Wild mushrooms	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	DDT (sum of p,p'-DDT, o,p'-DDT, p,p'-DDE and p,p'-TDE (DDD) expressed as DDT)
3. Pulses	0,05 (*)
Beans	
Lentils	
Peas	
Others	
4. Oil seeds	0,05 (*)
Linseed	
Peanuts	
Poppy seed	
Sesame seed	
Sunflower seed (with shell)	
Sunflower seed (without shell)	
Rape seed	
Soya bean	
Mustard	
Cotton seed	
Others	
5. Potatoes	0,05 (*)
Early potatoes	
Ware potatoes	
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,2 ► C1 ————— ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (*)

(*) Indicates lower limit of analytical determination.

(a) (b) (c) (d) Should levels not be adopted by ► **M7** 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*).

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts		► M54 0,05 (*) ◀					► M12 0,05 (*) ◀			► M12 0,05 (*) ◀
(I) CITRUS FRUIT	0,05 (*)		0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Grapefruit										
Lemons										
Limes										
Mandarines (including clementines and similar hybrids)										
Oranges										
Pomelos										
Others										
(II) TREE NUTS (shelled or unshelled)	0,05 (*)		0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Almonds										
Brazil nuts										
Cashew nuts										
Chestnuts										
Coconuts										
Hazelnuts										
Macadamia										
Pecans										
Pine nuts										
Pistachios										

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop- hosehyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin- alphos ◀	► M12 Fenvalerate and esfen- valerate ◀		► M12 Mecar- bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
Walnuts										
Others										
(III) POME FRUIT	0,05 (*)		0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,05 ◀	► M12 0,02 (*) ◀	
Apples										
Pears										
Quinces										
Others										
(IV) STONE FRUIT	0,05 (*)		0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)			► M12 0,02 (*) ◀	
Apricots								► M70 0,1 ◀		
Cherries										
Peaches (including nectarines and similar hybrids)								► M70 0,1 ◀		
Plums										
Others								► M70 0,02 (*) ◀		
(V) BERRIES AND SMALL FRUIT	0,05 (*)		0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)				
Table and wine grapes								► M70 0,1 ◀	► M12 0,02 (*) ◀	
Table grapes										
Wine grapes										
Strawberries (other than wild)								► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
Cane fruit								► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Blackberries										
Dewberries										
Loganberries										
Raspberries										
Others										
Other small fruit and berries (other than wild)								► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Bilberries (fruit of species <i>vaccinium myrtyllus</i>)										
Cranberries										
Currants (red, black and white)										
Gooseberries										
Others										
Wild berries and wild fruit									► M12 0,02 (*) ◀	
(VI) MISCELLANEOUS FRUIT	0,05 (*)		0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Avocados										
Bananas										
Dates										
Figs										
Kiwi										
Kumquats										

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
Litchis										
Mangoes										
Olives										
Passion fruit										
Pineapples										
Pomegranates										
Others										
2. Vegetables, fresh or uncooked, frozen or dry							► M12 0,05 (*) ◀			
(I) ROOT AND TUBER VEGETABLES	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Beetroot										
Carrots										
Celeriac										
Horseradish										
Jerusalem artichokes										
Parsnips										
Parsley root										
Radishes										
Salsify										
Sweet potatoes										
Swedes										
Turnips										

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
Yams										
Others										
(II) BULB VEGETABLES	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Garlic										
Onions										
Shallots										
Spring onions										
Others										
(III) FRUITING VEGETABLES	0,05 (*)		0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)				
Solanacea		► M54 0,05 (*) ◀								
Tomatoes								► M70 0,05 ◀	► M12 0,02 (*) ◀	
Peppers										
Aubergines								► M70 0,02 (*) ◀		
Others								► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Cucurbits — edible peel		► M54 0,05 (*) ◀						► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Cucumbers										
Gherkins										
Courgettes										
Others										
Cucurbits — inedible peel		► M54 0,05 (*) ◀						► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
Melons										
Squashes										
Watermelons										
Others										
Sweet corn		► M54 0,1 ◀						► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
(IV) BRASSICA VEGETABLES	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)				
Flowering brassica								► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Broccoli										
Cauliflower										
Others										
Head brassica									► M12 0,02 (*) ◀	
Brussels sprouts								► M70 0,05 ◀		
Head cabbage								► M70 0,1 ◀		
Others								► M70 0,02 (*) ◀		
Leafy brassica								► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Chinese cabbage										
Kale										
Others										
Kohlrabi								► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
(V) LEAF VEGETABLES AND FRESH HERBS	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Lettuce and similar										
Cress										
Lamb's lettuce										
Lettuce										
Scarole										
Others										
Spinach and similar										
Beet leaves (chord)										
Water cress										
Witloof										
Herbs										
Chervil										
Chives										
Parsley										
Celery leaves										
Others										
(VI) LEGUME VEGETABLES (fresh)	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)			► M12 0,02 (*) ◀	
Beans (with pods)										
Beans (without pods)										
Peas (with pods)								► M70 0,1 ◀		

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
Peas (without pods)								► M70 0,02 (*) ◀		
Others										
(VII) STEM VEGETABLES	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)		► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Asparagus										
Cardoons										
Celery										
Fennel										
Globe artichokes										
Leek										
Rhubarb										
Others								► M70 0,02 (*) ◀		
(VIII) FUNGI	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)			► M12 0,02 (*) ◀	
Cultivated mushrooms										
Wild mushrooms								► M70 0,02 (*) ◀		
3. Pulses	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)	► M12 0,05 (*) ◀	► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Beans										
Lentils										
Peas										
Others										

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)									
	Aminotriazole (Amitrole)	Atrazine	Binapacryl	Bromop-hoethyl	Captafol	Dichlorprop (including dichlorprop P)	► M12 Quin-alphos ◀	► M12 Fenvalerate and esfen-valerate ◀		► M12 Mecar-bam ◀
								sum of RR and SS isomers	sum of RS and SR isomers	
4. Oil seeds	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)	► M12 0,05 (*) ◀	► M70 0,05 (*) ◀	► M12 0,05 (*) ◀	
Linseed										
Peanuts										
Poppy seed										
Sesame seed										
Sunflower seed										
Rape seed										
Soya bean										
Mustard										
Cotton seed										
Others										
5. Potatoes	0,05 (*)	► M54 0,05 (*) ◀	0,05 (*)	0,05 (*)	0,02 (*)	0,05 (*)	► M12 0,05 (*) ◀	► M70 0,02 (*) ◀	► M12 0,02 (*) ◀	
Early potatoes										
Ware potatoes										
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,1 (*)	► M54 0,1 (*) ◀	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	► M12 0,1 (*) ◀	► M70 0,05 (*) ◀	► M12 0,05 (*) ◀	
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)	► M54 0,1 (*) ◀	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	► M12 0,1 (*) ◀	► M70 0,05 (*) ◀	► M12 0,05 (*) ◀	

(*) Indicates lower limit of analytical determination.

(a) (b) (c) (d) Should levels not be adopted by ► **M7** 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*)

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts					
(I) CITRUS FRUIT	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Grapefruit					
Lemons					
Limes					
Mandarines (including clementines and similar hybrids)					
Oranges					
Pomelos					
Others					
(II) TREE NUTS (shelled or unshelled)	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Almonds					
Brazil nuts					
Cashew nuts					
Chestnuts					
Coconuts					
Hazelnuts					
Macadamia					
Pecans					
Pine nuts					
Pistachios					
Walnuts					
Others					

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
(III) POME FRUIT	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Apples					
Pears					
Quinces					
Others					
(IV) STONE FRUIT	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Apricots					
Cherries					
Peaches (including nectarines and similar hybrids)					
Plums					
Others					
(V) BERRIES AND SMALL FRUIT	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Table and wine grapes					
Table grapes					
Wine grapes					
Strawberries (other than wild)					
Cane fruit					
Blackberries					
Dewberries					
Loganberries					
Raspberries					
Others					

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
Other small fruit and berries (other than wild)					
Bilberries (fruit of species <i>vaccinium myrtillus</i>)					
Cranberries					
Currants (red, black and white)					
Gooseberries					
Others					
Wild berries and wild fruit					
(VI) MISCELLANEOUS FRUIT	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Avocados					
Bananas					
Dates					
Figs					
Kiwis					
Kumquats					
Litchis					
Mangoes					
Olives					
Passion fruit					
Pineapples					
Pomegranates					
Others					

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
2. Vegetables, fresh or uncooked, frozen or dry					
(I) ROOT AND TUBER VEGETABLES	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Beetroot					
Carrots					
Celeriac					
Horseradish					
Jerusalem artichokes					
Parsnips					
Parsley root					
Radishes					
Salsify					
Sweet potatoes					
Swedes					
Turnips					
Yams					
Others					
(II) BULB VEGETABLES	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Garlic					
Onions					
Shallots					
Spring onions					
Others					

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
(III) FRUITING VEGETABLES	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Solanacea					
Tomatoes					
Peppers					
Aubergines					
Others					
Cucurbits — edible peel					
Cucumbers					
Gherkins					
Courgettes					
Others					
Cucurbits — inedible peel					
Melons					
Squashes					
Watermelons					
Others					
Sweet corn					
(IV) BRASSICA VEGETABLES	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Flowering brassica					
Broccoli					
Cauliflower					
Others					

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
Head brassica					
Brussels sprouts					
Head cabbage					
Others					
Leafy brassica					
Chinese cabbage					
Kale					
Others					
Kohlrabi					
(V) LEAF VEGETABLES AND FRESH HERBS	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Lettuce and similar					
Cress					
Lamb's lettuce					
Lettuce					
Scarole					
Others					
Spinach and similar					
Beet leaves (chord)					
Water cress					
Witloof					
Herbs					
Chervil					
Chives					

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
Parsley					
Celery leaves					
Others					
(VI) LEGUME VEGETABLES (fresh)	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Beans (with pods)					
Beans (without pods)					
Peas (with pods)					
Peas (without pods)					
Others					
(VII) STEM VEGETABLES	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Asparagus					
Cardoons					
Celery					
Fennel					
Globe artichokes					
Leek					
Rhubarb					
Others					
(VIII) FUNGI	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Cultivated mushrooms					
Wild mushrooms					
3. Pulses	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Beans					
Lentils					

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Dinoseb	Dioxathion	Endrin	1,2-dibromo ethane (ethylene dibromide)	Fenclorphos (sum of fenclorphos and fenclorphos oxon expressed as fenclorphos)
Peas					
Others					
4. Oil seeds	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Linseed					
Peanuts					
Poppy seed					
Sesame seed					
Sunflower seed					
Rape seed					
Soya bean					
Mustard					
Cotton seed					
Others					
5. Potatoes	0,05 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,01 (*)
Early potatoes					
Ware potatoes					
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,1 (*)	0,1 (*)	0,01 (*)	0,1 (*)	0,1 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)	0,1 (*)	0,1 (*)	0,01 (*)	0,1 (*)

(*) Indicates lower limit of analytical determination.

(a) (b) (c) (d) Should levels not be adopted by ►M7 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*)

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts		▶ M63 0,2 (*) (P) ◀		▶ M54 0,02 (*) ◀
(I) CITRUS FRUIT	0,01 (*)		0,05 (*)	
Grapefruit				
Lemons				
Limes				
Mandarines (including clementines and similar hybrids)				
Oranges				
Pomelos				
Others				
(II) TREE NUTS (shelled or unshelled)	0,01 (*)			
Almonds				
Brazil nuts				
Cashew nuts				
Chestnuts				
Coconuts				
Hazelnuts				
Macadamia				
Pecans				
Pine nuts				
Pistachios				
Walnuts				
Others				

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
(III) POME FRUIT	0,01 (*)		0,05 (*)	
Apples				
Pears				
Quinces				
Others				
(IV) STONE FRUIT	0,01 (*)			
Apricots				
Cherries				
Peaches (including nectarines and similar hybrids)				
Plums				
Others				
(V) BERRIES AND SMALL FRUIT	0,01 (*)			
Table and wine grapes				
Table grapes				
Wine grapes				
Strawberries (other than wild)			0,05 (*)	
Cane fruit (other than wild)			0,05 (*)	
Blackberries				
Dewberries				
Loganberries				
Raspberries				
Others				
Other small fruit and berries (other than wild)			0,05 (*)	
Bilberries (fruit of species <i>vaccinium myrtillus</i>)				

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
Cranberries Currants (red, black and white) Gooseberries Others Wild berries and wild fruit (VI) MISCELLANEOUS FRUIT	0,01 (*)		0,05 (*) 0,05 (*) (except figs)	
Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes Olives Passion fruit Pineapples Pomegranates Others 2. Vegetables, fresh or uncooked, frozen or dry				► <u>M54</u> 0,02 (*) ◀
(I) ROOT AND TUBER VEGETABLES Beetroot Carrots	0,01 (*)	► <u>M63</u> 30 (P) ◀	0,05 (*)	

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
Celeriac				
Horseradish				
Jerusalem artichokes				
Parsnips		► <u>M63</u> 30 (P) ◀		
Parsley root				
Radishes				
Salsify				
Sweet potatoes				
Swedes				
Turnips				
Yams				
Others		► <u>M63</u> 0,2 (*) (P) ◀		
(II) BULB VEGETABLES	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀	0,05 (*)	
Garlic		► <u>M63</u> 15 (P) ◀		
Onions		► <u>M63</u> 15 (P) ◀		
Shallots		► <u>M63</u> 15 (P) ◀		
Spring onions				
Others		► <u>M63</u> 0,2 (*) (P) ◀		
(III) FRUITING VEGETABLES	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀	0,05 (*)	
Solanacea				
Tomatoes				
Peppers				

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
Aubergines				
Others				
Cucurbits — edible peel				
Cucumbers				
Gherkins				
Courgettes				
Others				
Cucurbits — inedible peel				
Melons				
Squashes				
Watermelons				
Others				
Sweet corn				
(IV) BRASSICA VEGETABLES	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀	0,05 (*)	
Flowering brassica				
Broccoli				
Cauliflower				
Others				
Head brassica				
Brussels sprouts				
Head cabbage				
Others				

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
Leafy brassica				
Chinese cabbage				
Kale				
Others				
Kohlrabi				
(V) LEAF VEGETABLES AND FRESH HERBS	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀	0,05 (*)	
Lettuce and similar				
Cress				
Lamb's lettuce				
Lettuce				
Scarole				
Others				
Spinach and similar				
Beet leaves (chord)				
Water cress				
Witloof				
Herbs				
Chervil				
Chives				
Parsley				
Celery leaves				
Others				

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
(VI) LEGUME VEGETABLES (fresh)	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀	0,05 (*)	
Beans (with pods)				
Beans (without pods)				
Peas (with pods)				
Peas (without pods)				
Others				
(VII) STEM VEGETABLES	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀	0,05 (*)	
Asparagus				
Cardoons				
Celery				
Fennel				
Globe artichokes				
Leek				
Rhubarb				
Others				
(VIII) FUNGI	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀	0,05 (*)	
Cultivated mushrooms				
Wild mushrooms				
3. Pulses	0,01 (*)	► <u>M63</u> 0,2 (*) (P) ◀		► <u>M54</u> 0,02 (*) ◀
Beans				
Lentils				
Peas				
Others				

▼ M1

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Heptachlor (sum of heptachlor and heptachlor epoxide)	Maleic hydrazide	Methyl bromide	Paraquat
4. Oil seeds Linseed Peanuts Poppy seed Sesame seed Sunflower seed Rape seed Soya bean Mustard Cotton seed Others	0,01 (*)	► <u>M63</u> 0,5 (*) (P) ◀	0,1 (*)	► <u>M54</u> 0,02 (*) ◀
5. Potatoes Early potatoes Ware potatoes	0,01 (*)	► <u>M63</u> 50 (P) ◀	0,05 (*)	► <u>M54</u> 0,02 (*) ◀
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,02 (*)	► <u>M63</u> 0,5 (*) (P) ◀	0,05 (*)	► <u>M54</u> 0,05 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,01 (*)	► <u>M63</u> 0,5 (*) (P) ◀	0,05 (*)	► <u>M54</u> 0,05 (*) ◀

(*) Indicates lower limit of analytical determination.

► M63 (P) Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC. ◀

(a) (b) (c) (d) Should levels not be adopted by ► M7 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*)

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	TEPP	Camphchlor (Toxaphene)	2,4,5-T
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts			
(I) CITRUS FRUIT	0,01 (*)	0,1 (*)	0,05 (*)
Grapefruit			
Lemons			
Limes			
Mandarines (including clementines and similar hybrids)			
Oranges			
Pomelos			
Others			
(II) TREE NUTS (shelled or unshelled)	0,01 (*)	0,1 (*)	0,05 (*)
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			
(III) POME FRUIT	0,01 (*)	0,1 (*)	0,05 (*)
Apples			
Pears			
Quinces			
Others			
(IV) STONE FRUIT	0,01 (*)	0,1 (*)	0,05 (*)
Apricots			
Cherries			
Peaches (including nectarines and similar hybrids)			
Plums			
Others			
(V) BERRIES AND SMALL FRUIT	0,01 (*)	0,1 (*)	0,05 (*)
Table and wine grapes			
Table grapes			
Wine grapes			
Strawberries (other than wild)			

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	TEPP	Camphchlor (Toxaphene)	2,4,5-T
Cane fruit (other than wild)			
Blackberries			
Dewberries			
Loganberries			
Raspberries			
Others			
Other small fruit and berries (other than wild)			
Bilberries (fruit of species <i>vaccinium myrtyllus</i>)			
Cranberries			
Currants (red, black and white)			
Gooseberries			
Others			
Wild berries and wild fruit			
(VI) MISCELLANEOUS FRUIT	0,01 (*)	0,1 (*)	0,05 (*)
Avocados			
Bananas			
Dates			
Figs			
Kiwi			
Kumquats			
Litchis			
Mangoes			
Olives			
Passion fruit			
Pineapples			
Pomegranates			
Others			
2. Vegetables, fresh or uncooked, frozen or dry			
(I) ROOT AND TUBER VEGETABLES	0,01 (*)	0,1 (*)	0,05 (*)
Beetroot			
Carrots			
Celeriac			
Horseradish			
Jerusalem artichokes			
Parsnips			
Parsley root			
Radishes			
Salsify			
Sweet potatoes			

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	TEPP	Camphchlor (Toxaphene)	2,4,5-T
Swedes			
Turnips			
Yams			
Others			
(II) BULB VEGETABLES	0,01 (*)	0,1 (*)	0,05 (*)
Garlic			
Onions			
Shallots			
Spring onions			
Others			
(III) FRUITING VEGETABLES	0,01 (*)	0,1 (*)	0,05 (*)
Solanacea			
Tomatoes			
Peppers			
Aubergines			
Others			
Cucurbits — edible peel			
Cucumbers			
Gherkins			
Courgettes			
Others			
Cucurbits — inedible peel			
Melons			
Squashes			
Watermelons			
Others			
Sweet corn			
(IV) BRASSICA VEGETABLES	0,01 (*)	0,1 (*)	0,05 (*)
Flowering brassica			
Broccoli			
Cauliflower			
Others			
Head brassica			
Brussels sprouts			
Head cabbage			
Others			
Leafy brassica			
Chinese cabbage			
Kale			
Others			
Kohlrabi			

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	TEPP	Camphchlor (Toxaphene)	2,4,5-T
(V) LEAF VEGETABLES AND FRESH HERBS	0,01 (*)	0,1 (*)	0,05 (*)
Lettuce and similar			
Cress			
Lamb's lettuce			
Lettuce			
Scarole			
Others			
Spinach and similar			
Beet leaves (chord)			
Water cress			
Witloof			
Herbs			
Chervil			
Chives			
Parsley			
Celery leaves			
Others			
(VI) LEGUME VEGETABLES (fresh)	0,01 (*)	0,1 (*)	0,05 (*)
Beans (with pods)			
Beans (without pods)			
Peas (with pods)			
Peas (without pods)			
Others			
(VII) STEM VEGETABLES	0,01 (*)	0,1 (*)	0,05 (*)
Asparagus			
Cardoons			
Celery			
Fennel			
Globe artichokes			
Leek			
Rhubarb			
Others			
(VIII) FUNGI	0,01 (*)	0,1 (*)	0,05 (*)
Cultivated mushrooms			
Wild mushrooms			
3. Pulses	0,01 (*)	0,1 (*)	0,05 (*)
Beans			
Lentils			
Peas			
Others			

▼ **M1**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	TEPP	Camphchlor (Toxaphene)	2,4,5-T
4. Oil seeds Linseed Peanuts Poppy seed Sesame seed Sunflower seed Rape seed Soya bean Mustard Cotton seed Others	0,01 (*)	0,1 (*)	0,05 (*)
5. Potatoes Early potatoes Ware potatoes	0,01 (*)	0,1 (*)	0,05 (*)
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,02 (*)	0,1 (*)	0,05 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,02 (*)	0,1 (*)	0,05 (*)

(*) Indicates lower limit of analytical determination.

(a) (b) (c) (d) Should levels not be adopted by ► **M7** 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*).

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts			
(i) CITRUS FRUIT	0,02 (*)		► M44 0,05 (**) (P) ◀
grapefruit		► M70 0,1 ◀	
lemons		► M70 0,2 ◀	
limes		► M70 0,2 ◀	
mandarins (including clementines and similar hybrids)		► M70 0,2 ◀	
oranges		► M70 0,1 ◀	
pommelo		► M70 0,1 ◀	
others		► M70 0,02 (**) ◀	
(ii) TREE NUTS (shelled or unshelled)	0,05 (*)	► M70 0,05 (**) ◀	► M44 0,05 (**) (P) ◀
almonds			
brazil nuts			
cashew nuts			
chestnuts			
coconuts			
hazelnuts			
macadamia			
pecans			
pine nuts			
pistachios			
walnuts			
others			
(iii) POME FRUIT		► M70 0,1 ◀	► M44 0,05 (**) (P) ◀
apples	0,02 (*) (s)		
pears			
quinces			
others	0,02 (*)		

▼ **M2**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
(iv) STONE FRUIT	0,02 (*)		
apricots		► M70 0,2 ◀	► M44 0,2 (P) ◀
cherries		► M70 0,1 ◀	► M44 — ◀
peaches (including nectarines and similar hybrids)		► M70 0,2 ◀	► M44 0,2 (P) ◀
plums		► M70 0,1 ◀	► M44 — ◀
others		► M70 0,1 ◀	► M44 0,05 (**)(P) ◀
(v) BERRIES AND SMALL FRUITS	0,02 (*)		► M44 0,05 (**)(P) ◀
(a) <i>table and wine grapes</i>		► M70 0,2 ◀	► M44 — ◀
table grapes			
wine grapes			
(b) <i>strawberries (other than wild)</i>		► M70 0,5 ◀	► M44 — ◀
(c) <i>cane fruit (other than wild):</i>			► M44 — ◀
blackberries			
dewberries			
loganberries			
raspberries		► M70 0,2 ◀	
others		► M70 0,02 (**)* ◀	
(d) <i>other small fruit and berries (other than wild)</i>			► M44 — ◀
bilberries (fruits of species <i>vaccinium myrtilus</i>)			
cranberries			
currants (red, black and white)		► M70 0,1 ◀	
gooseberries		► M70 0,1 ◀	
others		► M70 0,02 (**)* ◀	
(e) <i>wild berries and wild fruit</i>		► M70 0,2 ◀	► M44 — ◀

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
(vi) MISCELLANEOUS	► <u>M34</u> 0,02 (*) ◀		
avocados			
bananas		► <u>M70</u> 0,1 ◀	► <u>M44</u> 0,1 (P) ◀
dates			
figs			
kiwi			
kumquats			
litchis			
mangoes		► <u>M70</u> 0,1 ◀	
▼ <u>M63</u>			
olives (table consumption)		► <u>M70</u> 0,5 ◀	
olives (oil extraction)		► <u>M70</u> 0,5 ◀	
▼ <u>M2</u>			
passion fruit			
pineapples			
pomegranate			
others		► <u>M70</u> 0,02 (***) ◀	► <u>M44</u> 0,05 (**) (P) ◀
2. Vegetables, fresh or uncooked, frozen or dry			
(i) ROOT AND TUBER VEGETABLES	0,02 (*)		► <u>M44</u> 0,05 (**) (P) ◀
Beetroot			
carrots			
celeriac		► <u>M70</u> 0,1 ◀	
horseradish			
jerusalem artichokes			
parsnip			
parsley root			
radishes		► <u>M70</u> 0,1 ◀	

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
salsify			
sweet potatoes			
swedes			
turnips			
yam			
others		► <u>M70</u> 0,02 (**) ◀	
(ii) BULB VEGETABLES	0,02 (*)		► <u>M44</u> 0,05 (**) (P) ◀
garlic			
onions			
shallots			
spring onions		► <u>M70</u> 0,05 ◀	
others		► <u>M70</u> 0,02 (**) ◀	
(iii) FRUITING	0,02 (*)		► <u>M44</u> 0,05 (**) (P) ◀
(a) <i>Solanacea</i>			
tomatoes		► <u>M70</u> 0,1 ◀	
peppers		► <u>M70</u> 0,1 ◀	
aubergines		► <u>M70</u> 0,5 ◀	► <u>M44</u> — ◀
okra		0,1	
others		► <u>M70</u> 0,02 (**) ◀	► <u>M44</u> — ◀
(b) <i>cucurbits — edible peel</i>		► <u>M70</u> 0,1 ◀	► <u>M44</u> — ◀
cucumbers			
gherkins			
courgettes			
others			

▼ M70▼ M2

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
(c) <i>cucurbits — inedible peel</i>		► <u>M70</u> 0,05 ◀	► <u>M44</u> — ◀
melons			
squashes			
watermelons			
others			
(d) <i>sweet corn</i>		► <u>M70</u> 0,05 ◀	► <u>M44</u> — ◀
(iv) BRASSICA VEGETABLES	0,02 (*)		► <u>M44</u> 0,05 (**) (P) ◀
(a) <i>flowering brassicas</i>		► <u>M70</u> 0,1 ◀	
broccoli			
cauliflower			
others			
(b) <i>head brassicas</i>			
brussels sprouts		► <u>M70</u> 0,05 ◀	
head cabbage		► <u>M70</u> 0,2 ◀	
others		► <u>M70</u> 0,02 (**) ◀	
(c) <i>leafy brassicas</i>		► <u>M70</u> 1 ◀	
chinese cabbage			
kale			
others			
(d) <i>kohlrabi</i>		► <u>M70</u> 0,02 (**) ◀	
(v) LEAF VEGETABLES AND FRESH HERBS	0,02 (*)		► <u>M44</u> 0,05 (**) (P) ◀
(a) <i>lettuce and similar</i>			
cress		► <u>M70</u> 1 ◀	
lamb's lettuce		► <u>M70</u> 1 ◀	
lettuce		► <u>M70</u> 0,5 ◀	

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
scarole		► <u>M70</u> 1 ◀	
▼ <u>M70</u>			
ruccola		1	
leaves and stems of brassica		1	
▼ <u>M2</u>			
others		► <u>M70</u> 1 ◀	
(b) <i>spinach and similar</i>		► <u>M70</u> 0,5 ◀	
▼ <u>M34</u>			
Spinach			
▼ <u>M2</u>			
Beet leaves (chard)			
beet leaves (chard)			
▼ <u>M22</u>			
Others			
▼ <u>M2</u>			
(c) <i>watercress</i>		► <u>M70</u> 0,02 (***) ◀	
(d) <i>witloof</i>		► <u>M70</u> 0,02 (***) ◀	
(e) <i>herbs</i>		► <u>M70</u> 1 ◀	
chervil			
chives			
parsley			
celery leaves			
others			
(vi) LEGUME VEGETABLES (fresh)	0,02 (*)		► <u>M44</u> 0,05 (***) (P) ◀
beans (with pods)		► <u>M70</u> 0,2 ◀	
beans (without pods)			
peas (with pods)		► <u>M70</u> 0,2 ◀	
peas (without pods)		► <u>M70</u> 0,2 ◀	
others		► <u>M70</u> 0,02 (***) ◀	

▼ **M2**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
(vii) STEM VEGETABLES	0,02 (*)		► M44 — ◀
asparagus			
cardoons			► M44 — ◀
celery		► M70 0,3 ◀	
fennel		► M70 0,3 ◀	► M44 — ◀
globe artichokes			
leek		► M70 0,3 ◀	► M44 0,1 ^(P) ◀
rhubarb			
others		► M70 0,02 (***) ◀	► M44 0,05 (***) ^(P) ◀
(viii) FUNGI	0,02 (*)		► M44 0,05 (***) ^(P) ◀
cultivated mushrooms		► M70 0,02 (***) ◀	
wild mushrooms		► M70 0,5 ◀	
3. Pulses	0,02 (*)		► M44 0,05 (***) ^(P) ◀
beans			
lentils			
peas			
others			
4. Oil seeds	0,05 (*)	► M70 0,05 (***) ◀	► M44 — ◀
linseed			► M44 — ◀
peanuts			► M44 0,2 ^(P) ◀
poppy seed			
sesame seed			
sunflower seed (with shell)			
rape seed			
soya bean			
mustard			

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Daminozide (sum of daminozide and 1,1-dimethylhydrazine, expressed as daminazide)	Lambda-cyhalothrin	Propiconazole
cotton seed			
others			► <u>M44</u> 0,1 (**) (P) ◀
5. Potatoes	0,02 (*)	► <u>M70</u> 0,02 (**) ◀	► <u>M44</u> 0,05 (**) (P) ◀
early and ware potatoes			
6. Tea (black tea processed from the leaves of <i>camellia sinensis</i>)	0,1 (*)	► <u>M70</u> 1 ◀	► <u>M44</u> 0,1 (**) (P) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)	► <u>M70</u> 10 ◀	► <u>M44</u> 0,1 (**) (P) ◀

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed ascarbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
1. Fruit, fresh, dried or uncooked, preserved by freezing not containing added sugar; nuts				
(i) CITRUS FRUIT	► <u>M49</u> 0,3 ◀	► <u>M12</u> 0,05 (*) ◀	► <u>M12</u> 0,05 (*) ◀	0,05 (*)
grapefruit				
lemons				
limes				
mandarins (including clementines and similar hybrids)				
oranges				
pommelo				
others				
(ii) TREE NUTS (shelled or unshelled)	► <u>M49</u> 0,02 (***) ◀	0,05 (*)	► <u>M12</u> 0,05 (*) ◀	0,05 (*)
almonds				
brazil nuts				
cashew nuts				
chestnuts				
coconuts				
hazelnuts	► <u>M49</u> ——— ◀		(b)	
macadamia				
pecans				
pine nuts				
pistachios				
walnuts				
others	► <u>M49</u> ——— ◀		0,05 (*)	

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed as carbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
(iii) POME FRUIT	► <u>M49</u> 0,02 (**) ◀	► <u>M12</u> 0,05 (*) ◀	0,05 (*)	0,05 (*)
apples				
pears				
quinces				
others				
(iv) STONE FRUIT	► <u>M49</u> 0,02 (**) ◀	► <u>M12</u> 0,05 (*) ◀	0,05 (*)	0,05 (*)
apricots				
cherries				
peaches (including nectarines and similar hybrids)				
plums				
others				
(v) BERRIES AND SMALL FRUIT	► <u>M49</u> 0,02 (**) ◀	0,05 (*)	0,05 (*)	0,05 (*)
(a) <i>Table and wine grapes</i>	► <u>M49</u> — ◀			
table grapes				
wine grapes				
(b) <i>strawberries</i> (other than wild)	► <u>M49</u> — ◀			
(c) <i>cane fruit</i> (other than wild)	► <u>M49</u> — ◀			
blackberries				
dewberries				
loganberries				
raspberries				
others				

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed as carbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
(d) <i>other small fruit and berries</i> (other than wild) bilberries (fruit of species <i>vaccinium myrtilus</i>) cranberries currants (red, black and white) gooseberries (<i>cynorrhodon</i>) others	► <u>M49</u> ——— ◀			
(e) <i>wild berries and wild fruit</i>	► <u>M49</u> ——— ◀			
(vi) MISCELLANEOUS avocados bananas dates figs kiwi kumquats litchis mangoes olives (table consumption) olives (oil extraction) passion fruit pineapples pomegranate others	► <u>M49</u> 0,02 (**) ◀	0,05 (*)	0,05 (*)	0,05 (*)

▼ **M2**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed ascarbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
2. Vegetables, fresh or uncooked, frozen or dry	► M49 0,02 (**) ◀			
(i) ROOT AND TUBER VEGETABLES			0,05 (*)	0,05 (*)
beetroot				
carrots	► M49 ——— ◀	0,1		
celeriac	► M49 ——— ◀			
horseradish				
jerusalem artichokes				
parsnip	► M49 ——— ◀	0,1		
parsley root				
radishes	► M49 ——— ◀			
salsify				
sweet potatoes				
swedes	► M49 ——— ◀	(b)		
turnips	► M49 ——— ◀	(b)		
yam				
others	► M49 ——— ◀	► M12 0,05 (*) ◀		
(ii) BULB VEGETABLES		► M12 0,05 (*) ◀	0,05 (*)	0,05 (*)
garlic	► M49 ——— ◀			
onions	► M49 ——— ◀	(b)		
shallots	► M49 ——— ◀			
spring onions				
others	► M49 ——— ◀	0,05 (*)		

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed ascarbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
(iii) FRUITING		► <u>M12</u> 0,05 (*) ◀	► <u>M12</u> 0,05 (*) ◀	
(a) <i>Solanacea</i>	► <u>M49</u> ——— ◀	0,05 (*)	0,05 (*)	0,05 (*)
tomatoes				
peppers				
aubergines				
others				
(b) <i>cucurbits — edible peel</i>	► <u>M49</u> ——— ◀	0,05 (*)	0,05 (*)	0,05 (*)
cucumbers				
gherkins				
courgettes				
others				
(c) <i>cucurbits — inedible peel</i>	► <u>M49</u> ——— ◀	(b)	(b)	0,05 (*)
melons	► <u>M49</u> ——— ◀			
squashes				
watermelons				
others	► <u>M49</u> ——— ◀			
(d) <i>sweet corn</i>	► <u>M49</u> ——— ◀	0,05 (*)	0,05 (*)	0,05 (*)
(iv) BRASSICA		► <u>M12</u> 0,05 (*) ◀	► <u>M12</u> 0,05 (*) ◀	
(a) <i>flowering brassicas</i>	► <u>M49</u> ——— ◀	(b)	(b)	0,1
broccoli				
cauliflower				
others				

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed as carbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
(b) <i>head brassicas</i> brussels sprouts head cabbage others	► <u>M49</u> ——— ◀	(b)	(b)	0,05
(c) <i>leafy brassicas</i> chinese cabbage kale others	► <u>M49</u> ——— ◀	(b)	0,05 (*)	0,05 (*)
(d) <i>kohlrabi</i>	► <u>M49</u> ——— ◀	(b)	0,05 (*)	0,05 (*)
(v) LEAF VEGETABLES AND FRESH HERBS	► <u>M49</u> ——— ◀	0,05 (*)	0,05 (*)	0,05 (*)
(a) <i>lettuce and similar</i> cress lamb's lettuce lettuce scarole others				
(b) <i>spinach and similar</i> beet leaves (chard)				
(c) <i>watercress</i>				
(d) <i>witloof</i>				

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed ascarbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
(e) <i>herbs</i>				
chervil				
chives				
parsley				
celery leaves				
others				
(vi) LEGUME VEGETABLES (fresh)	► <u>M49</u> ——— ◀	0,05 (*)	0,05 (*)	► <u>M12</u> 0,05 (*) ◀
beans (with pods)	► <u>M49</u> ——— ◀			(b)
beans (without pods)	► <u>M49</u> ——— ◀			(b)
peas (with pods)				
peas (without pods)				
others	► <u>M49</u> ——— ◀			0,05 (*)
(vii) STEM VEGETABLES	► <u>M49</u> ——— ◀	► <u>M12</u> 0,05 (*) ◀	0,05 (*)	► <u>M12</u> 0,05 (*) ◀
asparagus				
cardoons				
celery	► <u>M49</u> ——— ◀	(b)		(b)
fennel				
globe artichokes				
leek	► <u>M49</u> ——— ◀	(b)		
rhubarb				
others	► <u>M49</u> ——— ◀	0,05 (*)		0,05 (*)

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed ascarbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
(viii) FUNGI cultivates mushrooms wild mushrooms	► <u>M49</u> ——— ◀	0,05 (*)	0,05 (*)	0,05 (*)
3. Pulses	► <u>M49</u> 0,02 (**) ◀	0,05 (*)	0,05 (*)	► <u>M12</u> 0,05 (*) ◀
beans	► <u>M49</u> ——— ◀			(b)
lentils				
peas				
others	► <u>M49</u> ——— ◀			0,05 (*)
4. Oil seeds	► <u>M49</u> 0,1 ◀	► <u>M12</u> 0,05 (*) ◀		► <u>M12</u> 0,05 (*) ◀
linseed	► <u>M49</u> ——— ◀			
peanuts	► <u>M49</u> ——— ◀			
poppy seed				
sesame seed	► <u>M49</u> ——— ◀			
sunflower seed	► <u>M49</u> ——— ◀	(b)		
rape seed	► <u>M49</u> ——— ◀	(b)		(b)
soya bean	► <u>M49</u> ——— ◀			(b)
mustard				
cotton seed	► <u>M49</u> ——— ◀	(b)	(b)	(b)
others	► <u>M49</u> ——— ◀	0,05 (*)	0,05 (*)	0,05 (*)
5. Potatoes	► <u>M49</u> 0,02 (**) ◀	0,05 (*)	0,05 (*)	0,05 (*)
Early and ware potatoes				

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Carbofuran (sum of carbofuran and 3-hydroxy-carbofuran expressed as carbofuran)	Carbosulfan	Benfurocarb	Furathiocarb
6. Tea (black tea processed from the leaves of <i>camellia sinensis</i>)	► <u>M49</u> 0,05 (**) ◀	0,1 (*)	0,1 (*)	0,1 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	► <u>M49</u> 0,05 (**) ◀	► <u>M12</u> 1 ◀	5	5

▼ **M2**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
1. Fruit, fresh, dried or uncooked, preserved by freezing not containing added sugar; nuts					
(i) CITRUS FRUIT	► M70 0,02 (**) ◄	► M63 0,5 ◄	► M70 0,05 (**) ◄	0,02 (*)	► M54 0,05 (**) ◄
grapefruit					
lemons					
limes					
mandarins (including clementines and similar hybrids)					
oranges					
pommelo					
others					
(ii) TREE NUTS (shelled or unshelled)	► M70 0,02 (**) ◄	► M63 0,05 (**) ◄	► M70 0,05 (**) ◄	0,02 (*)	► M54 0,1 ◄
almonds					
brazil nuts					
cashew nuts					
chestnuts					
coconuts					
hazelnuts					
macadamia					
pecans					
pine nuts					
pistachios					
walnuts					
others					

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
(iii) POME FRUIT	▶ <u>M70</u> 0,2 ◀	▶ <u>M63</u> 1 ◀	▶ <u>M70</u> 0,05 (**) ◀	0,3	
apples					▶ <u>M54</u> 0,5 ◀
pears					
quinces					
others					▶ <u>M54</u> 0,05 (**) ◀
(iv) STONE FRUIT		▶ <u>M63</u> 0,05 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀	(a)	
apricots	▶ <u>M70</u> 0,3 ◀			▶ <u>M12</u> 0,5 ◀	
cherries	▶ <u>M70</u> 0,2 ◀			▶ <u>M12</u> 1 ◀	▶ <u>M54</u> 3 ◀
peaches (including nectarines and similar hybrids)	▶ <u>M70</u> 0,3 ◀			▶ <u>M12</u> 0,5 ◀	
plums	▶ <u>M70</u> 0,2 ◀				
others	▶ <u>M70</u> 0,02 (**) ◀			▶ <u>M12</u> 0,02 (*) ◀	▶ <u>M54</u> 0,05 (**) ◀
(v) BERRIES AND SMALL FRUIT					
(a) <i>table and wine grapes</i>	▶ <u>M70</u> 0,3 ◀		▶ <u>M70</u> 0,2 ◀	0,3	▶ <u>M54</u> 1 ◀
table grapes		▶ <u>M63</u> 2 ◀			
wine grapes		▶ <u>M63</u> 1 ◀			
(b) <i>strawberries</i> (other than wild)	▶ <u>M70</u> 0,02 (**) ◀	▶ <u>M63</u> 0,5 ◀	▶ <u>M70</u> 0,05 (**) ◀	0,3	▶ <u>M54</u> 0,05 (**) ◀

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
(c) <i>cane fruit</i> (other than wild)	▶ <u>M70</u> 0,02 (**) ◀	▶ <u>M63</u> 0,05 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀		▶ <u>M54</u> 0,05 (**) ◀
blackberries					
dewberries					
loganberries					
raspberries				▶ <u>M12</u> 0,1 ◀	
others				0,02 (*)	
(d) <i>other small fruit and berries</i> (other than wild)	▶ <u>M70</u> 0,02 (**) ◀	▶ <u>M63</u> 0,05 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀		
bilberries (fruit of species <i>vaccinium myrtyllus</i>)					
cranberries					
currants (red, black and white)				1	▶ <u>M54</u> 5 ◀
gooseberries				1	
others				0,02 (*)	▶ <u>M54</u> 0,05 (**) ◀
(e) <i>wild berries and wild fruit</i>		▶ <u>M63</u> 0,05 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀	0,02 (*)	▶ <u>M54</u> 0,05 (**) ◀
(vi) MISCELLANEOUS	▶ <u>M70</u> 0,02 (**) ◀	▶ <u>M63</u> 0,05 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀	0,02 (*)	
avocados					
bananas				▶ <u>M5</u> 0,3 ◀	
dates					
figs					
kiwi					
kumquats					

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
litchis					
mangoes					
olives (table consumption)					
olives (oil extraction)					
passion fruit					
pineapples					► <u>M54</u> 2 ◀
pomegranate					
others					► <u>M54</u> 0,05 (**) ◀
2. Vegetables, fresh or uncooked, frozen or dry					
(i) ROOT AND TUBER VEGETABLES	► <u>M70</u> 0,02 (**) ◀		► <u>M70</u> 0,05 (**) ◀	0,02 (*)	► <u>M54</u> 0,05 (**) ◀
beetroot					
carrots		► <u>M63</u> 0,1 ◀			
celeriac					
horseradish		► <u>M63</u> 0,1 ◀			
jerusalem artichokes					
parsnip		► <u>M63</u> 0,1 ◀			
parsley root					
radishes		► <u>M63</u> 0,1 ◀			
salsify					
sweet potatoes					
swedes					

▼ **M2**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
turnips					
yam					
others		► M63 0,05 (**) ◄			
(ii) BULB VEGETABLES	► M70 0,02 (**) ◄			0,02 (*)	► M54 0,05 (**) ◄
garlic		► M63 0,5 ◄			
onions		► M63 0,5 ◄	► M70 0,2 ◄		
shallots		► M63 0,5 ◄			
spring onions		► M63 0,2 ◄			
others		► M63 0,05 (**) ◄	► M70 0,05 (**) ◄		
(iii) FRUITING				(a)	
(a) <i>solanacea</i>					
tomatoes	► M70 0,05 (**) ◄	► M63 0,2 ◄	► M70 0,5 ◄	► M12 0,5 ◄	► M54 1 ◄
peppers	► M70 0,3 ◄	► M63 0,5 ◄	► M70 0,2 ◄	► M12 0,5 ◄	► M54 3 ◄
aubergines	► M70 0,1 ◄		► M70 0,5 ◄		
others	► M70 0,02 (**) ◄	► M63 0,05 (**) ◄	► M70 0,05 (**) ◄	► M12 0,02 (*) ◄	► M54 0,05 (**) ◄
(b) <i>cucurbits — edible peel</i>			► M70 0,05 (**) ◄	► M12 0,2 ◄	► M54 0,05 (**) ◄
cucumbers	► M70 0,1 ◄	► M63 0,5 ◄	► M70 0,05 (**) ◄		
gherkins					
courgettes					

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
others	► <u>M70</u> 0,02 (**) ◄	► <u>M63</u> 0,05 (**) ◄			
(c) <i>cucurbits — inedible peel</i>	► <u>M70</u> 0,02 (**) ◄			(a)	► <u>M54</u> 0,05 (**) ◄
melons		► <u>M63</u> 0,2 ◄	► <u>M70</u> 0,1 ◄		
squashes					
watermelons		► <u>M63</u> 0,2 ◄	► <u>M70</u> 0,1 ◄		
others		► <u>M63</u> 0,05 (**) ◄	► <u>M70</u> 0,05 (**) ◄		
(d) <i>sweet corn</i>	► <u>M70</u> 0,02 (**) ◄	► <u>M63</u> 0,05 (**) ◄	► <u>M70</u> 0,05 (**) ◄	0,02 (*)	► <u>M54</u> 0,05 (**) ◄
(iv) BRASSICA VEGETABLES			► <u>M70</u> 0,05 (**) ◄	0,02 (*)	► <u>M54</u> 0,05 (**) ◄
(a) <i>flowering brassicas</i>	► <u>M70</u> 0,05 ◄	► <u>M63</u> 0,2 ◄			
broccoli					
cauliflower					
others					
(b) <i>head brassicas</i>					
brussels sprouts					
head cabbage	► <u>M70</u> 0,3 ◄	► <u>M63</u> 1 ◄			
others	► <u>M70</u> 0,2 ◄	► <u>M63</u> 0,05 (**) ◄			
(c) <i>leafy brassicas</i>	► <u>M70</u> 0,3 ◄				
chinese cabbage					
kale		► <u>M63</u> 0,2 ◄			

▼ **M2**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
others		▶ M63 0,05 (**) ◀			
(d) <i>kohlrabi</i>	▶ M70 0,02 (**) ◀	▶ M63 0,05 (**) ◀			
(v) LEAF VEGETABLES AND FRESH HERBS				0,05 (*)	▶ M54 0,05 (**) ◀
(a) <i>lettuce and similar</i>	▶ M70 1 ◀				
cress					
lamb's lettuce		▶ M63 0,2 ◀			
lettuce		▶ M63 2 ◀	▶ M70 0,5 ◀		
scarole		▶ M63 1 ◀			
others		▶ M63 0,05 (**) ◀	▶ M70 0,05 (**) ◀		
(b) <i>spinach and similar</i>	▶ M70 0,02 (**) ◀	▶ M63 0,05 (**) ◀	▶ M70 0,05 (**) ◀		
beet leaves (chard)					
(c) <i>watercress</i>	▶ M70 0,02 (**) ◀	▶ M63 0,05 (**) ◀	▶ M70 0,05 (**) ◀		
(d) <i>witloof</i>	▶ M70 0,02 (**) ◀	▶ M63 0,3 ◀	▶ M70 0,05 (**) ◀		
(e) <i>herbs</i>	▶ M70 0,02 (**) ◀	▶ M63 2 ◀	▶ M70 0,05 (**) ◀		
chervil					
chives					
parsley					
celery leaves					
others					

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
(vi) LEGUME VEGETABLES (fresh)	► <u>M70</u> 0,05 ◀	► <u>M63</u> 0,05 (**)	► <u>M70</u> 0,05 (**)	► <u>M12</u> 0,02 (*)	► <u>M54</u> 0,05 (**)
beans (with pods)					
beans (without pods)					
peas (with pods)				(a)	
peas (without pods)				(a)	
others				0,02 (**)	
(vii) STEM VEGETABLES	► <u>M70</u> 0,02 (**)		► <u>M70</u> 0,05 (**)	► <u>M12</u> 0,02 (*)	► <u>M54</u> 0,05 (**)
Asparagus					
cardoons					
celery					
fennel					
globe artichokes				(a)	
leek		► <u>M63</u> 0,2 ◀			
rhubarb					
others		► <u>M63</u> 0,05 (**)		0,02 (*)	
(viii) FUNGI	► <u>M70</u> 0,02 (**)	► <u>M63</u> 0,05 (**)	► <u>M70</u> 0,05 (**)	0,02 (*)	► <u>M54</u> 0,05 (**)
cultivated mushrooms					
wild mushrooms					

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
3. Pulses	▶ <u>M70</u> 0,02 (**) ◀	▶ <u>M63</u> 0,05 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀	0,02 (*)	▶ <u>M54</u> 0,05 (**) ◀
beans					
lentils					
peas					
others					
4. Oil seeds		▶ <u>M63</u> 0,1 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀	0,02 (*)	
linseed					
peanuts					
poppy seed					
sesame seed					
sunflower seed					
rape seed	▶ <u>M70</u> 0,05 ◀				
soya bean					
mustard					
cotton seed					▶ <u>M54</u> 2 ◀
others					▶ <u>M54</u> 0,1 (**) ◀
5. Potatoes	▶ <u>M70</u> 0,02 (**) ◀	▶ <u>M63</u> 0,05 (**) ◀	▶ <u>M70</u> 0,05 (**) ◀	0,02 (*)	▶ <u>M54</u> 0,05 (**) ◀
early and ware potatoes					

▼ M2

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)				
	Cyfluthrin including other mixed isomeric constituents (sum of isomers)	Metalaxyl	Benalaxyl	Fenarimol	Ethephon
6. Tea (black tea processed from the leaves of <i>camellia sinensis</i>)	► <u>M70</u> 0,1 (**) ◀	► <u>M63</u> 0,1 (**) ◀	► <u>M70</u> 0,1 (**) ◀	0,05 (*)	► <u>M54</u> 0,1 (**) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► <u>M70</u> 20 ◀	► <u>M63</u> 10 ◀	► <u>M70</u> 0,1 (**) ◀	5	► <u>M54</u> 0,1 (**) ◀

x As from 1 January 1996.

(*) Indicates limit of analytical determination.

► M44 (**) Indicates lower limit of analytical determination.

(p) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 13 September 2009. ◀

(a) (b) (c) As from ► M7 at the latest 1 July 2000 ◀, and save for adoption of other levels, the following maximum limits shall apply:

- (a) 0,02 (*)
- (b) 0,05 (*)
- (c) 0,1 (*)

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	► <u>C2</u> Methidathion ◀	Methomyl Thiodicarb: sum of methomyl and thiodicarb expressed as methomyl	Amitraz residue: amitraz plus all its metabolites containing 2,4 dimethylaniline, expressed as amitraz
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts			► <u>M43</u> 0,05 (*) ◀
(i) CITRUS FRUIT	► <u>M70</u> 5 ◀		► <u>M43</u> — ◀
grapefruit		► <u>M57</u> 0,5 ◀	
lemons		► <u>M57</u> 1 ◀	
limes		► <u>M57</u> 1 ◀	
mandarins (including clementines and other hybrids)		► <u>M57</u> 1 ◀	
oranges		► <u>M57</u> 0,5 ◀	► <u>M22</u> — ◀
pomelos		► <u>M57</u> 0,5 ◀	
others		► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
(ii) TREE NUTS (shelled or unshelled)	► <u>M70</u> 0,05 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
almonds			
brazil nuts			
cashew nuts			
chestnuts			
coconuts			
hazelnuts			
Macadamia			
Pecans			
pine nuts			
pistachios			
walnuts			
others			
(iii) POME FRUIT	► <u>M70</u> 0,05 ◀	► <u>M57</u> 0,2 ◀	► <u>M43</u> — ◀
apples			
pears			
quinces			
others			
(iv) STONE FRUIT			► <u>M43</u> — ◀
apricots		► <u>M57</u> 0,2 ◀	
cherries	► <u>M70</u> 0,2 ◀	► <u>M57</u> 0,1 ◀	
peaches (including nectarines, and similar hybrids)		► <u>M57</u> 0,2 ◀	► <u>M22</u> — ◀
plums	► <u>M70</u> 0,2 ◀	► <u>M57</u> 0,5 ◀	
others	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	► <u>C2</u> Methidathion ◀	Methomyl Thiodicarb: sum of methomyl and thiodicarb expressed as methomyl	Amitraz residue: amitraz plus all its metabolites containing 2,4 dimethylaniline, expressed as amitraz
(v) BERRIES AND SMALL FRUIT			► <u>M43</u> — ◀
(a) <i>table and wine grapes</i>	► <u>M70</u> 0,02 (*) ◀		► <u>M22</u> — ◀
table grapes		► <u>M57</u> 0,05 (*) ◀	
wine grapes		► <u>M57</u> 1 ◀	
(b) <i>strawberries</i> (other than wild)	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
(c) <i>cane fruit</i> (other than wild)	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
blackberries			
dewberries			
loganberries			
raspberries			
others			
(d) <i>other small fruit and berries</i> (other than wild)	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	
bilberries (fruit of species <i>vaccinium myrtillus</i>)			
cranberries			
currants (red, black and white)			► <u>M22</u> — ◀
gooseberries			
others			► <u>M22</u> — ◀
(e) <i>wild berries and wild fruit</i>	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
(vi) MISCELLANEOUS		► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
avocados			
bananas			
dates			
figs			
kiwi			
kumquats			
lychees			
mangoes			
olives	► <u>M70</u> 1 ◀		
passion fruit			
pineapples	► <u>M70</u> 0,05 ◀		
pomegranates			
others	► <u>M70</u> 0,02 (*) ◀		

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	► <u>C2</u> Methidathion ◀	Methomyl Thiodicarb: sum of methomyl and thiodicarb expressed as methomyl	Amitraz residue: amitraz plus all its metabolites containing 2,4 dimethylaniline, expressed as amitraz
2. Vegetables, fresh or uncooked, frozen or dry			► <u>M43</u> 0,05 (*) ◀
(i) ROOT AND TUBER VEGETABLES	► <u>M70</u> 0,02 (*) ◀		► <u>M43</u> — ◀
beetroot			
carrots			
celeriac			
horse radish			
jerusalem artichokes			
parsnips			
parsley root			
radishes		► <u>M57</u> 0,5 ◀	
salsify			
sweet potatoes			
swedes			
turnips			
yam			
others		► <u>M57</u> 0,05 (*) ◀	
(ii) BULB VEGETABLES		► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
garlic			
onions	► <u>M70</u> 0,1 ◀		
shallots			
spring onions			
others	► <u>M70</u> 0,02 (*) ◀		
(iii) FRUITING VEGETABLES			
(a) <i>Solanacea</i>			
tomatoes	► <u>M70</u> 0,1 ◀	► <u>M57</u> 0,2 ◀	► <u>M43</u> — ◀
peppers		► <u>M57</u> 0,2 ◀	
aubergines		► <u>M57</u> 0,2 ◀	► <u>M43</u> — ◀
others	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
(b) <i>Cucurbits — edible peel</i>		► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
cucumbers	► <u>M70</u> 0,05 ◀		
gherkins			

▼ C2▼ M3▼ C2▼ M3

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	► <u>C2</u> Methidathion ◀	Methomyl Thiodicarb: sum of methomyl and thiodicarb expressed as methomyl	Amitraz residue: amitraz plus all its metabolites containing 2,4 dimethylaniline, expressed as amitraz
courgettes			
others	► <u>M70</u> 0,02 (*) ◀		
(c) <i>Cucurbits — inedible peel</i>	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
melons			
squashes			
watermelons			
others			
(d) <i>sweetcorn</i>	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
(iv) BRASSICA VEGETABLES			► <u>M43</u> — ◀
(a) <i>flowering brassica</i>	► <u>M70</u> 0,02 (*) ◀		► <u>M22</u> — ◀
broccoli		► <u>M57</u> 0,2 ◀	
cauliflower			
others		► <u>M57</u> 0,05 (*) ◀	
(b) <i>head brassica</i>	► <u>M70</u> 0,1 ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
brussels sprouts			
head cabbage			
others			
(c) <i>leaf brassica</i>	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
chinese cabbage			
kale			
others			
(d) <i>Kohlrabi</i>	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
(v) LEAF VEGETABLES AND FRESH HERBS	► <u>M70</u> 0,02 (*) ◀		► <u>M43</u> — ◀
(a) <i>lettuce and similar</i>			► <u>M22</u> — ◀
crisp			
lamb's lettuce			
lettuce		► <u>M57</u> 0,3 ◀	
scarole			
others		► <u>M57</u> 0,05 (*) ◀	
(b) <i>Spinach and similar</i>			► <u>M22</u> — ◀
spinach		0,05	
beet leaves (chard)			

▼ M57▼ M3

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	► <u>C2</u> Methidathion ◀	Methomyl Thiodicarb: sum of methomyl and thiodicarb expressed as methomyl	Amitraz residue: amitraz plus all its metabolites containing 2,4 dimethylaniline, expressed as amitraz
▼ <u>M57</u> others		0,05 (*)	
▼ <u>M3</u> (c) <i>water cress</i>		► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
(d) <i>witloof</i>		► <u>M57</u> 0,05 (*) ◀	► <u>M22</u> — ◀
(e) <i>herbs</i> chervil chives parsley celery leaves others		► <u>M57</u> 0,3 ◀	► <u>M22</u> — ◀
(vi) LEGUME VEGETABLES (fresh) beans (with pods) beans (without pods) peas (with pods) peas (without pods) others	► <u>M70</u> 0,1 ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
(vii) STEM VEGETABLES (fresh) asparagus cardoons celery fennel globe artichokes leek rhubarb others	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
(viii) FUNGI cultivated mushrooms wild mushrooms	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> — ◀
3. Pulses beans lentils peas others	► <u>M70</u> 0,1 ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> 0,05 (*) ◀
4. Oil seed linseed peanuts	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,1 ◀	

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	► <u>C2</u> Methidathion ◀	Methomyl Thiodicarb: sum of methomyl and thiodicarb expressed as methomyl	Amitraz residue: amitraz plus all its metabolites containing 2,4 dimethylaniline, expressed as amitraz
poppy seeds			
sesame seeds			
▼ <u>C2</u>			
sunflower seed	► <u>M70</u> 0,5 ◀		
rape seed	► <u>M70</u> 0,1 ◀		
▼ <u>M3</u>			
► <u>C2</u> soya bean ◀		► <u>M57</u> 0,1 ◀	
mustard seed			
cotton seed	► <u>M70</u> 1 ◀	► <u>M57</u> 0,1 ◀	► <u>M43</u> 1 (*) ◀
▼ <u>M70</u>			
hemp seed	0,1		
▼ <u>M3</u>			
others	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> 0,05 (*) ◀
5. Potatoes	► <u>M70</u> 0,02 (*) ◀	► <u>M57</u> 0,05 (*) ◀	► <u>M43</u> 0,05 (*) ◀
early and ware potatoes			
6. Tea (Dried leaves and stalks, fermented or otherwise of camellia sinensis)	► <u>M70</u> 0,5 ◀	► <u>M57</u> 0,1 (*) ◀	► <u>M43</u> 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► <u>M70</u> 5 ◀	► <u>M57</u> 10 ◀	► <u>M43</u> 0,1 (*) ◀

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Pirimiphosmethyl	Aldicarb residue: sum of aldicarb, its sulfoxide and its sulfone, expressed as aldicarb
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts		► M60 0,02 (*) ◀
(i) CITRUS FRUIT		
grapefruit		
lemons		
limes		
mandarins (including clementines and other hybrids)	2	
oranges		
pomelo		
others	1	
(ii) TREE NUTS (shelled or unshelled)	► M12 0,05 (*) ◀	
almonds	(b)	
brazil nuts		
cashew nuts		
chestnuts		
coconuts		
hazelnuts	(b)	
macadamia		
pecans		
pine nuts		
pistachios	(b)	
walnuts	(b)	
others	0,05 (*)	
(iii) POME FRUIT	► M12 0,05 (*) ◀	
apples		
pears		
quinces		
others		
(iv) STONE FRUIT	► M12 0,05 (*) ◀	
apricots		
cherries		
peaches (including nectarines and similar hybrids)		
plums		
others		
(v) BERRIES AND SMALL FRUIT		
(a) <i>table and wine grapes</i>	(b)	
table grapes	► M12 0,05 (*) ◀	
wine grapes	► M12 2 ◀	

▼ C2▼ M3

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Pirimiphosmethyl	Aldicarb residue: sum of aldicarb, its sulfoxide and its sulfone, expressed as aldicarb
(b) <i>strawberries</i> (other than wild)	► <u>M12</u> 0,05 (*) ◀	
(c) <i>cane fruit</i> (other than wild)	0,05 (*)	
blackberries		
dewberries		
loganberries		
raspberries		
others		
(d) <i>other small fruit and berries</i> (other than wild)	0,05 (*)	
bilberries (fruit of species <i>vaccinium myrtillus</i>)		
cranberries		
currants (red, black and white)		
gooseberries		
others		
(e) <i>wild berries and wild fruit</i>	0,05 (*)	
(vi) MISCELLANEOUS		
avocados		
bananas		
dates		
figs		
kiwi	2	
kumquats		
lychees		
mangoes		
olives	(b)	
passion fruit		
pineapples		
pomegranate		
others	► <u>M12</u> 0,05 (*) ◀	
2. Vegetables, fresh or uncooked, frozen or dry		
(i) ROOT AND TUBER VEGETABLES		► <u>M60</u> 0,02 (*) ◀
beetroot		
carrots	1	
celeriac		
horse radish		
jerusalem artichokes		
parsnips		
parsley root		
radishes		
salsify		

▼ C2▼ M3

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Pirimiphosmethyl	Aldicarb residue: sum of aldicarb, its sulfoxide and its sulfone, expressed as aldicarb
sweet potatoes		
swedes		
turnips		
yam		
others	0,05 (*)	
(ii) BULB VEGETABLES	► <u>M12</u> 0,05 (*) ◀	► <u>M60</u> 0,05 ◀
garlic		
onions		
shallots		
spring onions		
others		
(iii) FRUITING VEGETABLES		► <u>M60</u> 0,02 (*) ◀
(a) <i>solanacea</i>	(b)	
tomatoes	► <u>M12</u> 1 ◀	
peppers	► <u>M12</u> 1 ◀	
aubergines		
others	► <u>M12</u> 0,05 (*) ◀	
(b) <i>cucurbits — edible peel</i>	(b)	
cucumbers	► <u>M12</u> 0,1 ◀	
gherkins		
courgettes		
others	► <u>M12</u> 0,05 (*) ◀	
(c) <i>cucurbits — inedible peel</i>	(b)	
melons	► <u>M12</u> 1 ◀	
squashes		
watermelons		
others	► <u>M12</u> 0,05 (*) ◀	
(d) <i>sweetcorn</i>	0,05 (*)	
(iv) BRASSICA VEGETABLES		► <u>M60</u> 0,02 (*) ◀
(a) <i>Flowering brassica</i>	1	
broccoli		
cauliflower		
others		
(b) head brassica	2	
brussels sprouts		
head cabbage		
others	► <u>M12</u> 0,05 (*) ◀	
(c) <i>leafy brassica</i>	► <u>M12</u> 0,05 (*) ◀	
chinese cabbage		
kale		

▼ C2▼ M3

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Pirimiphosmethyl	Aldicarb residue: sum of aldicarb, its sulfoxide and its sulfone, expressed as aldicarb
others		
(d) <i>kohlrabi</i>	► <u>M12</u> 0,05 (*) ◀	
(v) LEAF VEGETABLES AND FRESH HERBS	► <u>M12</u> 0,05 (*) ◀	► <u>M60</u> 0,02 (*) ◀
(a) <i>lettuce and similar</i>	(b)	
cress		
lamb's lettuce		
lettuce		
scarole		
others		
(b) <i>spinach and similar</i>	(b)	
beet leaves (chard)		
(c) <i>water cress</i>	0,05 (*)	
(d) <i>Witloof</i>	0,05 (*)	
(e) <i>herbs</i>	(b)	
chervil		
chives		
parsley		
celery leaves		
others		
(vi) LEGUME VEGETABLES (fresh)	► <u>M12</u> 0,05 (*) ◀	► <u>M60</u> 0,02 (*) ◀
beans (with pods)		
beans (without pods)		
peas (with pods)		
peas (without pods)	0,05 (*)	
others	(b)	
(vii) STEM VEGETABLES (fresh)	► <u>M12</u> 0,05 (*) ◀	► <u>M60</u> 0,02 (*) ◀
asparagus		
cardoons		
celery		
fennel		
globe artichokes		
leek		
rhubarb		
others		
(viii) FUNGI		► <u>M60</u> 0,02 (*) ◀
cultivated mushrooms	2	
wild mushrooms	0,05 (*)	
3. Pulses	► <u>M12</u> 0,05 (*) ◀	► <u>M60</u> 0,02 (*) ◀
beans		
lentils		

▼ C2▼ M3

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Pirimiphosmethyl	Aldicarb residue: sum of aldicarb, its sulfoxide and its sulfone, expressed as aldicarb
peas others		
4. Oil seed	► M12 0,05 (*) ◀	► M60 0,05 (*) ◀
linseed	(b)	
peanuts	(b)	
poppy seeds		
sesame seeds		
▼ <u>C2</u> sunflower seed		
▼ <u>M3</u> rape seed	(b)	
soya bean	(b)	
mustard seed		
cotton seed	(b)	
others	0,05 (*)	
5. Potatoes	0,05 (*)	
early and ware potatoes		
6. Tea (Dried leaves and stalks, fermented or otherwise, <i>camellia sinensis</i>)	0,05 (*)	► M60 0,05 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (*)	► M60 0,05 (*) ◀

▼ **M3**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Thiabendazole
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts	
(i) CITRUS FRUIT	▶ M53 5 ◀
grapefruit	
lemons	
limes	
mandarins (including clementines and other hybrids)	
oranges	
pomelo	
others	
(ii) TREE NUTS (shelled or unshelled)	▶ M53 0,1 (*) ◀
almonds	
brazil nuts	
cashew nuts	
chestnuts	
coconuts	
hazelnuts	
macadamia	
pecans	
pine nuts	
pistachios	
walnuts	
others	
(iii) POME FRUIT	▶ M53 5 ◀
apples	▶ M53 5 ◀
pears	▶ M53 5 ◀
quinces	
others	▶ M53 0,05 (*) ◀
(iv) STONE FRUIT	▶ M53 0,05 (*) ◀
apricots	
cherries	
peaches (including, nectarines and similar hybrids)	
plums	
others	
(v) BERRIES AND SMALL FRUIT	▶ M53 0,05 (*) ◀
(a) <i>table and wine grapes</i>	
table grapes	
wine grapes	
(b) <i>strawberries (other than wild)</i>	
(c) <i>cane fruit (other than wild)</i>	
blackberries	
dewberries	
loganberries	

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Thiabendazole
(ii) BULB VEGETABLES	► <u>M53</u> 0,05 (*) ◀
garlic	
onions	
shallots	
spring onions	
others	
(iii) FRUITING VEGETABLES	► <u>M53</u> 0,05 (*) ◀
(a) solanacea	
tomatoes	
peppers	
aubergines	
others	
(b) cucurbits — edible peel	
cucumbers	
gherkins	
courgettes	
others	
(c) cucurbits — inedible peel	
melons	
squashes	
watermelons	
others	
(d) <i>sweetcorn</i>	
(iv) BRASSICA VEGETABLES	
(a) flowering brassica	
Broccoli	► <u>M53</u> 5 ◀
cauliflower	
others	► <u>M53</u> 0,05 (*) ◀
(b) head brassica	► <u>M53</u> 0,05 (*) ◀
Brussels sprouts	
head cabbage	
others	
(c) <i>leafy brassica</i>	► <u>M53</u> 0,05 (*) ◀
chinese cabbage	
kale	
others	
(d) <i>kohlrabi</i>	► <u>M53</u> 0,05 (*) ◀
(v) LEAF VEGETABLES AND FRESH HERBS	► <u>M53</u> 0,05 (*) ◀
(a) lettuce and similar	
cress	
lamb's lettuce	
lettuce	

▼ C2▼ M3

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Thiabendazole
scarole	
others	
(b) <i>spinach and similar</i>	
beet leaves (chard)	
(c) <i>water cress</i>	
(d) <i>witloof</i>	► <u>M53</u> 1 ◀
(e) <i>herbs</i>	
chervil	
chives	
parsley	
celery leaves	
others	
(vi) LEGUME VEGETABLES (fresh)	► <u>M53</u> 0,05 (*) ◀
beans (with pods)	
beans (without pods)	
peas (with pods)	
peas (without pods)	
others	
(vii) STEM VEGETABLES (fresh)	► <u>M53</u> 0,05 (*) ◀
asparagus	
cardoons	
celery	
fennel	
globe artichokes	
leek	
rhubarb	
others	
(viii) FUNGI	
cultivated mushrooms	► <u>M53</u> 10 ◀
wild mushrooms	► <u>M53</u> 0,05 (*) ◀
3. Pulses	► <u>M53</u> 0,05 (*) ◀
beans	
lentils	
peas	
others	
4. Oil Seed	► <u>M53</u> 0,05 (*) ◀
linseed	
peanuts	
poppy seeds	
sesame seeds	
▼ <u>C2</u>	
sunflower seed	
▼ <u>M3</u>	
rape seed	
soya bean	

▼ M3

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Thiabendazole
mustard seed	
cotton seed	
others	
5. Potatoes	
early potatoes	▶ <u>M53</u> 0,05 (*) ◀
ware potatoes	▶ <u>M53</u> 15 ◀
6. Tea (Dried leaves and stalks fermented or otherwise, <i>camellia sinensis</i>)	▶ <u>M53</u> 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	▶ <u>M53</u> 0,1 (*) ◀

▼ C2▼ M3

(*) Indicates lower limit of analytical determination.

▶ M43 (*) Should this level not be confirmed or amended by a directive, with effect from 1 July 2007, the appropriate lower limit of analytical determination shall apply. ◀

(a) (b) Should levels not be adopted by 1 July 2000, the following maximum levels shall apply:

(a) 0,02 (*),

(b) 0,05 (*).

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Triforine	Endosulfan (Sum of alpha and beta endosulfan and endosulfan sulphate expressed as endosulfan)	Fentin (Fentin, expressed as triphenyltin cation)
1. Fruit, fresh, dried or uncooked preserved by freezing, not containing added sugar; nuts			
(i) CITRUS FRUIT	0,05 (*)	► M52 0,05 (*) ◀	0,05 (*)
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)	► M12 0,05 (*) ◀	► M52 0,1 (*) ◀	0,05 (*)
Almonds	(a)		
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others	0,05 (*)		
(iii) POME FRUIT	2		0,05 (*)
Apples			
Pears		► M52 0,3 ◀	
Quinces			
Other		► M52 0,05 (*) ◀	
(iv) STONE FRUIT		► M52 0,05 (*) ◀	0,05 (*)
Apricots	► M12 2 ◀		
Cherries	2		
Peaches (including nectarines and similar hybrids)	► M12 2 ◀		
Plums	► M12 1 ◀		
Others	0,05 (*)		
(v) BERRIES AND SMALL FRUIT			0,05 (*)
(a) <i>Table and wine grapes</i>	► M12 0,05 (*) ◀	► M52 0,5 ◀	
(b) <i>Strawberries</i> (other than wild)	► M12 0,05 (*) ◀	► M52 0,05 (*) ◀	

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Triforine	Endosulfan (Sum of alpha and beta endosulfan and endosulfan sulphate expressed as endosulfan)	Fentin (Fentin, expressed as triphenyltin cation)
(c) <i>Cane fruit (other than wild)</i>	0,05 (*)	► $\frac{M52}{(*)}$ 0,05 (*) ◀	
Blackberries			
Dewberries			
Loganberries			
Raspberries			
Others			
(d) <i>Other small fruit and berries (other than wild)</i>		► $\frac{M52}{(*)}$ 0,05 (*) ◀	
Bilberries (fruit of species <i>Vaccinium myrtilus</i>)			
Cranberries			
Currants (red, black and white)	2		
Gooseberries	2		
Others	► $\frac{M12}{(*)}$ 0,05 (*) ◀		
(e) <i>Wild berries and wild fruit</i>	0,05 (*)	► $\frac{M52}{(*)}$ 0,05 (*) ◀	
(vi) MISCELLANEOUS	0,05 (*)	► $\frac{M52}{(*)}$ 0,05 (*) ◀	0,05 (*)
Avocados			
Bananas			
Dates			
Figs			
Kiwis			
Kumquats			
Litchis			
Mangoes			
Olives			
Passion fruit			
Pineapples			
Pomegranates			
Others			
2. Vegetables, fresh or uncooked, frozen or dry			
(i) ROOT AND TUBER VEGETABLES	► $\frac{M12}{(*)}$ 0,05 (*) ◀	► $\frac{M52}{(*)}$ 0,05 (*) ◀	0,05 (*)
Beetroot			
Carrots			
Celeriac			
Horseradish			
Jerusalem artichokes			
Parsnips			
Parsley root			

▼ M5

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Triforine	Endosulfan (Sum of alpha and beta endosulfan and endosulfan sulphate expressed as endosulfan)	Fentin (Fentin, expressed as triphenyltin cation)
Radishes			
Salsify			
Sweet potatoes			
Swedes	(a)		
Turnips			
Yams			
Others	0,05 (*)		
(ii) BULB VEGETABLES	► <u>M12</u> 0,05 (*) ◀	► <u>M52</u> 0,05 (*) ◀	0,05 (*)
Garlic			
Onions			
Shallots			
Spring onions			
Others			
(iii) FRUITING VEGETABLES			0,05 (*)
(a) <i>Solanacea</i>	► <u>M12</u> 0,05 (*) ◀		
Tomatoes		► <u>M52</u> 0,5 ◀	
Peppers		► <u>M52</u> 1 ◀	
Aubergines			
Others		► <u>M52</u> 0,05 (*) ◀	
(b) <i>Cucurbits — edible peel</i>	0,5	► <u>M52</u> 0,05 (*) ◀	
Cucumbers			
Gherkins			
Courgettes			
Others			
(c) <i>Cucurbits — inedible peel</i>	► <u>M12</u> 0,05 (*) ◀	► <u>M52</u> 0,05 (*) ◀	
Melons			
Squashes			
Watermelons			
Others			
(d) <i>Sweetcorn</i>	0,05 (*)	► <u>M52</u> 0,05 (*) ◀	
(iv) BRASSICA VEGETABLES	► <u>M12</u> 0,05 (*) ◀	► <u>M52</u> 0,05 (*) ◀	0,05 (*)
(a) <i>Flowering brassica</i>			
Broccoli			
Cauliflower			
Others			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Triforine	Endosulfan (Sum of alpha and beta endosulfan and endosulfan sulphate expressed as endosulfan)	Fentin (Fentin, expressed as triphenyltin cation)
(b) <i>Head brassica</i> Brussels sprouts Head cabbage Others			
(c) <i>Leafy brassica</i> Chinese cabbage Kale Others			
(d) <i>Kohlrabi</i>			
(v) LEAF VEGETABLES AND FRESH HERBS	► M12 0,05 (*) ◀	► M52 0,05 (*) ◀	0,05 (*)
(a) <i>Lettuce and similar</i> Cress Lamb's lettuce Lettuce Scarole Others	(a) 0,05 (*)		
(b) <i>Spinach and similar</i> Spinach Beet leaves (chard) Others	(a) 0,05 (*)		
(c) <i>Watercress</i>	0,05 (*)		
(d) <i>Witloof</i>	0,05 (*)		
(e) <i>Herbs</i> Chervil Chives Parsley Celery leaves Others	(a) 0,05 (*)		
(vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others	► M12 0,05 (*) ◀	► M52 0,05 (*) ◀	0,05 (*)
(vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes	► M12 0,05 (*) ◀ (a) (a) (a)	► M52 0,05 (*) ◀	0,05 (*)

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Triforine	Endosulfan (Sum of alpha and beta endosulfan and endosulfan sulphate expressed as endosulfan)	Fentin (Fentin, expressed as triphenyltin cation)
Leeks	(a)		
Rhubarb			
Others	0,05 (*)		
(viii) FUNGI	0,05 (*)	► <u>M52</u> 0,05 (*) ◀	0,05 (*)
(a) <i>Cultivated mushrooms</i>			
(b) <i>Wild mushrooms</i>			
3. Pulses	0,05 (*)	► <u>M52</u> 0,05 (*) ◀	0,05 (*)
Beans			
Lentils			
Peas			
Others			
4. Oil seed	0,05 (*)		0,05 (*)
Linseed			
Peanuts			
Poppy seeds			
Sesame seeds			
Sunflower seed			
Rapeseed			
Soya bean		► <u>M52</u> 0,5 ◀	
Mustard seed			
Cotton seed		► <u>M52</u> 5 ◀	
Others		► <u>M52</u> 0,1 (*) ◀	
5. Potatoes	0,05 (*)	► <u>M52</u> 0,05 (*) ◀	0,1
Early and ware potatoes			
6. Tea (Black tea processed from the leaves of <i>Camellia sinensis</i>)	0,1 (*)	► <u>M52</u> 30 ◀	0,1 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	30	► <u>M52</u> 0,1 (*) ◀	0,5

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Phorate (Sum of phorate, its oxygen analogue and their sulphoxides and sulphones expressed as phorate)	Dicofol (Sum of P, P'- and O, P'- isomers)	Chlormequat
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts	► M12 0,05 (*) ◀		
(i) CITRUS FRUIT	0,05 (*)	► M18 2 ◀	► M70 0,05 (*) ◀
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)	0,05 (*)	0,05 (*)	► M70 0,1 (*) ◀
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			
(iii) POME FRUIT	0,05 (*)	► M18 0,02 (*) ◀	
Apples			
Pears			► M70 0,2 (*) ◀
Quinces			
Other			► M70 0,05 (*) ◀
(iv) STONE FRUIT	0,05 (*)	► M18 0,02 (*) ◀	► M70 0,05 (*) ◀
Apricots			
Cherries			
Peaches (including nectarines and similar hybrids)			
Plums			
Others			

▼ M5

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Phorate (Sum of phorate, its oxygen analogue and their sulphoxides and sulphones expressed as phorate)	Dicofol (Sum of P, P'- and O, P'- isomers)	Chloromequat
(v) BERRIES AND SMALL FRUIT			► <u>M70</u> 0,05 (*) ◀
(a) <i>Table and wine grapes</i>	0,05 (*)	► <u>M18</u> 2 ◀	
▼ <u>M12</u>			
Table grapes		► <u>M18</u> ◀	
Wine grapes		► <u>M18</u> ◀	
▼ <u>M5</u>			
(b) <i>Strawberries</i> (other than wild)	(a)	► <u>M18</u> 0,02 (*) ◀	
(c) <i>Cane fruit</i> (other than wild)	0,05 (*)	► <u>M18</u> 0,02 (*) ◀	
Blackberries			
Dewberries			
Loganberries			
Raspberries			
Others			
(d) <i>Other small fruit and berries</i> (other than wild)	0,05 (*)	► <u>M18</u> 0,02 (*) ◀	
Bilberries (fruit of species <i>Vaccinium myrtilus</i>)			
Cranberries			
Currants (red, black and white)		(b)	
Gooseberries			
Others		0,02 (*)	
(e) <i>Wild berries and wild fruit</i>	0,05 (*)	► <u>M18</u> 0,02 (*) ◀	
(vi) MISCELLANEOUS	0,05 (*)	► <u>M18</u> 0,02 (*) ◀	
Avocados			
Bananas		2 (b)	
Dates			
Figs		(b)	
Kiwis			
Kumquats			
Litchis			
Mangoes			
Olives			► <u>M70</u> 0,1 (*) ◀
Passion fruit			
Pineapples			
Pomegranates			
Others		0,02 (*)	► <u>M70</u> 0,05 (*) ◀

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Phorate (Sum of phorate, its oxygen analogue and their sulphoxides and sulphones expressed as phorate)	Dicofol (Sum of P, P'- and O, P'- isomers)	Chloromequat
2. Vegetables, fresh or uncooked, frozen or dry	► M12 0,05 (*) ◀		
(i) ROOT AND TUBER VEGETABLES		0,02 (*)	► M70 0,05 (*) ◀
Beetroot	(a)		
Carrots	(a)		
Celeriac			
Horseradish			
Jerusalem artichokes			
Parsnips	(a)		
Parsley root			
Radishes			
Salsify			
Sweet potatoes			
Swedes			
Turnips			
Yams			
Others	0,05 (*)		
(ii) BULB VEGETABLES	0,05 (*)	► M18 0,02 (*) ◀	► M70 0,05 (*) ◀
Garlic		(b)	
Onions			
Shallots			
Spring onions			
Others		0,02 (*)	
(iii) FRUITING VEGETABLES			► M70 0,05 (*) ◀
(a) <i>Solanacea</i>	(a)	► M18 — ◀	
Tomatoes		► M18 1 ◀	
Peppers		0,5 (b)	
Aubergines			
Others		► M18 0,02 (*) ◀	
(b) <i>Cucurbits — edible peel</i>		► M18 0,2 ◀	
Cucumbers	0,05 (*)		
Gherkins			
Courgettes			
Others	(a)		
(c) <i>Cucurbits — inedible peel</i>	0,05 (*)	► M18 0,5 ◀	
Melons			
Squashes			
Watermelons			
Others			

▼ M5

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Phorate (Sum of phorate, its oxygen analogue and their sulphoxides and sulphones expressed as phorate)	Dicofol (Sum of P, P'- and O, P'- isomers)	Chlormequat
(d) <i>Sweetcorn</i>	(a)	0,02 (*)	
(iv) BRASSICA VEGETABLES		0,02 (*)	► <u>M70</u> 0,05 (*) ◀
(a) <i>Flowering brassica</i>	(a)		
Broccoli			
Cauliflower			
Others			
(b) <i>Head brassica</i>	(a)		
Brussels sprouts			
Head cabbage			
Others			
(c) <i>Leafy brassica</i>	(a)		
Chinese cabbage			
Kale			
Others			
(d) <i>Kohlrabi</i>	0,05 (*)		
(v) LEAF VEGETABLES AND FRESH HERBS		0,02 (*)	► <u>M70</u> 0,05 (*) ◀
(a) <i>Lettuce and similar</i>	(a)		
Cress			
Lamb's lettuce			
Lettuce			
Scarole			
Others			
(b) <i>Spinach and similar</i>	0,05 (*)		
Spinach			
Beet leaves (chard)			
Others			
(c) <i>Watercress</i>	0,05 (*)		
(d) <i>Witloof</i>	0,05 (*)		
(e) <i>Herbs</i>	(a)		
Chervil			
Chives			
Parsley			
Celery leaves			
Others			
(vi) LEGUME VEGETABLES (fresh)	(a)	► <u>M18</u> 0,02 (*) ◀	► <u>M70</u> 0,05 (*) ◀
Beans (with pods)		0,5 (b)	
Beans (without pods)		0,5 (b)	
Peas (with pods)		0,5 (b)	
Peas (without pods)		0,5 (b)	
Others		0,02 (*)	

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Phorate (Sum of phorate, its oxygen analogue and their sulphoxides and sulphones expressed as phorate)	Dicofol (Sum of P, P'- and O, P'- isomers)	Chlormequat
(vii) STEM VEGETABLES (fresh)		► M18 0,02 (*) ◀	► M70 0,05 (*) ◀
Asparagus			
Cardoons			
Celery	(a)		
Fennel			
Globe artichokes		(b)	
Leeks			
Rhubarb			
Others	0,05 (*)	0,02 (*)	
(viii) FUNGI	0,05 (*)	► M18 0,02 (*) ◀	
(a) <i>Cultivated mushrooms</i>		(b)	► M70 10 ◀
(b) <i>Wild mushrooms</i>		0,02 (*)	► M70 0,05 (*) ◀
3. Pulses	► M12 0,05 (*) ◀	► M18 0,02 (*) ◀	► M70 0,05 (*) ◀
Beans	(a)	(b)	
Lentils			
Peas			
Others	0,05 (*)	0,02 (*)	
4. Oilseed			
Linseed	(a)		► M70 7 ◀
Peanuts	0,1		
Poppy seeds			
Sesame seeds			
Sunflower seed			
Rapeseed	(a)		► M70 7 ◀
Soya bean			
Mustard seed			
Cotton seed		0,1	
Others	► M12 0,05 (*) ◀	0,05 (*)	► M70 0,1 (*) ◀
5. Potatoes	(a)	0,02 (*)	► M70 0,05 (*) ◀
Early and Ware potatoes			
6. Tea (Black tea processed from the leaves of <i>Camellia sinensis</i>)	0,1 (*)	(d) (laid down in Directive 93/58/EEC)	► M70 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)	50	► M70 0,1 (*) ◀

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Propyzamide	Propoxur	Disulfoton (Sum of disulfoton, disulfoton sulphoxide and disulfoton sulphone expressed as disulfoton)
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts	► M42 0,02 (*)(P) ◀		► M12 0,02 (*) ◀
(i) CITRUS FRUIT	► M42 — ◀	3 (a)	0,02 (*)
Grapefruit			
Lemons		► M12 0,3 ◀	
Limes		► M12 0,3 ◀	
Mandarins (including clementines and other hybrids)		► M12 0,3 ◀	
Oranges			
Pomelos			
Others		► M12 0,05 (*) ◀	
(ii) TREE NUTS (shelled or unshelled)	► M42 — ◀	0,05 (*)	0,02 (*)
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Other			
(iii) POME FRUIT	► M42 — ◀	► M12 0,05 (*) ◀	0,02 (*)
Apples			
Pears			
Quinces			
Other			
(iv) STONE FRUIT	► M42 — ◀	► M12 0,05 (*) ◀	0,02 (*)
Apricots			
Cherries			
Peaches (including nectarines and similar hybrides)			
Plums			
Others			
(v) BERRIES AND SMALL FRUIT	► M42 — ◀		
(a) <i>Table and wine grapes</i>	► M42 — ◀	► M12 0,05 (*) ◀	0,02 (*)

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Propyzamide	Propoxur	Disulfoton (Sum of disulfoton, disulfoton sulphoxide and disulfoton sulphone expressed as disulfoton)
(b) <i>Strawberries</i> (other than wild)	► M42 — ◀	► M12 0,05 (*) ◀	(b)
(c) <i>Cane fruit</i> (other than wild)	► M42 — ◀	► M12 0,05 (*) ◀	0,02 (*)
Blackberries		3 (a)	
Dewberries			
Loganberries			
Raspberries		3 (a)	
Others		0,05 (*)	
(d) <i>Other small fruit and berries</i> (other than wild)			0,02 (*)
Bilberries (fruit of species <i>Vaccinium myrtilus</i>)			
Cranberries			
Currants (red, black and white)	► M42 — ◀	0,2	
Gooseberries	► M42 — ◀	► M12 0,2 ◀	
Others	► M42 — ◀	0,05 (*)	
(e) <i>Wild berries and wild fruit</i>	► M42 — ◀	0,05 (*)	0,02 (*)
(vi) MISCELLANEOUS	► M42 — ◀	► M12 0,05 (*) ◀	
Avocados			
Bananas			
Dates			
Figs			
Kiwis			
Kumquats			
Litchis			
Mangoes			
Olives		3 (a)	
Passion fruit			
Pineapples			(b)
Pomegranates			
Others		0,05 (*)	0,02 (*)
2. Vegetables, fresh or uncooked, frozen or dry			► M12 0,02 (*) ◀
(i) ROOT AND TUBER VEGETABLES	► M42 0,02 (*) (P) ◀	► M12 0,05 (*) ◀	
Beetroot		3 (a)	
Carrots			3 (b)
Celeriac		3 (a)	
Horseradish			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Propyzamide	Propoxur	Disulfoton (Sum of disulfoton, disulfoton sulphoxide and disulfoton sulphone expressed as disulfoton)
Jerusalem artichokes			
Parsnips			(b)
Parsley root			
Radishes			
Salsify			
Sweet potatoes			
Swedes			
Turnips			
Yams			
Others		0,05 (*)	0,02 (*)
(ii) BULB VEGETABLES	► M42 0,02 (*) (P) ◀	► M12 0,05 (*) ◀	0,02 (*)
Garlic			
Onions			
Shallots			
Spring onions			
Others			
(iii) FRUITING VEGETABLES	► M42 0,02 (*) (P) ◀	► M12 0,05 (*) ◀	
(a) <i>Solanacea</i>			0,02 (*)
Tomatoes		(^x)	
Peppers		3 (a)	
Aubergines		3 (a)	
Others		3 (a)	
(b) <i>Cucurbits — edible peel</i>			0,02 (*)
Cucumbers		(^x)	
Gherkins		3 (a)	
Courgettes		(a)	
Others			
(c) <i>Cucurbits — inedible peel</i>		3 (a)	
Melons			
Squashes			(b)
Watermelons			
Others			0,02 (*)
(d) <i>Sweet corn</i>		0,05 (*)	0,02 (*)
(iv) BRASSICA VEGETABLES	► M42 0,02 (*) (P) ◀	3 (a)	
(a) <i>Flowering brassica</i>	► M42 — ◀	► M12 0,5 ◀	
Broccoli (including calabrese)			(b)
Cauliflower			(b)
Others			0,02 (*)

▼ M5

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Propyzamide	Propoxur	Disulfoton (Sum of disulfoton, disulfoton sulphoxide and disulfoton sulphone expressed as disulfoton)
(b) Head brassica			
Brussels sprouts			(b)
Head cabbage	► <u>M42</u> — ◀	► <u>M12</u> 0,5 ◀	(b)
Others	► <u>M42</u> — ◀	► <u>M12</u> 0,05 (*) ◀	0,02 (*)
(c) <i>Leafy brassica</i>	► <u>M42</u> — ◀	► <u>M12</u> 0,05 (*) ◀	0,02 (*)
Chinese cabbage			
Kale			
Others			
(d) <i>Kohlrabi</i>	► <u>M42</u> — ◀	► <u>M12</u> 0,05 (*) ◀	(b)
(v) LEAF VEGETABLES AND FRESH HERBS		► <u>M12</u> 0,05 (*) ◀	
(a) <i>Lettuce and similar</i>	► <u>M42</u> 1 (P) ◀		0,02 (*)
Cress		0,05 (*)	
Lamb's lettuce			
Lettuce			
Scarole			
Others		3 (a)	
(b) <i>Spinach and similar</i>	► <u>M42</u> 0,02 (*) (P) ◀	3 (a)	0,02 (*)
Spinach			
Beet leaves (chard)			
Others			
(c) <i>Watercress</i>	► <u>M42</u> 0,02 (*) (P) ◀	0,05 (*)	0,02 (*)
(d) <i>Witloof</i>	► <u>M42</u> 0,02 (*) (P) ◀	0,05 (*)	0,02 (*)
(e) <i>Herbs</i>	► <u>M42</u> 1 (P) ◀	3 (a)	(b)
Chervil			
Chives			
Parsley			
Celery leaves			
Others			
(vi) LEGUME VEGETABLES (fresh)	► <u>M42</u> 0,02 (*) (P) ◀	► <u>M12</u> 0,05 (*) ◀	
Beans (with pods)	► <u>M42</u> — ◀	3 (a)	
Beans (without pods)	► <u>M42</u> — ◀		
Peas (with pods)		3 (a)	

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Propyzamide	Propoxur	Disulfoton (Sum of disulfoton, disulfoton sulphoxide and disulfoton sulphone expressed as disulfoton)
Peas (without pods)			0,02 (*)
Others	► M42 — ◀	0,05 (*)	(b)
(vii) STEM VEGETABLES (fresh)	► M42 0,02 (*) (P) ◀		
Asparagus			
Cardoons		3 (a)	
Celery		3 (a)	(b)
Fennel		3 (a)	
Globe artichokes	► M42 — ◀	3 (a)	
Leeks		1	
Rhubarb			
Others	► M42 — ◀	► M12 0,05 (*) ◀	0,02 (*)
(viii) FUNGI	► M42 0,02 (*) (P) ◀	0,05 (*)	0,02 (*)
(a) <i>Cultivated mushrooms</i>			
(b) <i>Wild mushrooms</i>			
3. Pulses	► M42 0,02 (*) (P) ◀	0,05 (*)	► M12 0,02 (*) ◀
Beans			(b)
Lentils			
Peas			
Others			0,02 (*)
4. Oilseed	► M42 0,05 (*) (P) ◀	0,05 (*)	
Linseed	► M42 — ◀		
Peanuts	► M42 — ◀		
Poppy seeds			
Sesame seeds			
Sunflower seed			
Rapeseed	► M42 — ◀		
Soya bean			
Mustard seed			
Cotton seed	► M42 — ◀		0,05 (*)
Others	► M42 — ◀		0,02 (*)
5. Potatoes	► M42 0,02 (*) (P) ◀	0,05 (*)	► M12 0,02 (*) ◀
Early and ware potatoes			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Propyzamide	Propoxur	Disulfoton (Sum of disulfoton, disulfoton sulphoxide and disulfoton sulphone expressed as disulfoton)
6. Tea (Black tea processed from the leaves of <i>Camellia sinensis</i>)	► M42 0,05 (*) (P) ◀	0,1 (*)	0,05 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	► M42 0,05 (*) (P) ◀	0,1 (*)	► M12 0,05 (*) ◀

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Fenbutatin oxide	Triazophos	Diazinon
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts		► M54 0,01 (*) ◀	
(i) CITRUS FRUIT	► M51 5 ◀		► M64 0,01 (*) ◀
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)	► M51 0,05 (*) ◀		
Almonds			► M64 0,05 ◀
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			► M64 0,01 (*) ◀
(iii) POME FRUIT	► M51 2 ◀		► M64 0,01 (*) ◀
Apples			
Pears			
Quinces			
Others			
(iv) STONE FRUIT	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
Apricots			
Cherries			
Peaches (including nectarines and similar hybrids)			
Plums			
Others			
(v) BERRIES AND SMALL FRUIT			
(a) <i>Table and wine grapes</i>	► M51 2 ◀		► M64 0,01 (*) ◀

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Fenbutatin oxide	Triazophos	Diazinon
(b) <i>Strawberries</i> (other than wild)	► M51 1 ◀		► M64 0,01 (*) ◀
► M12 (c) <i>Cane fruit</i> (other than wild) ◀			► M64 0,01 (*) ◀
Blackberries	► M51 5 ◀		
Dewberries			
Loganberries			
Raspberries	► M51 5 ◀		
Others	► M51 0,05 (*) ◀		
(d) <i>Other small fruit and berries</i> (others than wild)	► M51 0,05 (*) ◀		
Bilberries (fruit of species <i>Vaccinium myrtyllus</i>)			
Cranberries			► M64 0,2 ◀
Currants (red, black and white)			
Gooseberries			
Others			► M64 0,01 (*) ◀
(e) <i>Wild berries and wild fruit</i>	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
(vi) MISCELLANEOUS			
Avocados			
Bananas	► M51 3 ◀		
Dates			
Figs			
Kiwis			
Kumquats			
Litchis			
Mangoes			
Olives			
Passion fruit			
Pineapples			► M64 0,3 ◀
Pomegranates			
Others	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
2. Vegetables, fresh or uncooked, frozen or dry			
(i) ROOT AND TUBER	► M51 0,05 (*) ◀	► M54 0,01 (*) ◀	
Beetroot			
Carrots			
Celeriac			
Horseradish			
Jerusalem artichokes			
Parsnips			
Parsley root			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Fenbutatin oxide	Triazophos	Diazinon
Radishes			► M64 0,1 ◀
Salsify			
Sweet potatoes			
Swedes			
Turnips			
Yams			
Others			► M64 0,01 (*) ◀
(ii) BULB VEGETABLES	► M51 0,05 (*) ◀		
Garlic			
Onions			► M64 0,05 ◀
Shallots			
Spring onions			
Others			► M64 0,01 (*) ◀
(iii) FRUITING VEGETABLES			
(a) <i>Solanacea</i>	► M51 1 ◀		
Tomatoes			
Peppers			► M64 0,05 ◀
Aubergines			
Others			► M64 0,01 (*) ◀
(b) <i>Cucurbits — edible peel</i>			► M64 0,01 (*) ◀
Cucumbers	► M51 0,5 ◀		
Gherkins			
Courgettes	► M51 0,5 ◀		
Others	► M51 0,05 (*) ◀		
(c) <i>Cucurbits — inedible peel</i>	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
Melons			
Squashes			
Watermelons			
Others			
(d) <i>Sweetcorn</i>	► M51 0,05 (*) ◀		► M64 0,02 (*) ◀
(iv) BRASSICA VEGETABLES	► M51 0,05 (*) ◀		
(a) <i>Flowering brassica</i>			► M64 0,01 (*) ◀
Broccoli			
Cauliflower			
Others			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Fenbutatin oxide	Triazophos	Diazinon
(b) <i>Head brassica</i>			
Brussels sprouts			
Head cabbage			► M64 0,5 ◀
Others			► M64 0,01 (*) ◀
(c) <i>Leafy brassica</i>			
Chinese cabbage			► M64 0,05 ◀
Kale			
Others			► M64 0,01 (*) ◀
(d) <i>Kohlrabi</i>			► M64 0,2 ◀
(v) LEAF VEGETABLES AND FRESH HERBS	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
(a) Lettuce and similar			
Cress			
Lamb's lettuce			
Lettuce			
Scarole			
Others			
(b) Spinach and similar			
Spinach			
Beet leaves (chard)			
Others			
(c) <i>Watercress</i>			
(d) <i>Witloof</i>			
(e) Herbs			
Chervil			
Chives			
Parsley			
Celery leaves			
Others			
(vi) LEGUME VEGETABLES	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
Beans (with pods)			
Beans (without pods)			
Peas (with pods)			
Peas (without pods)			
Others			
(vii) STEM VEGETABLES (fresh)	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
Asparagus			
Cardoons			
Celery			
Fennel			
Globe artichokes			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Fenbutatin oxide	Triazophos	Diazinon
Leeks			
Rhubarb			
Others			
(viii) FUNGI	► M51 0,05 (*) ◀		► M64 0,01 (*) ◀
(a) <i>Cultivated mushrooms</i>			
(b) <i>Wild mushrooms</i>			
3. Pulses	► M51 0,05 (*) ◀	► M54 0,01 (*) ◀	► M64 0,01 (*) ◀
Beans			
Lentils			
Peas			
Others			
4. Oilseed	► M51 0,05 (*) ◀	► M54 0,01 (*) ◀	► M64 0,02 (*) ◀
Linseed			
Peanuts			
Poppy seeds			
Sesame seeds			
Sunflower seeds			
Rapeseed			
Soya bean			
Mustard seed			
Cotton seed			
Others			
5. Potatoes	► M51 0,05 (*) ◀	► M54 0,01 (*) ◀	► M64 0,01 (*) ◀
Early and ware potatoes			
6. Tea (Black tea processed from the leaves of <i>Camellia sinensis</i>)	► M51 0,1 (*) ◀	► M54 0,02 (*) ◀	► M64 0,02 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► M51 0,1 (*) ◀	► M54 0,02 (*) ◀	► M64 0,5 ◀

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Mecarbam		
1. Fruit, fresh, dried or uncooked preserved by freezing not containing added sugar; nuts			
(i) CITRUS FRUIT	2 (a)		
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)	0,05 (*)		
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			
(iii) POME FRUIT	0,05 (*)		
Apples			
Pears			
Quinces			
Other			
(iv) STONE FRUIT	0,05 (*)		
Apricots			
Cherries			
Peaches (including nectarines and similar hybrids)			
Plum			
Others			
(v) BERRIES AND SMALL FRUIT	0,05 (*)		
(a) <i>Table and wine grapes</i>			
(b) <i>Strawberries</i> (other than wild)			
(c) <i>Cane fruit</i> (other than wild)			
Blackberries			
Dewberries			
Loganberries			
Raspberries			
Others			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Mecarbam		
(d) <i>Other small fruit and berries</i> (other than wild) Bilberries (fruit of species <i>Vaccinium myrtilus</i>) Cranberries Currants (red, black and white) Gooseberries Others			
(e) <i>Wildberries and wild fruit</i>			
(vi) MISCELLANEOUS	0,05 (*)		
Avocados			
Bananas			
Dates			
Figs			
Kiwis			
Kumquats			
Litchis			
Mangoes			
Olives			
Passion fruit			
Pineapples			
Pomegranates			
Others			
2. Vegetables, fresh or uncooked, frozen or dry			
(i) ROOT AND TUBER	0,05 (*)		
Beetroot			
Carrots			
Celeriac			
Horseradish			
Jerusalem artichokes			
Parsnips			
Parsley root			
Radishes			
Salsify			
Sweet potatoes			
Swedes			
Turnips			
Yams			
Others			
(ii) BULB VEGETABLES	0,05 (*)		
Garlic			
Onions			
Shallots			
Spring onions			
Others			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Mecarbam		
(iii) FRUITING VEGETABLES	0,05 (*)		
(a) Solanacea			
Tomatoes			
Peppers			
Aubergines			
Others			
(b) Cucurbits — edible peel			
Cucumbers			
Gherkins			
Courgettes			
Others			
(c) Cucurbits — inedible peel			
Melons			
Squashes			
Watermelons			
Others			
(d) <i>Sweetcorn</i>			
(iv) BRASSICA VEGETABLES	0,05 (*)		
(a) Flowering brassica			
Broccoli			
Cauliflower			
Others			
(b) Head brassica			
Brussels sprouts			
Head cabbage			
Others			
(c) Leafy brassica			
Chinese cabbage			
Kale			
Others			
(d) <i>Kohlrabi</i>			
(v) LEAF VEGETABLES AND FRESH HERBS	0,05 (*)		
(a) Lettuce and similar			
Cress			
Lamb's lettuce			
Lettuce			
Scarole			
Others			
(b) Spinach and similar			
Spinach			
Beet leaves (chard)			
Others			
(c) <i>Watercress</i>			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Mecarbam		
(d) <i>Witloof</i>			
(e) Herbs			
Chervil			
Chives			
Parsley			
Celery leaves			
Others			
(vi) LEGUME VEGETABLES (fresh)	0,05 (*)		
Beans (with pods)			
Beans (without pods)			
Peas (with pods)			
Peas (without pods)			
Others			
(vii) STEM VEGETABLES (fresh)	0,05 (*)		
Asparagus			
Cardoons			
Celery			
Fennel			
Globe artichokes			
Leek			
Rhubarb			
Others			
(viii) FUNGI	0,05 (*)		
(a) <i>Cultivated mushrooms</i>			
(b) <i>Wild mushrooms</i>			
3. Pulses	0,05 (*)		
Beans			
Lentils			
Peas			
Others			
4. Oil seed	0,05 (*)		
Linseed			
Peanuts			
Poppy seeds			
Sesame seeds			
Sunflower seed			
Rapeseed			
Soya bean			
Mustard seed			
Cotton seed			
Others			
5. Potatoes	0,05 (*)		
Early and ware potatoes			

▼ **M5**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Mecarbam		
6. Tea (Black tea processed from the leaves of <i>Camellia sinensis</i>)	0,05 (*)		
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)		

(*) Indicates lower limit of analytical determination.

► **M42** (P) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 24 June 2009. ◀

(s) See Article 1 and Article 2 (2).

(a) (b) (c) (d) Should levels not be adopted by ► **M7** at the latest 1 July 2000 ◀, the following levels shall apply as indicated thereafter:

(a) 0,05 (*)

(b) 0,02 (*)

(c) 0,1 (*)

(d) 0,01 (*)

► **M34** (t) ► **M53** A temporary MRL of 0,2 mg/kg shall apply until 31 July 2009. ◀ ◀

▼ **M23**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Flupyrsulfuron-methyl	Azoxystrobin	Pymetrozine
1. Fruits, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,02 (P) (*)		
(i) CITRUS FRUIT		► M66 1 ◀	► M70 0,3 ◀
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)		► M66 0,1 (*) ◀	► M70 0,02 (*) ◀
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			
(iii) POME FRUIT		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
Apples			
Pears			
Quinces			
Others			
(iv) STONE FRUIT		► M66 0,05 (*) ◀	
Apricots			► M70 0,05 ◀
Cherries			
Peaches (including nectarines and similar hybrids)			► M70 0,05 ◀
Plums			
Others			► M70 0,02 ◀
(v) BERRIES AND SMALL FRUIT			
(a) Table and wine grapes		► M66 2 ◀	► M70 0,02 (*) ◀
Table grapes			
Wine grapes			
(b) Strawberries (other than wild)		► M66 2 ◀	► M70 0,5 ◀

▼ **M23**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Flupyr sulfuron-methyl	Azoxystrobin	Pymetrozine
(c) Cane fruit (other than wild)			
Blackberries		► M66 3 ◀	► M70 3 ◀
Dewberries			
Loganberries			
Raspberries		► M66 3 ◀	► M70 3 ◀
Others		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
(d) Other small fruit and berries (other than wild)		► M66 0,05 (*) ◀	
Bilberries			
Cranberries			
Currants (red, black and white)			► M70 0,1 ◀
Gooseberries			
Others			► M70 0,02 (*) ◀
(e) Wild berries and wild fruit		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
(vi) MISCELLANEOUS			► M70 0,02 (*) ◀
Avocados			
Bananas		► M66 2 ◀	
Dates			
Figs			
Kiwi			
Kumquats			
Litchis			
Mangoes		► M66 0,2 ◀	
Olives			
▼ M47			
Papaya		► M66 0,2 ◀	
▼ M23			
Passion fruit			
Pineapples			
Pomegranate			
Others		► M66 0,05 (*) ◀	
2. Vegetables, fresh or uncooked, frozen or dry	0,02 (p) (*)		
(i) ROOT AND TUBER VEGETABLES			► M70 0,02 (*) ◀
Beetroot			
Carrots		► M66 0,2 ◀	
Celeriac		► M66 0,3 ◀	
Horseradish		► M66 0,2 ◀	
Jerusalem artichokes			
Parsnips		► M66 0,2 ◀	
Parsley root		► M66 0,2 ◀	

▼ **M23**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Flupyrsulfuron-methyl	Azoxystrobin	Pymetrozine
Radishes		► M66 0,2 ◀	
Salsify		► M66 0,2 ◀	
Sweet potatoes			
Swedes			
Turnips			
Yam			
Others		► M66 0,05 (*) ◀	
(ii) BULB VEGETABLES			► M70 0,02 (*) ◀
Garlic			
Onions			
Shallots			
Spring onions		► M66 2 ◀	
Others		► M66 0,05 (*) ◀	
(iii) FRUITING VEGETABLES			
(a) Solanacea		► M66 2 ◀	
Tomatoes			► M70 0,5 ◀
Peppers			► M70 1 ◀
Aubergines			► M70 0,5 ◀
Okra			1
Others			► M70 0,02 (*) ◀
(b) Cucurbits — edible peel		► M66 1 ◀	► M70 0,5 ◀
Cucumbers			
Gherkins			
Courgettes			
Others			
(c) Cucurbits — inedible peel		► M66 0,5 ◀	► M70 0,2 ◀
Melons			
Squashes			
Watermelons			
Others			
(d) Sweetcorn		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
(iv) BRASSICA VEGETABLES			
(a) Flowering brassica		► M66 0,5 ◀	► M70 0,02 (*) ◀
Broccoli			
Cauliflower			
Others			

▼ **M70**▼ **M23**

▼ **M23**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Flupyrsulfuron-methyl	Azoxystrobin	Pymetrozine
b) Head brassica		► M66 0,3 ◀	
Brussels sprouts			► M70 0,05 ◀
Head cabbage			► M70 0,02 (*) ◀
Others			► M70 0,02 (*) ◀
(c) Leafy brassica		► M66 5 ◀	► M70 0,2 ◀
Chinese cabbage			
Kale			
Others			
(d) Kohlrabi		► M66 0,2 ◀	► M70 0,02 (*) ◀
(v) LEAF VEGETABLES AND FRESH HERBS			
(a) Lettuce and similar		► M66 3 ◀	► M70 2 ◀
Cress			
Lamb's lettuce			
Lettuce			
Scarole			
Others			
(b) Spinach and similar		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
Spinach			
Beet leaves (chard)			
Others			
(c) Water cress		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
(d) Witloof		► M66 0,2 ◀	► M70 0,02 (*) ◀
(e) Herbs		► M66 3 ◀	► M70 2 ◀
Chervil			
Chives			
Parsley			
Celery leaves			
Others			
(vi) LEGUME VEGETABLES (fresh)			
Beans (with pods)		► M66 1 ◀	► M70 2 ◀
Beans (without pods)		► M66 0,2 ◀	
Peas (with pods)		► M66 0,5 ◀	
Peas (without pods)		► M66 0,2 ◀	
Others		► M66 0,05 (*) ◀	► M70 1 ◀
(vii) STEM VEGETABLES (fresh)			► M70 0,02 (*) ◀
Asparagus			
Cardoons			
Celery		► M66 5 ◀	
Fennel		► M66 5 ◀	

▼ **M23**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)		
	Flupyr sulfuron-methyl	Azoxystrobin	Pymetrozine
Globe artichokes		► M66 1 ◀	
Leek		► M66 2 ◀	
Rhubarb			
Others		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
(viii) FUNGI		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
(a) Cultivated mushrooms			
(b) Wild mushrooms			
3. Pulses	0,02 (P) (*)	► M66 0,1 ◀	► M70 0,02 (*) ◀
Beans			
Lentils			
Peas			
Others			
4. Oil ls seeds	0,05 (P) (*)		
Linseed			
Peanuts			
Poppy seed			
Sesame seed			
Sunflower seed			
Rapeseed		► M66 0,5 ◀	
Soya bean		► M66 0,5 ◀	
Mustard seed			
Cotton seed			► M70 0,05 ◀
Others		► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
5. Potatoes	0,02 (P) (*)	► M66 0,05 (*) ◀	► M70 0,02 (*) ◀
Early potatoes			
Ware potatoes			
6. Tea (leaves and stems, dried, fermented or otherwise, of <i>Camellia sinensis</i>)	0,05 (P) (*)	► M66 0,1 (*) ◀	► M70 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (P) (*)	► M66 20 ◀	► M70 15 ◀

(P) Indicates provisional maximum residue level. For those agricultural products listed in Annex II to Directive 90/642/EEC where the maximum residue levels for flupyr sulfuron-methyl, pymetrozine and azoxystrobin are indicated as '(P)', this shall mean that they are provisional in accordance with the provisions of Article 4(1)(f) of Directive 91/414/EEC.

By 1 December 2005, provisional maximum residue levels for flupyr sulfuron-methyl and pymetrozine shall cease to be provisional and shall become definitive in the sense of Article 3 of Directive 90/624/EEC. For azoxystrobin this shall be 1 August 2003.

(*) Indicates lower limit of analytical determination.

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Dip-henylamine ◀
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts (I) CITRUS FRUIT Grapefruit Lemons Limes Mandarins (including clem-entines and other hybrids) Oranges Pomelos Others (II) TREE NUTS (shelled or unshelled) Almonds Brazil nuts Cashew nuts Chestnuts Coconuts Hazelnuts Macadamia Pecans Pine nuts Pistachios Walnuts Others	0,01 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,02 (*)	0,05 (*)	0,05 (*)	0,01 (*)	0,05 (*)	0,01 (*)	► M14 0,05 (*) ◀

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Dip-henylamine ◀
(III) POME FRUIT											
Apples											► M14 5 ◀
Pears											► M14 10 ◀
Quinces											
Others											
(IV) STONE FRUIT											
Apricots											
Cherries											
Peaches (including nectarines and similar hybrids)											
Plums											
Others											
(V) BERRIES AND SMALL FRUIT											
(a) Table and wine grapes											
Table grapes											
Wine grapes											
(b) Strawberries (other than wild)											
(c) Cane fruit (other than wild)											
Blackberries											
Dewberries											
Loganberries											

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Diphenylamine ◀
Raspberries											
Others											
(d) Other small fruit and berries (other than wild)											
Bilberries (fruit of species <i>Vaccinium myrtillus</i>)											
Cranberries											
Currants (red, black and white)											
Gooseberries											
Others											
(e) Wild berries and wild fruit											
(VI) MISCELLANEOUS											
Avocados											
Bananas											
Dates											
Figs											
Kiwi											
Kumquats											
Litchis											
Mangoes											
Olives											
Passion fruit											
Pineapples											
Pomegranate											
Others											

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Diphenylamine ◀
2. Vegetables, fresh or uncooked, frozen or dry	0,01 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,02 (*)	0,05 (*)	0,05 (*)	0,01 (*)	0,05 (*)	0,01 (*)	► M14 0,05 (*) ◀
(I) ROOT AND TUBER VEGETABLES											
Beetroot											
Carrots											
Celeriac											
Horseradish											
Jerusalem artichokes											
Parsnips											
Parsley root											
Radishes											
Salsify											
Sweet potatoes											
Swedes											
Turnips											
Yam											
Others											
(II) BULB VEGETABLES											
Garlic											
Onions											
Shallots											
Spring onions											
Others											

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Diphenylamine ◀
(III) FRUITING VEGETABLES											
(a) Solanacea											
Tomatoes											
Peppers											
Chili peppers											
Aubergines											
Others											
(b) Cucurbits — edible peel											
Cucumbers											
Gherkins											
Courgettes											
Others											
(c) Cucurbits — inedible peel											
Melons											
Squashes											
Watermelons											
Others											
(d) Sweet corn											
(IV) BRASSICA VEGETABLES											
(a) Flowering brassica											
Broccoli											
Cauliflower											
Others											

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenseide	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Diphenylamine ◀
(b) Head brassica											
Brussels sprouts											
Head cabbage											
Others											
(c) Leafy brassica											
Chinese cabbage											
Kale											
Others											
(d) Kohlrabi											
(V) LEAF VEGETABLES AND FRESH HERBS											
(a) Lettuce and similar											
Cress											
Lamb's lettuce											
Lettuce											
Scarole (broad-leaf endive)											
Others											
(b) Spinach and similar											
Spinach											
Beet leaves (chard)											
Others											
(c) Water cress											

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Diphenylamine ◀
(d) Witloof											
(e) Herbs											
Chervil											
Chives											
Parsley											
Celery leaves											
Others											
(VI) LEGUME VEGETABLES (fresh)											
Beans (with pods)											
Beans (without pods)											
Peas (with pods)											
Peas (without pods)											
Others											
(VII) STEM VEGETABLES (fresh)											
Asparagus											
Cardoons											
Celery											
Fennel											
Globe artichokes											
Leek											
Rhubarb											
Others											

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Diphenylamine ◀
(VIII) FUNGI											
(a) Cultivated mushrooms											
(b) Wild mushrooms											
3. Pulses	0,01 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,02 (*)	0,05 (*)	0,05 (*)	0,01 (*)	0,05 (*)	0,01 (*)	► M14 0,05 (*) ◀
Beans											
Lentils											
Peas											
Others											
4. Oil seed	0,01 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,02 (*)	0,05 (*)	0,05 (*)	0,01 (*)	0,05 (*)	0,01 (*)	► M14 0,05 (*) ◀
Linseed											
Peanuts											
Poppy seed											
Sesame seed											
Sunflower seed											
Rape seed											
Soya bean											
Mustard seed											
Cotton seed											
Others											
5. Potatoes	0,01 (*)	0,05 (*)	0,01 (*)	0,01 (*)	0,02 (*)	0,05 (*)	0,05 (*)	0,01 (*)	0,05 (*)	0,01 (*)	► M14 0,05 (*) ◀
Early potatoes											
Ware potatoes											

▼ **M11**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)										
	Methoxychlor	Barban	Aramite	Chlorfenson	Chlorobenzilate	Chlorbufam	Chloroxuron	Chlorbenside	Diallate	1,1-Dichloro-2,2-bis (4-ethyl-phenyl-) ethane	► M14 Diphenylamine ◀
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	► M14 0,05 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	► M14 0,05 (*) ◀

(*) Indicates lower limit of analytical determination.

▼ **M15**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Kresoxim methyl
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	
(i) CITRUS FRUIT	► M47 0,05 (*) ◀
Grapefruit	
Lemons	
Limes	
Mandarins (including clementines and other hybrids)	
Oranges	
Pomelos	
Others	
(ii) TREE NUTS (SHELLED OR UNSHELLED)	► M47 0,1 (*) ◀
Almonds	
Brazil nuts	
Cashew nuts	
Chestnuts	
Coconuts	
Hazelnuts	
Macadamia	
Pecans	
Pine nuts	
Pistachios	
Walnuts	
Others	
(iii) POME FRUIT	► M47 0,2 ◀
Apples	
Pears	
Quinces	
Others	
(iv) STONE FRUIT	► M47 0,05 (*) ◀
Apricots	
Cherries	
Peaches (including nectarines and similar hybrids)	
Plums	
Others	
(v) BERRIES AND SMALL FRUIT	
(a) Table and wine grapes	► M47 1 ◀
Table grapes	
Wine grapes	
(b) Strawberries (other than wild)	► M47 1 ◀
(c) Cane fruit (other than wild)	► M47 0,05 (*) ◀
Blackberries	
Dewberries	
Loganberries	
Raspberries	
Others	

▼ **M15**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Kresoxim methyl
(d) Other small fruit and berries (other than wild)	► M20 ——— ◀
Bilberries	
Cranberries	
Currants (red, black and white)	► M47 1 ◀
Gooseberries	► M47 1 ◀
Others	► M47 0,05 (*) ◀
(e) Wild berries and wild fruit	► M47 0,05 (*) ◀
(vi) MISCELLANEOUS	
Avocados	
Bananas	
Dates	
Figs	
Kiwi	
Kumquats	
Litchis	
Mangoes	
Olives	► M47 0,2 ◀
Passion fruit	
Pineapples	
Pomegranates	
Others	► M47 0,05 (*) ◀
2. Vegetables, fresh or uncooked, frozen or dry	
(i) ROOT AND TUBER VEGETABLES	► M47 0,05 (*) ◀
Beetroot	
Carrots	
Celeriac	
Horseradish	
Jerusalem artichokes	
Parsnips	
Parsley root	
Radishes	
Salsify	
Sweet potatoes	
Swedes	
Turnips	
Yam	
Others	
(ii) BULB VEGETABLES	► M47 0,05 (*) ◀
Garlic	
Onions	
Shallots	
Spring onions	
Others	

▼ **M15**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Kresoxim methyl
(iii) FRUITING VEGETABLES	
(a) Solanacea	
Tomatoes	► M47 0,5 ◀
Peppers	► M47 1 ◀
Aubergines	► M47 0,5 ◀
Others	► M47 0,05 (*) ◀
(b) Cucurbits — edible peel	► M47 0,05 (*) ◀
Cucumbers	
Gherkins	
Courgettes	
Others	
(c) Cucurbits — inedible peel	► M47 0,2 ◀
Melons	
Squashes	
Watermelons	
Others	
(d) Sweetcorn	► M47 0,05 (*) ◀
(iv) BRASSICA VEGETABLES	► M47 0,05 (*) ◀
(a) Flowering brassica	
Broccoli	
Cauliflower	
Others	
(b) Head brassica	
Brussels sprouts	
Head cabbage	
Others	
(c) Leafy brassica	
Chinese cabbage	
Kale	
Others	
(d) Kohlrabi	
(v) LEAF VEGETABLES AND FRESH HERBS	► M47 0,05 (*) ◀
(a) Lettuce and similar	
Cress	
Lamb's lettuce	
Lettuce	
Scarole	
Others	
(b) Spinach and similar	
Spinach	
Beet leaves (chard)	
Others	
(c) Watercress	
(d) Witloof	

▼ **M15**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Kresoxim methyl
(e) Herbs	
Chervil	
Chives	
Parsley	
Celery leaves	
Others	
(vi) LEGUME VEGETABLES (fresh)	► M47 0,05 (*) ◀
Beans (with pods)	
Beans (without pods)	
Peas (with pods)	
Peas (without pods)	
Others	
(vii) STEM VEGETABLES (fresh)	► M47 — ◀
Asparagus	
Cardoons	
Celery	
Fennel	
Globe artichokes	
Leeks	► M47 5 ◀
Rhubarb	
Others	► M47 0,05 (*) ◀
(viii) FUNGI	► M47 0,05 (*) ◀
(a) Cultivated mushrooms	
(b) Wild mushrooms	
3. Pulses	► M47 0,05 (*) ◀
Beans	
Lentils	
Peas	
Others	
4. Oil seeds	► M47 0,1 (*) ◀
Linseed	
Peanuts	
Poppy seeds	
Sesame seeds	
Sunflower seed	
Rape seed	
Soya bean	
Mustard seed	
Cotton seed	
Others	
5. Potatoes	► M47 0,05 (*) ◀
Early potatoes	
Ware potatoes	
6. Tea (leaves and stems dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	► M47 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► M47 0,1 (*) ◀

(*) Indicates lower limit of analytical determination.

▼ **M16**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Spiroxamine
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	
(i) CITRUS FRUIT	► M70 0,05 (*) ◀
Grapefruit	
Lemons	
Limes	
Mandarins (including clementines and other hybrids)	
Oranges	
Pomelos	
Others	
(ii) TREE NUTS (SHELLED OR UNSHELLED)	► M70 0,05 (*) ◀
Almonds	
Brazil nuts	
Cashew nuts	
Chestnuts	
Coconuts	
Hazelnuts	
Macadamia	
Pecans	
Pine nuts	
Pistachios	
Walnuts	
Others	
(iii) POME FRUIT	► M70 0,05 (*) ◀
Apples	
Pears	
Quinces	
Others	
(iv) STONE FRUIT	► M70 0,05 (*) ◀
Apricots	
Cherries	
Peaches (including nectarines and similar hybrids)	
Plums	
Others	
(v) BERRIES AND SMALL FRUIT	
(a) Table and wine grapes	► M70 1 ◀
Table grapes	
Wine grapes	
(b) Strawberries (other than wild)	► M70 0,05 (*) ◀
(c) Cane fruit (other than wild)	► M70 0,05 (*) ◀
Blackberries	
Dewberries	
Loganberries	
Raspberries	
Others	

▼ **M16**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Spiroxamine
(d) Other small fruit and berries (other than wild)	► M70 0,05 (*) ◀
Bilberries	
Cranberries	
Currants (red, black and white)	
Gooseberries	
Others	
(e) Wild berries and wild fruit	► M70 0,05 (*) ◀
(vi) MISCELLANEOUS	
Avocados	
Bananas	► M70 3 ◀
Dates	
Figs	
Kiwi fruit	
Kumquats	
Litchis	
Mangoes	
Olives	
Passion fruit	
Pineapples	
Pomegranates	
Others	► M70 0,05 (*) ◀
2. Vegetables, fresh or uncooked, frozen or dry	► M70 0,05 (*) ◀
(i) ROOT AND TUBER VEGETABLES	
Beetroot	
Carrots	
Celeriac	
Horseradish	
Jerusalem artichokes	
Parsnips	
Parsley root	
Radishes	
Salsify	
Sweet potatoes	
Swedes	
Turnips	
Yam	
Others	
(ii) BULB VEGETABLES	
Garlic	
Onions	
Shallots	
Spring onions	
Others	

▼ **M16**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Spiroxamine
<p>(iii) FRUITING VEGETABLES</p> <p>(a) Solanacea</p> <p>Tomatoes</p> <p>Peppers</p> <p>Aubergines</p> <p>Others</p> <p>(b) Cucurbits — edible peel</p> <p>Cucumbers</p> <p>Gherkins</p> <p>Courgettes</p> <p>Others</p> <p>(c) Cucurbits — inedible peel</p> <p>Melons</p> <p>Squashes</p> <p>Watermelons</p> <p>Others</p> <p>(d) Sweetcorn</p> <p>(iv) BRASSICA VEGETABLES</p> <p>(a) Flowering brassica</p> <p>Broccoli</p> <p>Cauliflower</p> <p>Others</p> <p>(b) Head brassica</p> <p>Brussels sprouts</p> <p>Head cabbage</p> <p>Others</p> <p>(c) Leafy brassica</p> <p>Chinese cabbage</p> <p>Kale</p> <p>Others</p> <p>(d) Kohlrabi</p> <p>(v) LEAF VEGETABLES AND FRESH HERBS</p> <p>(a) Lettuce and similar</p> <p>Cress</p> <p>Lamb's lettuce</p> <p>Lettuce</p> <p>Scarole</p> <p>Others</p> <p>(b) Spinach and similar</p> <p>Spinach</p> <p>Beet leaves (chard)</p> <p>Others</p>	

▼ **M16**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Spiroxamine
(c) Watercress	
(d) Witloof	
(e) Herbs	
Chervil	
Chives	
Parsley	
Celery leaves	
Others	
(vi) LEGUME VEGETABLES (fresh)	
Beans (with pods)	
Beans (without pods)	
Peas (with pods)	
Peas (without pods)	
Others	
(vii) STEM VEGETABLES (fresh)	
Asparagus	
Cardoons	
Celery	
Fennel	
Globe artichokes	
Leeks	
Rhubarb	
Others	
(viii) FUNGI	
(a) Cultivated mushrooms	
(b) Wild mushrooms	
3. Pulses	► M70 0,05 (*) ◀
Beans	
Lentils	
Peas	
Others	
4. Oil seeds	► M70 0,05 (*) ◀
Linseed	
Peanuts	
Poppy seeds	
Sesame seeds	
Sunflower seed	
Rape seed	
Soya bean	
Mustard seed	
Cotton seed	
Others	

▼ **M16**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)
	Spiroxamine
5. Potatoes Early potatoes Ware potatoes	► M70 0,05 (*) ◀
6. Tea (leaves and stems dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	► M70 0,1 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► M70 0,1 (*) ◀
(*) Indicates lower limit of analytical determination.	

▼M17

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)							
	Azinphosethyl	Chlozolinate	Dinoterb	DNOC	Monolinuron	Propham	Pyrazophos	Tecnazene
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	► M54 0,02 (*) ◀	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)
(I) CITRUS FRUIT								
Grapefruit								
Lemons								
Limes								
Mandarins (including clementines and other hybrids)								
Oranges								
Pomelos								
Others								
(II) TREE NUTS (shelled or unshelled)								
Almonds								
Brazil nuts								
Cashew nuts								
Chestnuts								
Coconuts								
Hazelnuts								
Macadamia								
Pecans								
Pine nuts								
Pistachios								
Walnuts								
Others								

▼M17

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)							
	Azinphosethyl	Chlozolinate	Dinoterb	DNOC	Monolinuron	Propham	Pyrazophos	Tecnazene
(III) POME FRUIT								
Apples								
Pears								
Quinces								
Others								
(IV) STONE FRUIT								
Apricots								
Cherries								
Peaches (including nectarines and similar hybrids)								
Plums								
Others								
(V) BERRIES AND SMALL FRUIT								
(a) Table and wine grapes								
Table grapes								
Wine grapes								
(b) Strawberries (other than wild)								
(c) Cane fruit (other than wild)								
Blackberries								
Dewberries								
Loganberries								
Raspberries								
Others								
(d) Other small fruit and berries (other than wild)								
Bilberries								
Cranberries								

▼M17

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)							
	Azinphosethyl	Chlozolinate	Dinoterb	DNOC	Monolinuron	Propham	Pyrazophos	Tecnazene
Currants (red, black and white)								
Gooseberries								
Others								
(e) Wild berries and wild fruit								
(VI) MISCELLANEOUS								
Avocados								
Bananas								
Dates								
Figs								
Kiwi								
Kumquats								
Litchis								
Mangoes								
Olives								
Passion fruit								
Pineapples								
Pomegranate								
Others								
2. Vegetables, fresh or uncooked, frozen or dry	► M54 0,02 (*) ◀	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)
(I) ROOT AND TUBER VEGETABLES								
Beetroot								
Carrots								
Celeriac								
Horseradish								
Jerusalem artichokes								
Parsnips								

▼M17

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)							
	Azinphosethyl	Chlozolinate	Dinoterb	DNOC	Monolinuron	Propham	Pyrazophos	Tecnazene
Parsley root								
Radishes								
Salsify								
Sweet potatoes								
Swedes								
Turnips								
Yam								
Others								
(II) BULB VEGETABLES								
Garlic								
Onions								
Shallots								
Spring onions								
Others								
(III) FRUITING VEGETABLES								
(a) Solanacea								
Tomatoes								
Peppers								
Aubergines								
Others								
(b) Cucurbits — edible peel								
Cucumbers								
Gherkins								
Courgettes								
Others								
(c) Cucurbits — inedible peel								
Melons								
Squashes								

▼M17

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)							
	Azinphosethyl	Chlozolinate	Dinoterb	DNOC	Monolinuron	Propham	Pyrazophos	Tecnazene
Watermelons								
Others								
(d) Sweet corn								
(IV) BRASSICA VEGETABLES								
(a) Flowering brassica								
Broccoli (including Calabrese)								
Cauliflower								
Others								
(b) Head brassica								
Brussels sprouts								
Head cabbage								
Others								
(c) Leafy brassica								
Chinese cabbage								
Kale								
Others								
(d) Kohlrabi								
(V) LEAF VEGETABLES AND FRESH HERBS								
(a) Lettuce and similar								
Cress								
Lamb's lettuce								
Lettuce								
Scarole (broad-leaf endive)								
Others								
(b) Spinach and similar								
Spinach								
Beet leaves (chard)								
Others								

▼M17

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)							
	Azinphosethyl	Chlozolinate	Dinoterb	DNOC	Monolinuron	Propham	Pyrazophos	Tecnazene
(c) Water cress								
(d) Witloof								
(e) Herbs								
Chervil								
Chives								
Parsley								
Celery leaves								
Others								
(VI) LEGUME VEGETABLES (fresh)								
Beans (with pods)								
Beans (without pods)								
Peas (with pods)								
Peas (without pods)								
Others								
(VII) STEM VEGETABLES (fresh)								
Asparagus								
Cardoons								
Celery								
Fennel								
Globe artichokes								
Leek								
Rhubarb								
Others								
(VIII) FUNGI								
(a) Cultivated mushrooms								
(b) Wild mushrooms								

▼ **M17**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)							
	Azinphosethyl	Chlozolinate	Dinoterb	DNOC	Monolinuron	Propham	Pyrazophos	Tecnazene
3. Pulses Beans Lentils Peas Others	► M54 0,02 (*) ◀	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)
4. Oilseeds Linseed Peanuts Poppy seed Sesame seed Sunflower seed Rape seed Soya bean Mustard seed Cotton seed Others	► M54 0,02 (*) ◀	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)
5. Potatoes Early potatoes Ware potatoes	► M54 0,02 (*) ◀	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)	0,05 (*)
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	► M54 0,05 (*) ◀	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	► M54 0,05 (*) ◀	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)	0,1 (*)

(*) Indicates lower limit of analytical determination.

▼ **M19**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Azimsulfuron	Prohexadione (prohexadione and its salts expressed as prohexadione)
1. Fruits, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,02 (P) (*)	0,05 (P) (*)
(i) CITRUS FRUIT		
Grapefruit		
Lemons		
Limes		
Mandarins (including clementines and other hybrids)		
Oranges		
Pomelos		
Others		
(ii) TREE NUTS (shelled or unshelled)		
Almonds		
Brazil nuts		
Cashew nuts		
Chestnuts		
Coconuts		
Hazelnuts		
Macadamia		
Pecans		
Pine nuts		
Pistachios		
Walnuts		
Others		
(iii) POME FRUIT		
Apples		
Pears		
Quinces		
Others		
(iv) STONE FRUIT		
Apricots		
Cherries		
Peaches (including nectarines and similar hybrids)		
Plums		
Others		
(v) BERRIES AND SMALL FRUIT		
(a) Table and wine grapes		
Table grapes		
Wine grapes		
(b) Strawberries (other than wild)		

▼ **M19**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Azimsulfuron	Prohexadione (prohexadione and its salts expressed as prohexadione)
(c) Cane fruit (other than wild)		
Blackberries		
Dewberries		
Loganberries		
Raspberries		
Others		
(d) Other small fruit and berries (other than wild)		
Bilberries		
Cranberries		
Currants (red, black and white)		
Gooseberries		
Others		
(e) Wild berries and wild fruit		
(vi) MISCELLANEOUS		
Avocados		
Bananas		
Dates		
Figs		
Kiwi		
Kumquats		
Litchis		
Mangoes		
Olives		
Passion fruit		
Pineapples		
Pomegranate		
Others		
2. Vegetables, fresh or uncooked, frozen or dry	0,02 (P) (*)	0,05 (P) (*)
(i) ROOT AND TUBER VEGETABLES		
Beetroot		
Carrots		
Celeriac		
Horseradish		
Jerusalem artichokes		
Parsnips		
Parsley root		
Radishes		
Salsify		
Sweet potatoes		
Swedes		
Turnips		
Yam		
Others		

▼ **M19**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Azimsulfuron	Prohexadione (prohexadione and its salts expressed as prohexadione)
(ii) BULB VEGETABLES		
Garlic		
Onions		
Shallots		
Spring onions		
Others		
(iii) FRUITING VEGETABLES		
(a) Solanacea		
Tomatoes		
Peppers		
Aubergines		
Others		
(b) Cucurbits — edible peel		
Cucumbers		
Gherkins		
Courgettes		
Others		
(c) Cucurbits — inedible peel		
Melons		
Squashes		
Watermelons		
Others		
(d) Sweetcorn		
(iv) BRASSICA VEGETABLES		
(a) Flowering brassica		
Broccoli		
Cauliflower		
Others		
(b) Head brassica		
Brussels sprouts		
Head cabbage		
Others		
(c) Leafy brassica		
Chinese cabbage		
Kale		
Others		
(d) Kohlrabi		
(v) LEAF VEGETABLES AND FRESH HERBS		
(a) Lettuce and similar		
Cress		
Lamb's lettuce		
Lettuce		
Scarole		
Others		

▼ **M19**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Azimsulfuron	Prohexadione (prohexadione and its salts expressed as prohexadione)
(b) Spinach and similar Spinach Beet leaves (chard) Others		
(c) Watercress		
(d) Witloof		
(e) Herbs Chervil Chives Parsley Celery leaves Others		
(vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others		
(vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek Rhubarb Others		
(viii) FUNGI (a) Cultivated mushrooms (b) Wild mushrooms		
3. Pulses	0,02 (P) (*)	0,05 (P) (*)
Beans		
Lentils		
Peas		
Others		
4. Oils seeds	0,1 (P) (*)	0,1 (P) (*)
Linseed		
Peanuts		
Poppy seeds		
Sesame seeds		
Sunflower seed		
Rapeseed		
Soya bean		
Mustard seed		

▼ **M19**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Azimsulfuron	Prohexadione (prohexadione and its salts expressed as prohexadione)
Cotton seed		
Others		
5. Potatoes	0,02 (P) (*)	0,05 (P) (*)
Early potatoes		
Ware potatoes		
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (P) (*)	0,1 (P) (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (P) (*)	0,1 (P) (*)

(*) Indicates lower limit of analytical determination.

(P) Indicates provisional maximum residue level established in accordance with Article 4(1)(f) of Directive 91/414/EEC: all provisional maximum residue levels for these pesticide residues will be treated as definitive in accordance with Article 10 of the Directive with effect from four years after the entry into force of this Directive.

▼ **M21**

Groups and examples of individual products to which the maximum residue levels apply	Pesticide residues and maximum residue levels (mg/kg)
	Fluroxypyr including its esters expressed as fluroxypyr
<p>1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts</p> <p>(i) CITRUS FRUIT</p> <p>Grapefruit</p> <p>Lemons</p> <p>Limes</p> <p>Mandarins (including clementines and other hybrids)</p> <p>Oranges</p> <p>Pomelos</p> <p>Others</p> <p>(ii) TREE NUTS (shelled or unshelled)</p> <p>Almonds</p> <p>Brazil nuts</p> <p>Cashew nuts</p> <p>Chestnuts</p> <p>Coconuts</p> <p>Hazelnuts</p> <p>Macadamia</p> <p>Pecans</p> <p>Pine nuts</p> <p>Pistachios</p> <p>Walnuts</p> <p>Others</p> <p>(iii) POME FRUIT</p> <p>Apples</p> <p>Pears</p> <p>Quinces</p> <p>Others</p> <p>(iv) STONE FRUIT</p> <p>Apricots</p> <p>Cherries</p> <p>Peaches (including nectarines and similar hybrids)</p> <p>Plums</p> <p>Others</p> <p>(v) BERRIES AND SMALL FRUIT</p> <p>(a) Table and wine grapes</p> <p>Table grapes</p> <p>Wine grapes</p> <p>(b) Strawberries (other than wild)</p>	0,05 (*) (P)

▼ **M21**

Groups and examples of individual products to which the maximum residue levels apply	Pesticide residues and maximum residue levels (mg/kg)
	Fluroxypyr including its esters expressed as fluroxypyr
(c) Cane fruit (other than wild) Blackberries Dewberries Loganberries Raspberries Others (d) Other small fruit and berries (other than wild) Bilberries Cranberries Currants (red, black and white) Gooseberries Others (e) Wild berries and wild fruit (vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes Olives Passion fruit Pineapples Pomegranates Others	
2. Vegetables, fresh or uncooked, frozen or dry	0,05 (*) (P)
(i) ROOT AND TUBER VEGETABLES Beetroot Carrots Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips Yam Others	

▼ **M21**

Groups and examples of individual products to which the maximum residue levels apply	Pesticide residues and maximum residue levels (mg/kg)
	Fluroxypyr including its esters expressed as fluroxypyr
<ul style="list-style-type: none"> (ii) BULB VEGETABLES <ul style="list-style-type: none"> Garlic Onions Shallots Spring onions Others (iii) FRUITING VEGETABLES <ul style="list-style-type: none"> (a) Solanacea <ul style="list-style-type: none"> Tomatoes Peppers Aubergines Others (b) Cucurbits — edible peel <ul style="list-style-type: none"> Cucumbers Gherkins Courgettes Others (c) Cucurbits — inedible peel <ul style="list-style-type: none"> Melons Squashes Watermelons Others (d) Sweetcorn (iv) BRASSICA VEGETABLES <ul style="list-style-type: none"> (a) Flowering brassica <ul style="list-style-type: none"> Broccoli Cauliflower Others (b) Head brassica <ul style="list-style-type: none"> Brussels sprouts Head cabbage Others (c) Leafy brassica <ul style="list-style-type: none"> Chinese cabbage Kale Others (d) Kohlrabi (v) LEAF VEGETABLES AND FRESH HERBS <ul style="list-style-type: none"> (a) Lettuce and similar <ul style="list-style-type: none"> Cress Lamb's lettuce Lettuce Scarole Others 	

▼ **M21**

Groups and examples of individual products to which the maximum residue levels apply	Pesticide residues and maximum residue levels (mg/kg)
	Fluroxypyr including its esters expressed as fluroxypyr
(b) Spinach and similar Spinach Beet leaves (chard) Others (c) Watercress (d) Witloof (e) Herbs Chervil Chives Parsley Celery leaves Others (vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others (vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek Rhubarb Others (viii) FUNGI (a) Cultivated mushrooms (b) Wild mushrooms	
3. Pulses Beans Lentils Peas Others	0,05 (*) (P)
4. Oil seeds Linseed Peanuts Poppy seeds Sesame seeds Sunflower seeds Rape seed	0,05 (*) (P)

▼ **M21**

Groups and examples of individual products to which the maximum residue levels apply	Pesticide residues and maximum residue levels (mg/kg)
	Fluroxypyr including its esters expressed as fluroxypyr
Soya bean Mustard seed Cotton seed Others	
5. Potatoes Early potatoes Ware potatoes	0,05 (*) (P)
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*) (P)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) (P)

(*) Indicates lower limit of analytical determination.

(P) Indicates provisional maximum residue level established in accordance with Article 4(1)(f) of Directive 91/414/EEC: all provisional maximum residue levels for these pesticide residues will be treated as definitive in accordance with Article 10 of the Directive with effect from four years after the entry into force of this Directive.

▼ **M24**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Bentazone (sum of bentazone and the conjugates of 6-OH- and 8-OH-bentazone expressed as bentazone)	Pyridate (sum of pyridate, its hydrolysis product CL 9673 (6-chloro-4-hydroxy-3-phenylpyridazin) and hydrolysable conjugates of CL 9673 expressed as pyridate)
1. Fruits, fresh, dried or uncooked, preserved by freezing, not containing added sugar: nuts	0,1 (P) (*)	0,05 (P) (*)
(i) CITRUS FRUIT		
Grapefruit		
Lemons		
Limes		
Mandarins (including clementines and other hybrids)		
Oranges		
Pomelos		
Others		
(ii) TREE NUTS (shelled or unshelled)		
Almonds		
Brazil nuts		
Cashew nuts		
Chestnuts		
Coconuts		
Hazelnuts		
Macadamia		
Pecans		
Pine nuts		
Pistachios		
Walnuts		
Others		
(iii) POME FRUIT		
Apples		
Pears		
Quinces		
Others		
(iv) STONE FRUIT		
Apricots		
Cherries		
Peaches (including nectarines and similar hybrids)		
Plums		
Others		
(v) BERRIES AND SMALL FRUIT		
(a) Table and wine grapes		
Table grapes		
Wine grapes		

▼ **M24**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Bentazone (sum of bentazone and the conjugates of 6-OH- and 8-OH-bentazone expressed as bentazone)	Pyridate (sum of pyridate, its hydrolysis product CL 9673 (6-chloro-4-hydroxy-3-phenylpyridazin) and hydrolysable conjugates of CL 9673 expressed as pyridate)
(b) Strawberries (other than wild)		
(c) Cane fruit (other than wild)		
Blackberries		
Dewberries		
Loganberries		
Raspberries		
Others		
(d) Other small fruit and berries (other than wild)		
Bilberries		
Cranberries		
Currants (red, black and white)		
Gooseberries		
Others		
(e) Wild berries and wild fruit		
(vi) MISCELLANEOUS		
Avocados		
Banans		
Dates		
Figs		
Kiwi		
Kumquats		
Litchis		
Mangoes		
Olives		
Passion fruit		
Pineapples		
Pomegranate		
Others		
2. Vegetables, fresh or uncooked, frozen or dry		
(i) ROOT AND TUBER VEGETABLES	0,1 (P) (*)	0,05 (P) (*)
Beetroot		
Carrots		
Celeriac		
Horseradish		
Jerusalem artichokes		
Parsnips		
Parsley root		
Radishes		
Salsify		

▼ M24

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Bentazone (sum of bentazone and the conjugates of 6-OH- and 8-OH-bentazone expressed as bentazone)	Pyridate (sum of pyridate, its hydrolysis product CL 9673 (6-chloro-4-hydroxy-3-phenylpyridazin) and hydrolysable conjugates of CL 9673 expressed as pyridate)
Sweet potatoes		
Swedes		
Turnips		
Yam		
Others		
(ii) BULB VEGETABLES	0,1 (P) (*)	0,05 (P) (*)
Garlic		
Onions		
Shallots		
Spring onions		
Others		
(iii) FRUITING VEGETABLES	0,1 (P) (*)	0,05 (P) (*)
(a) Solanacea		
Tomatoes		
Peppers		
Aubergines		
Others		
(b) Cucurbits — edible peel		
Cucumbers		
Gherkins		
Courgettes		
Others		
(c) Cucurbits — inedible peel		
Melons		
Squashes		
Watermelons		
Others		
(d) Sweetcorn		
(iv) BRASSICA VEGETABLES	0,1 (P) (*)	
(a) Flowering brassica		0,05 (P) (*)
Broccoli		
Cauliflower		
Others		
(b) Head brassica		0,05 (P) (*)
Brussels sprouts		
Head cabbage		
Others		

▼ M24

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Bentazone (sum of bentazone and the conjugates of 6-OH- and 8-OH-bentazone expressed as bentazone)	Pyridate (sum of pyridate, its hydrolysis product CL 9673 (6-chloro-4-hydroxy-3-phenylpyridazin) and hydrolysable conjugates of CL 9673 expressed as pyridate)
(c) Leafy brassica		
Chinese cabbage		
Kale		0,2 (P)
Others		0,05 (P) (*)
(d) Kohlrabi		0,05 (P) (*)
(v) LEAF VEGETABLES AND FRESH HERBS		0,05 (P) (*)
(a) Lettuce and similar	0,1 (P) (*)	
Cress		
Lamb's lettuce		
Lettuce		
Scarole		
Others		
(b) Spinach and similar	0,1 (P) (*)	
Spinach		
Beet leaves (chard)		
Others		
(c) Water cress	0,1 (P) (*)	
(d) Witloof	0,1 (P) (*)	
(e) Herbs	0,1 (P) (*)	
Chervil		
Chives		
Parsley		
Celery leaves		
Others		
(vi) LEGUME VEGETABLES (fresh)		0,05 (P) (*)
Beans (with pods)		
Beans (without pods)		
Peas (with pods)	0,5 (P)	
Peas (without pods)	0,2 (P)	
Others	0,1 (P) (*)	
(vii) STEM VEGETABLES (fresh)	0,1 (P) (*)	
Asparagus		
Cardoons		
Celery		
Fennel		
Globe artichokes		
Leek		1 (P)
Rhubarb		
Others		0,05 (P) (*)

▼ **M24**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)	
	Bentazone (sum of bentazone and the conjugates of 6-OH- and 8-OH-bentazone expressed as bentazone)	Pyridate (sum of pyridate, its hydrolysis product CL 9673 (6-chloro-4-hydroxy-3-phenylpyridazin) and hydrolysable conjugates of CL 9673 expressed as pyridate)
(viii) FUNGI	0,1 (P) (*)	0,05 (P) (*)
(a) Cultivated mushrooms		
(b) Wild mushrooms		
3. Pulses	0,1 (P) (*)	0,05 (P) (*)
Beans		
Lentils		
Peas		
Others		
4. Oils seeds		0,05 (P) (*)
Linseed		
Peanuts		
Poppy seed		
Sesame seed		
Sunflower seed		
Rapeseed		
Soya bean	0,1 (P)	
Mustard seed		
Cotton seed		
Others	0,1 (P) (*)	
5. Potatoes	0,1 (P) (*)	0,05 (P) (*)
Early potatoes		
Ware potatoes		
6. Tea (leaves and stems, dried, fermented or otherwise, of <i>Camellia sinensis</i>)	0,1 (P)	0,1 (P) (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (P) (*)	0,1 (P) (*)

► **C5** (P) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 4 years from date of coming into force of the Directive introducing this amendment. ◀

(*) Indicates lower limit of analytical determination.

▼ **M25**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Lindane	Quintozene (sum of quintozene, and pentachloroaniline expressed as quintozene)	Permethrin (sum of isomers)	Parathion
1. Fruits, fresh, dried or uncooked, preserved by freezing, not containing added sugar: nuts	0,01 (*)	0,02 (*)	0,05 (*)	
(i) CITRUS FRUIT				
Grapefruit				
Lemons				
Limes				
Mandarins (including clementines and other hybrids)				
Oranges				
Pomelos				
Others				
(ii) TREE NUTS (shelled or unshelled)				
Almonds				
Brazil nuts				
Cashew nuts				
Chestnuts				
Coconuts				
Hazelnuts				
Macadamia				
Pecans				
Pine nuts				
Pistachios				
Walnuts				
Others				
(iii) POME FRUIT				
Apples				
Pears				
Quinces				
Others				
(iv) STONE FRUIT				
Apricots				
Cherries				
Peaches (including nectarines and similar hybrids)				
Plums				
Others				
(v) BERRIES AND SMALL FRUIT				
(a) Table and wine grapes				
Table grapes				
Wine grapes				

▼ M25

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Lindane	Quintozene (sum of quintozene, and pentachloroaniline expressed as quintozene)	Permethrin (sum of isomers)	Parathion
(b) Strawberries (other than wild)				
(c) Cane fruit (other than wild)				
Blackberries				
Dewberries				
Loganberries				
Raspberries				
Others				
(d) Other small fruit and berries (other than wild)				
Bilberries				
Cranberries				
Currants (red, black and white)				
Gooseberries				
Others				
(e) Wild berries and wild fruit				
(vi) MISCELLANEOUS				
Avocados				
Banans				
Dates				
Figs				
Kiwi				
Kumquats				
Litchis				
Mangoes				
Olives				
Passion fruit				
Pineapples				
Pomegranate				
Others				
2. Vegetables, fresh or uncooked, frozen or dry	0,01 (*)	0,02 (*)	0,05 (*)	
(i) ROOT AND TUBER VEGETABLES				
Beetroot				
Carrots				
Celeriac				
Horseradish				
Jerusalem artichokes				
Parsnips				
Parsley root				
Radishes				

▼ **M25**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Lindane	Quintozene (sum of quintozene, and pentachloroaniline expressed as quintozene)	Permethrin (sum of isomers)	Parathion
Salsify				
Sweet potatoes				
Swedes				
Turnips				
Yam				
Others				
(ii) BULB VEGETABLES				
Garlic				
Onions				
Shallots				
Spring onions				
Others				
(iii) FRUITING VEGETABLES				
(a) Solanacea				
Tomatoes				
Peppers				
Aubergines				
Others				
(b) Cucurbits — edible peel				
Cucumbers				
Gherkins				
Courgettes				
Others				
(c) Cucurbits — inedible peel				
Melons				
Squashes				
Watermelons				
Others				
(d) Sweetcorn				
(iv) BRASSICA VEGETABLES				
(a) Flowering brassica				
Broccoli				
Cauliflower				
Others				
(b) Head brassica				
Brussels sprouts				
Head cabbage				
Others				

▼ M25

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Lindane	Quintozene (sum of quintozene, and pentachloroaniline expressed as quintozene)	Permethrin (sum of isomers)	Parathion
(c) Leafy brassica Chinese cabbage Kale Others				
(d) Kohlrabi				
(v) LEAF VEGETABLES AND FRESH HERBS				
(a) Lettuce and similar Cress Lamb's lettuce Lettuce Scarole (broad-leaf endive) Others				
(b) Spinach and similar Spinach Beet leaves (chard) Others				
(c) Watercress				
(d) Witloof				
(e) Herbs Chervil Chives Parsley Celery leaves Others				
(vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others				
(vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek Rhubarb Others				

▼ **M25**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Lindane	Quintozene (sum of quintozene, and pentachloroaniline expressed as quintozene)	Permethrin (sum of isomers)	Parathion
(viii) FUNGI				
(a) Cultivated mushrooms				
(b) Wild mushrooms				
3. Pulses	0,01 (*)	0,02 (*)	0,05 (*)	0,05 (*)
Beans				
Lentils				
Peas				
Others				
4. Oils seeds	0,01 (*)		0,05 (*)	
Linseed				
Peanuts		0,05 (#)		
Poppy seed				
Sesame seed				
Sunflower seed				
Rapeseed				
Soya bean				
Mustard seed				
Cotton seed				
Others		0,02 (*)		
5. Potatoes	0,01 (*)	0,02 (*)	0,05 (*)	
Early potatoes				
Ware potatoes				
6. Tea (leaves and stems, dried, fermented or otherwise, of <i>Camellia sinensis</i>)	0,05 (*)	0,05 (*)	0,1 (*)	
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (*)	0,05 (*)	0,1 (*)	

(*) Indicates lower limit of analytical determination.

Indicates that the MRL is based on Codex MRL.

▼ **M26**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)		
	Formothion	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts		0,02 (*)	
(i) CITRUS FRUIT	0,02 (*)		► <u>M70</u> 0,02 (*) ◀
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)	0,05 (*)		► <u>M70</u> 0,05 (*) ◀
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			
(iii) POME FRUIT	0,02 (*)		► <u>M70</u> 0,02 (*) ◀
Apples			
Pears			
Quinces			
Others			
(iv) STONE FRUIT	0,02 (*)		
Apricots			
Cherries			► <u>M70</u> 1 ◀
Peaches (including nectarines and similar hybrids)			
Plums			
Others			► <u>M70</u> 0,02 (*) ◀

▼ **M26**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)		
	Formothion	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
(v) BERRIES AND SMALL FRUIT	0,02 (*)		► M70 0,02 (*) ◀
(a) Table and wine grapes			
Table grapes			
Wine grapes			
(b) Strawberries (other than wild)			
(c) Cane fruit (other than wild)			
Blackberries			
Dewberries			
Loganberries			
Raspberries			
Others			
(d) Other small fruit and berries (other than wild)			
Bilberries			
Cranberries			
Currants (red-, black- and white-)			
Gooseberries			
Others			
(e) Wild berries and wild fruit			
(vi) MISCELLANEOUS	0,02 (*)		
Avocados			
Bananas			
Dates			
Figs			
Kiwi			
Kumquats			
Litchis			
Mangoes			
Olives			► M70 2 ◀
Passion fruit			
Pineapples			
Pomegranate			
Others			► M70 0,02 (*) ◀
2. Vegetables, fresh or uncooked, frozen or dry	0,02 (*)		
(i) ROOT AND TUBER VEGETABLES		0,02 (*)	
Beetroot			
Carrots			
Celeriac			► M70 0,1 ◀

▼ **M26**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)		
	Formothion	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
Horseradish			
Jerusalem artichokes			
Parsnips			
Parsley root			
Radishes			
Salsify			
Sweet potatoes			
Swedes			
Turnips			
Yam			
Others			► M70 0,02 (*) ◀
(ii) BULB VEGETABLES		0,02 (*)	
Garlic			
Onions			
Shallots			
Spring onions			► M70 2 ◀
Others			
(iii) FRUITING VEGETABLES		0,02 (*)	► M70 0,02 (*) ◀
(a) Solanacea			
Tomatoes			
Peppers			
Aubergines			
Others			
(b) Cucurbits — edible peel			
Cucumbers			
Gherkins			
Courgettes			
Others			
(c) Cucurbits — inedible peel			
Melons			
Squashes			
Watermelons			
Others			
(d) Sweet corn			

▼ **M26**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)		
	Formothion	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
(iv) BRASSICA VEGETABLES			
(a) Flowering brassica		0,02 (*)	
Broccoli (including Calabrese)			
Cauliflower			► <u>M70</u> 0,2 ◀
Others			► <u>M70</u> 0,02 (*) ◀
(b) Head brassica			
Brussels sprouts		0,05	► <u>M70</u> 0,3 ◀
Head cabbage		0,05	► <u>M70</u> 1 ◀
Others		0,02 (*)	► <u>M70</u> 0,02 (*) ◀
(c) Leafy brassica		0,02 (*)	► <u>M70</u> 0,02 (*) ◀
Chinese cabbage			
Kale			
Others			
(d) Kohlrabi		0,05	► <u>M70</u> 0,02 (*) ◀
(v) LEAF VEGETABLES AND FRESH HERBS			
(a) Lettuce and similar		0,05	
Cress			
Lamb's lettuce			
Lettuce			► <u>M70</u> 0,5 ◀
Scarole (broad-leaf endive)			
Others			► <u>M70</u> 0,02 (*) ◀
(b) Spinach and similar		0,02 (*)	► <u>M70</u> 0,02 (*) ◀
Spinach			
Beet leaves (chard)			
Others			
(c) Water cress		0,02 (*)	► <u>M70</u> 0,02 (*) ◀
(d) Witloof		0,02 (*)	► <u>M70</u> 0,02 (*) ◀
(e) Herbs		0,02 (*)	► <u>M70</u> 0,02 (*) ◀
Chervil			
Chives			
Parsley			
Celery leaves			
Others			

▼ **M26**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)		
	Formothion	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
(vi) LEGUME VEGETABLES (fresh)		0,02 (*)	
Beans (with pods)			
Beans (without pods)			
Peas (with pods)			► M70 1 ◀
Peas (without pods)			
Others			► M70 0,02 (*) ◀
(vii) STEM VEGETABLES (fresh)		0,02 (*)	► M70 0,02 (*) ◀
Asparagus			
Cardoons			
Celery			
Fennel			
Globe artichokes			
Leek			
Rhubarb			
Others			
(viii) FUNGI		0,02 (*)	► M70 0,02 (*) ◀
Cultivated mushrooms			
Wild mushrooms			
3. Pulses	0,02 (*)	0,02 (*)	► M70 0,02 (*) ◀
Beans			
Lentils			
Peas			
Others			
4. Oilseeds	0,05 (*)	0,05 (*)	► M70 0,05 (*) ◀
Linseed			
Peanuts			
Poppy seed			
Sesame seed			
Sunflower seed			
Rape seed			
Soya bean			
Mustard seed			
Cotton seed			
Others			

▼ **M26**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)		
	Formothion	Oxydemeton-methyl (sum of oxydemeton-methyl and demeton-S-methylsulfone expressed as oxydemeton-methyl)	Dimethoate (sum of dimethoate and omethoate expressed as dimethoate)
5. Potatoes Early potatoes Ware potatoes	0,02 (*)	0,02 (*)	► M70 0,02 (*) ◀
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,05 (*)	0,05 (*)	► M70 0,05 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (*)	0,05 (*)	► M70 0,05 (*) ◀

(*) Indicates lower limit of analytical determination.

▼ **M27**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)
	Metsulfuron methyl
<p>1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts</p> <p>(i) CITRUS FRUIT</p> <ul style="list-style-type: none"> Grapefruit Lemons Limes Mandarins (including clementines and other hybrids) Oranges Pomelos Others <p>(ii) TREE NUTS (shelled or unshelled)</p> <ul style="list-style-type: none"> Almonds Brazil nuts Cashew nuts Chestnuts Coconuts Hazelnuts Macadamia Pecans Pine nuts Pistachios Walnuts Others <p>(iii) POME FRUIT</p> <ul style="list-style-type: none"> Apples Pears Quinces Others <p>(iv) STONE FRUIT</p> <ul style="list-style-type: none"> Apricots Cherries Peaches (including nectarines and similar hybrids) Plums Others <p>(v) BERRIES AND SMALL FRUIT</p> <ul style="list-style-type: none"> (a) Table and wine grapes <ul style="list-style-type: none"> Table grapes Wine grapes (b) Strawberries (other than wild) (c) Cane fruit (other than wild) <ul style="list-style-type: none"> Blackberries Dewberries Loganberries Raspberries Others 	0,05 (*) (a)

▼ **M27**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)
	Metsulfuron methyl
(d) Other small fruit and berries (other than wild) Bilberries Cranberries Currants (red, black and white) Gooseberries Others (e) Wild berries and wild fruit (vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes Olives Passion fruit Pineapples Pomegranate Others	
2. Vegetables, fresh or uncooked, frozen or dry (i) ROOT AND TUBER VEGETABLES Beetroot Carrots Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips Yam Others (ii) BULB VEGETABLES Garlic Onions Shallots Spring onions Others	0,05 (*) ^(a)

▼ **M27**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)
	Metsulfuron methyl
<ul style="list-style-type: none"> (iii) FRUITING VEGETABLES <ul style="list-style-type: none"> (a) Solanacea <ul style="list-style-type: none"> Tomatoes Peppers Aubergines Others (b) Cucurbits — edible peel <ul style="list-style-type: none"> Cucumbers Gherkins Courgettes Others (c) Cucurbits — inedible peel <ul style="list-style-type: none"> Melons Squashes Watermelons Others (d) Sweet corn (iv) BRASSICA VEGETABLES <ul style="list-style-type: none"> (a) Flowering brassica <ul style="list-style-type: none"> Broccoli Cauliflower Others (b) Head brassica <ul style="list-style-type: none"> Brussels sprouts Head cabbage Others (c) Leafy brassica <ul style="list-style-type: none"> Chinese cabbage Kale Others (d) Kohlrabi (v) LEAF VEGETABLES AND FRESH HERBS <ul style="list-style-type: none"> (a) Lettuce and similar <ul style="list-style-type: none"> Cress Lamb's lettuce Lettuce Scarole Others (b) Spinach and similar <ul style="list-style-type: none"> Spinach Beet leaves (chard) Others (c) Watercress (d) Witloof 	

▼ **M27**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)
	Metsulfuron methyl
(e) Herbs Chervil Chives Parsley Celery leaves Others	
(vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others	
(vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek Rhubarb Others	
(viii) FUNGI (a) Cultivated mushrooms (b) Wild mushrooms	
3. Pulses Beans Lentils Peas Others	0,05 (*) ^(a)
4. Oil seeds Linseed Peanuts Poppy seeds Sesame seeds Sunflower seed Rape seed Soya bean Mustard seed Cotton seed Others	0,1 (*) ^(a)
5. Potatoes Early potatoes Ware potatoes	0,05 (*) ^(a)

▼ **M27**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)
	Metsulfuron methyl
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*) ^(a)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) ^(a)

(*) Indicates lower limit of analytical determination

^(a) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 4 years from date of coming into force of the Directive introducing this amendment.

▼ **M28**

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacryfos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)	
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts					► M38 — ◀		► M70 0,05 (*) ◀		0,05 (*)	► M32 — ◀	0,05 (*)				0,05 (*)				
(i) CITRUS FRUIT	► M57 0,01 (*) ◀	0,2	► M57 0,1 ◀	0,05 (*)	► M38 2 (i) ◀	► M70 0,5 ◀		► M53 0,05 (*) ◀		► M32 0,02 (*) ◀		► M53 3 ◀	► M58 0,05 (*) ◀	10		0,1 (*)	0,05 (*)	► M51 0,1 (*) ◀	
Grapefruit																			
Lemons																			
Limes																			
Mandarins (including clementines and other hybrids)																			
Oranges																			
Pomelos																			
Others																			
(ii) TREE NUTS (shelled or unshelled)	► M57 0,01 (*) ◀	0,1 (*)	► M57 0,05 (*) ◀	0,1 (*)	► M38 0,05 (*) ◀	► M70 0,05 (*) ◀		► M53 0,05 (*) ◀		► M32 0,05 (*) ◀		► M53 0,05 (*) ◀	► M58 0,05 (*) ◀	0,1 (*)		0,2 (*)	0,1 (*)	► M51 0,2 (*) ◀	
Almonds																			
Brazil nuts																			
Cashew nuts																			

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Chestnuts																		
Coconuts																		
Hazelnuts																		
Macadamia																		
Pecans																		
Pine nuts																		
Pistachios																		
Walnuts																		
Others																		
(iii) POME FRUIT	► M57 0,01 (*) ◄		► M57 0,3 ◄	2	► M38 2 (i) ◄	► M70 0,5 ◄		► M53 0,05 (*) ◄				► M53 0,5 ◄	► M58 0,2 ◄	0,05 (*)		0,1 (*)	0,05 (*)	
Apples		0,2								► M32 0,1 ◄								► M51 0,2 ◄
Pears		0,1								► M32 0,1 ◄								
Quinces																		
Others		0,05 (*)								► M32 0,02 (*) ◄								► M51 0,1 (*) ◄

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentazine	Cyromazine	Fenpropimorph	Flucythrinate	Hexaconazole	Methacryfos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)	
(iv) STONE FRUIT	► <u>M57</u> 0,01 (*) ◀		► <u>M57</u> 0,2 ◀		► <u>M38</u> 0,05 (*) ◀			► <u>M53</u> 0,05 (*) ◀		► <u>M32</u> 0,02 (*) ◀				0,05 (*)		0,1 (*)	0,05 (*)	► <u>M51</u> 0,1 (*) ◀	
Apricots				1								► <u>M53</u> 0,3 ◀	► <u>M58</u> 0,1 ◀						
Cherries				1								► <u>M53</u> 1 ◀							
Peaches (including nectarines and similar hybrids)				1								► <u>M53</u> 0,5 ◀	► <u>M58</u> 0,1 ◀						
Plums		0,3		2		► <u>M70</u> 0,2 ◀						► <u>M53</u> 0,5 ◀							
Others		0,05 (*)		0,05 (*)		► <u>M70</u> 0,02 (*) ◀						► <u>M53</u> 0,02 (*) ◀	► <u>M58</u> 0,05 (*) ◀						
(v) BERRIES AND SMALL FRUIT				0,05 (*)										0,05 (*)		0,1 (*)	0,05 (*)		
(a) Table and wine grapes	► <u>M57</u> 0,01 (*) ◀		► <u>M57</u> 0,2 ◀		► <u>M38</u> 2 (*) ◀			► <u>M53</u> 0,05 (*) ◀		► <u>M32</u> 0,1 ◀		► <u>M53</u> 1 ◀	► <u>M58</u> 0,2 ◀						► <u>M51</u> 2 ◀
Table grapes		0,05 (*)				► <u>M70</u> 0,02 (*) ◀													
Wine grapes		0,3				► <u>M70</u> 1 ◀													

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
(b) Strawberries (other than wild)	► <u>M57</u> 0,1 ◄	0,05 (*)	► <u>M57</u> 0,5 ◄		► <u>M38</u> 0,05 (*) ◄	► <u>M70</u> 2 ◄		► <u>M53</u> 1 ◄		► <u>M32</u> 0,2 ◄		► <u>M53</u> 1 ◄	► <u>M58</u> 0,5 ◄					► <u>M51</u> 0,5 ◄
(c) Cane fruit (other than wild)		0,05 (*)			► <u>M38</u> 0,05 (*) ◄			► <u>M53</u> 1 ◄		► <u>M32</u> 0,02 (*) ◄			► <u>M58</u> 0,05 (*) ◄					► <u>M51</u> 0,1 (*) ◄
Blackberries	► <u>M57</u> 0,1 ◄		► <u>M57</u> 0,3 ◄			► <u>M70</u> 3 ◄						► <u>M53</u> 1 ◄						
Dewberries																		
Loganberries																		
Raspberries	► <u>M57</u> 0,1 ◄		► <u>M57</u> 0,3 ◄			► <u>M70</u> 3 ◄						► <u>M53</u> 1 ◄						
Others	► <u>M57</u> 0,01 (*) ◄		► <u>M57</u> 0,05 (*) ◄			► <u>M70</u> 0,3 ◄						► <u>M53</u> 0,02 (*) ◄						
(d) Other small fruit and berries (other than wild)	► <u>M57</u> 0,01 (*) ◄	0,05 (*)			► <u>M38</u> 0,05 (*) ◄			► <u>M53</u> 1 ◄		► <u>M32</u> 0,02 (*) ◄								► <u>M51</u> 1 ◄
Bilberries (fruit of species <i>Vaccinium myrtillus</i>)																		
Cranberries																		

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacryfos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Currants (red, black and white)			► M57 0,5 ◄			► M70 0,5 ◄						► M53 1 ◄	► M58 0,5 ◄					
Gooseberries												► M53 1 ◄						
Others			► M57 0,05 (*) ◄			► M70 0,02 (*) ◄						► M53 0,02 (*) ◄	► M58 0,05 (*) ◄					
(e) Wild berries and wild fruit	► M57 0,01 (*) ◄	0,05 (*)	► M57 0,05 (*) ◄		► M38 0,05 (*) ◄	► M70 0,02 (*) ◄		► M53 0,05 (*) ◄		► M32 0,02 (*) ◄		► M53 0,02 (*) ◄	► M58 0,05 (*) ◄	0,05 (*)				► M51 0,1 (*) ◄
(vi) MISCELLANEOUS		0,05 (*)			► M38 0,05 (*) ◄								► M58 0,05 (*) ◄			0,1 (*)	0,05 (*)	
Avocados														5				
Bananas			► M57 0,1 ◄	3		► M70 2 ◄		► M53 2 ◄		► M32 0,1 ◄		► M53 2 ◄						► M51 0,2 ◄
Dates																		
Figs																		
Kiwi																		
Kumquat																		
Litchis																		
Mangoes			► M57 0,3 ◄											5				
Olives																		

▼ **M28**

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Passion fruit																		
Pineapples														5				► M51 3 ◀
Papaya	► M57 0,05 ◀		► M57 0,5 ◀											5				
Others	► M57 0,01 (*) ◀		► M57 0,05 (*) ◀	0,05 (*)		► M70 0,02 (*) ◀		► M53 0,05 (*) ◀		► M32 0,02 (*) ◀		► M53 0,02 (*) ◀		0,05 (*)				► M51 0,1 (*) ◀
2. Vegetables, fresh or uncooked, frozen or dry					► M38 — ◀				0,05 (*)	► M32 — ◀	0,05 (*)					0,1 (*)		
(i) ROOT AND TUBER VEGETABLES	► M57 0,01 (*) ◀	0,05 (*)	► M57 0,05 (*) ◀	0,05 (*)	► M38 0,05 (*) ◀	► M70 0,02 (*) ◀		► M53 0,05 (*) ◀		► M32 0,02 (*) ◀			► M58 0,05 (*) ◀	0,05 (*)	0,05 (*)		0,05 (*)	► M51 0,1 (*) ◀
Beetroot																		
Carrots							► M70 1 ◀											► M53 0,2 ◀
Celeriac																		
Horseradish																		► M53 0,2 ◀
Jerusalem artichokes																		
Parsnip																		► M53 0,2 ◀

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Parsley root												► M53 0,2 ◄						
Radishes																		
Salsify																		
Sweet potatoes																		
Swedes																		
Turnips																		
Yam																		
Others							► M70 0,05 (*)					► M53 0,02 (*)						
(ii) BULB VEGETABLES	► M57 0,01 (*)	0,05 (*)	► M57 0,05 (*)	0,05 (*)	► M38 0,05 (*)	► M70 0,02 (*)	► M70 0,05 (*)	► M53 0,05 (*)		► M32 0,02 (*)		► M53 0,02 (*)	► M58 0,05 (*)		0,05 (*)		0,05 (*)	
Garlic														0,5				
Onions																		► M51 0,5 ◄
Shallots														5				► M51 1 ◄
Spring onions																		► M51 1 ◄
Others														0,05 (*)				► M51 0,1 (*)

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentazine	Cyromazine	Fenpropimorph	Flucythrinate	Hexaconazole	Methacryfos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
(iii) FRUITING VEGETABLES								► M53 0,05 (*) ◄						0,05 (*)			0,05 (*)	
(a) Solanacea		0,05 (*)	► M57 0,2 ◄					► M70 1 ◄										
Tomatoes	► M57 0,02 ◄			3	► M38 1 (†) ◄	► M70 0,3 ◄				► M32 0,1 ◄		► M53 0,3 ◄	► M58 0,1 ◄					► M51 0,3 ◄
Peppers	► M57 0,05 ◄											► M53 0,5 ◄	► M58 0,2 ◄					► M51 0,5 ◄
Chilli Peppers														5				
Aubergines	► M57 0,02 ◄											► M53 0,3 ◄	► M58 0,1 ◄					
Others	► M57 0,01 (*) ◄			0,05 (*)	► M38 0,05 (*) ◄	► M70 0,02 (*) ◄				► M32 0,02 (*) ◄		► M53 0,02 (*) ◄	► M58 0,05 (*) ◄		0,05 (*)			► M51 0,1 (*) ◄
(b) Cucurbits - edible peel	► M57 0,02 ◄	0,05 (*)	► M57 0,1 ◄	0,5	► M38 0,05 (*) ◄	► M70 0,02 (*) ◄	► M70 1 ◄			► M32 0,02 (*) ◄		► M53 0,1 ◄	► M58 0,1 ◄		0,05 (*)			► M51 0,1 (*) ◄
Cucumbers																		
Gherkins																		

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentazine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Courgettes																		
Others																		
(c) Cucurbits - inedible peel	► <u>M57</u> 0,01 (*) ◄	0,05 (*)	► <u>M57</u> 0,05 (*) ◄	0,05 (*)	► <u>M38</u> 0,05 (*) ◄					► <u>M32</u> 0,02 (*) ◄		► <u>M53</u> 0,2 ◄	► <u>M58</u> 0,1 ◄		0,05 (*)			► <u>M51</u> 0,1 (*) ◄
Melons						► <u>M70</u> 0,1 ◄	► <u>M70</u> 0,3 ◄											
Squashes																		
Watermelons							► <u>M70</u> 0,3 ◄											
Others						► <u>M70</u> 0,02 (*) ◄	► <u>M70</u> 0,05 (*) ◄											
(d) Sweetcorn	► <u>M57</u> 0,01 (*) ◄	0,05 (*)	► <u>M57</u> 0,05 (*) ◄	0,05 (*)	► <u>M38</u> 0,05 (*) ◄	► <u>M70</u> 0,02 (*) ◄	► <u>M70</u> 0,05 (*) ◄			► <u>M32</u> 0,02 (*) ◄		► <u>M53</u> 0,02 (*) ◄	► <u>M58</u> 0,05 (*) ◄		0,05 (*)			► <u>M51</u> 0,1 (*) ◄
(iv) BRASSICA VEGETABLES	► <u>M57</u> 0,01 (*) ◄	0,05 (*)		0,05 (*)	► <u>M38</u> 0,05 (*) ◄	► <u>M70</u> 0,02 (*) ◄	► <u>M70</u> 0,05 (*) ◄			► <u>M32</u> 0,02 (*) ◄		► <u>M53</u> 0,02 (*) ◄	► <u>M58</u> 0,05 (*) ◄	0,05 (*)	0,05 (*)		0,05 (*)	► <u>M51</u> 0,1 (*) ◄
(a) Flowering brassica			► <u>M57</u> 0,2 ◄							► <u>M53</u> 0,05 (*) ◄								
Broccoli (including calabrese)																		

▼ **M28**

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacryfos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Cauliflower																		
Others																		
(b) Head brassica			► <u>M57</u> 1 ◄															
Brussels sprouts								► <u>M53</u> 0,5 ◄										
Head cabbage								► <u>M53</u> 0,05 (*) ◄										
Others								► <u>M53</u> 0,05 (*) ◄										
(c) Leafy brassica			► <u>M57</u> 0,05 (*) ◄					► <u>M53</u> 0,05 (*) ◄										
Chinese cabbage																		
Kale																		
Others																		
(d) Kohlrabi			► <u>M57</u> 0,05 (*) ◄					► <u>M53</u> 0,05 (*) ◄										
(v) LEAF VEGETABLES AND FRESH		0,05 (*)		0,05 (*)	► <u>M38</u> 0,05 (*) ◄	► <u>M70</u> 0,02 (*) ◄		► <u>M53</u> 0,05 (*) ◄		► <u>M32</u> 0,02 (*) ◄			► <u>M58</u> 0,05 (*) ◄		0,05 (*)		0,05 (*)	► <u>M51</u> 0,1 (*) ◄
(a) Lettuce and similar	► <u>M57</u> 0,1 ◄		► <u>M57</u> 2 ◄				► <u>M70</u> 15 ◄							5				
Cress																		

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Lamb's lettuce												► <u>M53</u> 5 ◄						
Lettuce																		
Scarole (broad-leaf endive)																		
Others												► <u>M53</u> 0,02 (*) ◄						
(b) Spinach and similar	► <u>M57</u> 0,01 (*) ◄		► <u>M57</u> 0,05 (*) ◄									► <u>M53</u> 0,02 (*) ◄		0,05 (*)				
Spinach																		
Beet leaves (chard)							► <u>M70</u> 20 ◄											
Others							► <u>M70</u> 0,05 (*) ◄											
(c) Water cress	► <u>M57</u> 0,01 (*) ◄		► <u>M57</u> 0,05 (*) ◄				► <u>M70</u> 0,05 (*) ◄					► <u>M53</u> 0,02 (*) ◄		0,05 (*)				
(d) Witloof	► <u>M57</u> 0,01 (*) ◄		► <u>M57</u> 0,05 (*) ◄				► <u>M70</u> 0,05 (*) ◄					► <u>M53</u> 0,02 (*) ◄		0,05 (*)				
(e) Herbs	► <u>M57</u> 1 ◄		► <u>M57</u> 0,05 (*) ◄				► <u>M70</u> 15 ◄					► <u>M53</u> 0,02 (*) ◄		5				
Chervil																		

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)	
Chives																			
Parsley																			
Celery leaves																			
Others																			
(vi) LEGUME VEGETABLES (fresh)	► <u>M57</u> 0,01 (*) ◄			0,05 (*)		► <u>M70</u> 0,02 (*) ◄		► <u>M53</u> 0,05 (*) ◄		► <u>M32</u> 0,02 (*) ◄			► <u>M58</u> 0,05 (*) ◄	0,05 (*)	0,05 (*)		0,05 (*)	► <u>M51</u> 0,1 (*) ◄	
Beans (with pods)		0,5	► <u>M57</u> 0,50 ◄		► <u>M38</u> 1 (1) ◄		► <u>M70</u> 5 ◄					► <u>M53</u> 0,3 (*) ◄							
Beans (without pods)																			
Peas (with pods)			► <u>M57</u> 0,1 ◄				► <u>M70</u> 5 ◄												
Peas (without pods)																			
Others		0,05 (*)	► <u>M57</u> 0,05 (*) ◄		► <u>M38</u> 0,05 (*) ◄		► <u>M70</u> 0,05 (*) ◄					► <u>M53</u> 0,02 (*) ◄							
(vii) STEM VEGETABLES (fresh)	► <u>M57</u> 0,01 (*) ◄	0,05 (*)	► <u>M57</u> 0,05 (*) ◄	0,05 (*)	► <u>M38</u> 0,05 (*) ◄	► <u>M70</u> 0,02 (*) ◄				► <u>M32</u> 0,02 (*) ◄				0,05 (*)	0,05 (*)		0,05 (*)		
Asparagus																			
Cardoons																			
Celery							► <u>M70</u> 2 ◄												

▼ M28

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentezine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
Fennel																		
Globe artichokes							► M70 2 ◀					► M53 0,5 ◀	► M58 0,2 ◀					► M51 1 ◀
Leek								► M53 1 ◀										
Rhubarb																		
Others							► M70 0,05 (*)	► M53 0,05 (*)				► M53 0,02 (*)	► M58 0,05 (*)					► M51 0,1 (*)
(viii) FUNGI	► M57 0,01 (*)	0,05 (*)	► M57 0,05 (*)	0,05 (*)	► M38 0,05 (*)	► M70 0,02 (*)		► M53 0,05 (*)		► M32 0,02 (*)		► M53 0,02 (*)	► M58 0,05 (*)		0,05 (*)	0,1 (*)	0,05 (*)	► M51 0,1 (*)
(a) Cultivated mushrooms							► M70 5 ◀							2				
(b) Wild mushrooms							► M70 0,05 (*)							0,05 (*)				
3. Pulses	► M57 0,01 (*)	0,05 (*)	► M57 0,05 (*)	0,05 (*)	► M38 0,05 (*)	► M70 0,02 (*)	► M70 0,05 (*)	► M53 0,05 (*)	0,05 (*)	► M32 0,02 (*)	0,05 (*)	► M53 0,02 (*)	► M58 0,05 (*)		0,05 (*)	0,1 (*)	0,05 (*)	► M51 0,1 (*)
Beans																		
Lentils																		
Peas														0,3				
Others														0,05 (*)				

▼ **M28**

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentazine	Cyromazine	Fenproimorph	Flucythrinate	Hexaconazole	Methacryfos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
4. Oil seed	► <u>M57</u> 0,02 (*) ◄	0,05 (*)	► <u>M57</u> 0,1 (*) ◄	0,1 (*)	► <u>M38</u> 0,1 (*) ◄	► <u>M70</u> 0,05 (*) ◄	► <u>M70</u> 0,05 (*) ◄	► <u>M53</u> 0,05 (*) ◄	0,05 (*)	► <u>M32</u> 0,05 (*) ◄	0,05 (*)	► <u>M53</u> 0,05 (*) ◄	► <u>M58</u> 0,05 (*) ◄			0,2 (*)	0,1 (*)	► <u>M51</u> 0,2 (*) ◄
Linseed														0,5				
Peanuts																		
Poppy seed																		
Sesame seed																		
Sunflower seed														0,5				
Rape seed														0,5				
Soya bean																		
Mustard seed																		
Cotton seed															2			
Others														0,1 (*)	0,05 (*)			
5. Potatoes	► <u>M57</u> 0,01 (*) ◄	0,05 (*)	► <u>M57</u> 0,05 (*) ◄	0,05 (*)	► <u>M38</u> 0,05 (*) ◄	► <u>M70</u> 0,02 (*) ◄	► <u>M70</u> 1 ◄	► <u>M53</u> 0,05 (*) ◄	0,05 (*)	► <u>M32</u> 0,02 (*) ◄	0,05 (*)	► <u>M53</u> 0,02 (*) ◄	► <u>M58</u> 0,05 (*) ◄	0,05 (*)	0,05 (*)	0,1 (*)	0,05 (*)	► <u>M51</u> 0,1 (*) ◄
Early and ware potatoes																		

▼ **M28**

Groups and examples of individual products to which the MRLs would apply	Abamectin (sum of avermectin B1a, avermectin B1b and delta-8,9 isomer of avermectin B1a)	Azocyclotin and Cyhexatin (sum of azocyclotin and cyhexatin expressed as cyhexatin)	Bifenthrin	Bitertanol	Bromopropylate	Clofentazine	Cyromazine	Fenpropimorph	Flucythrinate	Hexaconazole	Methacrifos	Myclobutanil	Penconazole	Prochloraz (sum of prochloraz and its metabolites containing the 2,4,6-Trichlorophenol moiety expressed as prochloraz)	Profenofos	Resmethrin, including other mixtures of constituent isomers (sum of isomers)	Tridemorph	Triadimefon and Triadimenol (sum of triadimefon and triadimenol)
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	► M57 0,02 (*) ◀	0,1 (*)	► M57 5 ◀	0,1 (*)	► M38 0,1 (*) ◀	► M70 0,05 (*) ◀	► M70 0,05 (*) ◀	► M53 0,1 (*) ◀	0,1 (*)	► M32 0,05 (*) ◀	0,1 (*)	► M53 0,05 (*) ◀	► M58 0,1 (*) ◀	0,1 (*)	0,1 (*)	0,2 (*)	20	► M51 0,2 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	► M57 0,05 ◀	0,1 (*)	► M57 10 ◀	0,1 (*)	► M38 0,1 (*) ◀	► M70 0,05 (*) ◀	► M70 0,05 (*) ◀	► M53 10 ◀	0,1 (*)	► M32 0,05 (*) ◀	0,1 (*)	► M53 2 ◀	► M58 0,5 ◀	0,1 (*)	0,1 (*)	0,2 (*)	0,1 (*)	► M51 10 ◀

(*) Indicates lower limit of analytical determination.

► **M38** (*) Indicates that the maximum level has been established temporarily until 31 December 2008 to accommodate an essential use in accordance with Regulation 2076/2002/EC. ◀

▼ **M29**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)		
	Triasulfuron	Thifensulfuron methyl	2,4-D (sum of 2,4-D and its esters) expressed as 2,4-D)
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)
(i) CITRUS FRUIT			► M36 1 (P) ◀
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)			
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			
(iii) POME FRUIT			
Apples			
Pears			
Quinces			
Others			
(iv) STONE FRUIT			
Apricots			
Cherries			
Peaches (including nectarines and similar hybrids)			
Plums			
Others			
(v) BERRIES AND SMALL FRUIT			
(a) Table and wine grapes			
Table grapes			
Wine grapes			
(b) Strawberries (other than wild)			

▼ **M29**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)		
	Triasulfuron	Thifensulfuron methyl	2,4-D (sum of 2,4-D and its esters) expressed as 2,4-D)
(c) Cane fruit (other than wild) Blackberries Dewberries Loganberries Raspberries Others			
(d) Other small fruit and berries (other than wild) Bilberries Cranberries Currants (red, black and white) Gooseberries Others			
(e) Wild berries and wild fruit			
(vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes Olives Passion fruit Pineapples Pomegranate Others			
2. Vegetables, fresh or uncooked, frozen or dry	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)
(i) ROOT AND TUBER VEGETABLES Beetroot Carrots Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips			

▼ **M29**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)		
	Triasulfuron	Thifensulfuron methyl	2,4-D (sum of 2,4-D and its esters) expressed as 2,4-D)
Yam			
Others			
(ii) BULB VEGETABLES			
Garlic			
Onions			
Shallots			
Spring onions			
Others			
(iii) FRUITING VEGETABLES			
(a) <i>Solanacea</i>			
Tomatoes			
Peppers			
Aubergines			
Others			
(b) <i>Cucurbits</i> — edible peel			
Cucumbers			
Gherkins			
Courgettes			
Others			
(c) <i>Cucurbits</i> — inedible peel			
Melons			
Squashes			
Watermelons			
Others			
(d) Sweet corn			
(iv) BRASSICA VEGETABLES			
(a) Flowering brassica			
Broccoli			
Cauliflower			
Others			
(b) Head brassica			
Brussels sprouts			
Head cabbage			
Others			
(c) Leafy brassica			
Chinese cabbage			
Kale			
Others			
(d) Kohlrabi			
(v) LEAF VEGETABLES AND FRESH HERBS			
(a) Lettuce & similar			
Cress			

▼ **M29**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)		
	Triasulfuron	Thifensulfuron methyl	2,4-D (sum of 2,4-D and its esters) expressed as 2,4-D)
Lamb's lettuce			
Lettuce			
Scarole			
Others			
(b) Spinach & similar			
Spinach			
Beet leaves (chard)			
Others			
(c) Water cress			
(d) Witloof			
(e) Herbs			
Chervil			
Chives			
Parsley			
Celery leaves			
Others			
(vi) LEGUME VEGETABLES (fresh)			
Beans (with pods)			
Beans (without pods)			
Peas (with pods)			
Peas (without pods)			
Others			
(vii) STEM VEGETABLES (fresh)			
Asparagus			
Cardoons			
Celery			
Fennel			
Globe artichokes			
Leek			
Rhubarb			
Others			
(viii) FUNGI			
(a) Cultivated mushrooms			
(b) Wild mushrooms			
3. Pulses	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)
Beans			
Lentils			
Peas			
Others			
4. Oil seeds	0,05 (*) (P)	0,05 (*) (P)	0,1 (*) (P)
Linseed			
Peanuts			
Poppy seeds			

▼ **M29**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)		
	Triasulfuron	Thifensulfuron methyl	2,4-D (sum of 2,4-D and its esters) expressed as 2,4-D)
Sesame seeds			
Sunflower seed			
Rape seed			
Soya bean			
Mustard seed			
Cotton seed			
Others			
5. Potatoes	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)
Early potatoes			
Ware potatoes			
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*) (P)	0,1 (*) (P)	0,1 (*) (P)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) (P)	0,1 (*) (P)	0,1 (*) (P)

(*) Indicates lower limit of analytical determination

(P) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 4 years from date of coming into force of this Directive.

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																				
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphonate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fen-tin hydr-oxide
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,05 (*) (p)	0,02 (*) (p)			0,05 (*) (p)		0,05 (*) (p)	0,02 (*) (p)	0,05 (*) (p)				0,05 (*) (p)			► M48 0,05 (*) (p) ◀	0,05 (*) (p)	► M46 0,05 (*) (p) ◀	► M63 0,05 (*) ◀	0,05 (*)	0,05 (*)
(i) CITRUS FRUIT			► M70 0,02 (*) ◀	0,01 (*) (p)		► M70 0,5 ◀				► M41 0,05 (*) (q) ◀			► M70 0,05 (*) (p) ◀	► M70 0,02 (*) (p) ◀		0,02 (*) (p)	0,01 (*) (p)				
Grapefruit																					
Lemons																					
Limes																					
Mandarins (including clementines and other hybrids)																					
Oranges																					
Pomelos																					
Others																					

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	ipropalcarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide	
(ii) TREE NUTS (shelled or unshelled)			► M70 0,02 (*) ◀	0,1 (*) (p)		► M70 0,05 (*) ◀		► M41 0,05 (*) (q) ◀			► M70 0,05 (*) (p) ◀			0,1 (*) (p)	0,01 (*) (p)							
Almonds																						
Brazil nuts																						
Cashew nuts																						
Chestnuts																						
Coconuts																						
Hazelnuts												► M70 0,1 (*) (p) ◀										
Macadamia																						
Pecans																						
Pine nuts																						
Pistachios																						

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
Walnuts																						
Others												► M70 0,02 (*) (p) ◀										
(iii) POME FRUIT			► M70 0,02 (*) ◀	0,01 (*) (p)		► M70 1 ◀		► M41 0,05 (*) (q) ◀			► M70 0,05 (*) (p) ◀	► M70 0,02 (*) (p) ◀		0,02 (*) (p)	0,01 (*) (p)							
Apples																						
Pears																						
Quinces																						
Others																						
(iv) STONE FRUIT			► M70 0,02 (*) ◀	0,01 (*) (p)		► M70 0,05 (*) ◀		► M41 0,05 (*) (q) ◀				► M70 0,02 (*) (p) ◀		0,02 (*) (p)	0,01 (*) (p)							
Apricots											► M70 5 (p) ◀											
Cherries											► M70 5 (p) ◀											

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphonate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
Peaches (including nectarines and similar hybrids)											► M70 5 (P) ◀											
Plums											► M70 1 (P) ◀											
Others											► M70 0,05 (*) (P) ◀											
(v) BERRIES AND SMALL FRUIT				0,01 (*) (P)								► M70 0,02 (*) (P) ◀		0,02 (*) (P)	0,01 (*) (P)							
(a) Table and wine grapes			► M70 2 ◀							► M41 2 (9) ◀	► M70 5 (P) ◀											
Table grapes						► M70 2 ◀																
Wine grapes						► M70 1 ◀																
(b) Strawberries (other than wild)			► M70 0,02 (*) ◀			► M70 0,5 ◀				► M41 0,05 (*) (9) ◀	► M70 5 (P) ◀											

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprotalcarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
(c) Cane fruit (other than wild)			► M70 0,02 (*) ◀			► M70 0,05 (*) ◀		► M41 0,05 (*) ◀ (9) ◀			► M70 10 (P) ◀											
Blackberries																						
Dewberries																						
Loganberries																						
Raspberries																						
Others																						
(d) Other small fruit and berries (other than wild)			► M70 0,02 (*) ◀			► M70 0,05 (*) ◀		► M41 0,05 (*) ◀ (9) ◀			► M70 5 (P) ◀											
Bilberries																						
Cranberries																						
Currants (red, black and white)																						
Gooseberries																						

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitrrole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide	
Others																						
(e) Wild berries and wild fruit											► M70 0,05 (*) (9) ◀											
(vi) MISCELLANEOUS			► M70 0,02 (*) ◀	0,01 (*) (p)		► M70 0,05 (*) ◀																
Avocados																						
Bananas																						
Dates																						
Figs																						
Kiwi																						
Kumquats																						
Litchis																						

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
Mangoes												► M70 0,5 (p) ◀										
Olives														0,05 (p)								
Passion fruit																						
Pineapples																						
Papaya																						
Others											► M70 0,05 (*) ◀ (p) ◀	► M70 0,02 (*) ◀ (p) ◀			0,01 (*) (p)							
2. Vegetables, fresh or uncooked, frozen or dry	0,05 (*) (p)	0,02 (*) (p)		0,01 (*) (p)	0,05 (*) (p)		0,05 (*) (p)		0,02 (*) (p)	0,05 (*) (p)				0,05 (*) (p)	0,02 (*) (p)	0,01 (*) (p)	► M48 0,05 (*) (p) ◀	0,05 (*) (p)	► M46 — ◀	► M63 0,05 (*) ◀	0,05 (*)	0,05 (*)
(i) ROOT AND TUBER VEGETABLES			► M70 0,02 (*) ◀						► M41 0,05 (*) (q) ◀		► M70 0,05 (*) ◀ (p) ◀	► M70 0,02 (*) ◀ (p) ◀				► M48 — ◀						
Beetroot																						► M46 0,1 (p) ◀
Carrots						► M70 0,1 ◀																

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 meta- laxyl- M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclo- nil- ide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide	
Celeriac																						
Horseradish						► M70 0,1 ◀																
Jerusalem artichokes																						
Parsnips						► M70 0,1 ◀																
Parsley root																						
Radishes						► M70 0,1 ◀																
Salsify																						
Sweet potatoes																						
Swedes																						
Turnips																						
Yam																						
Others						► M70 0,05 (*) ◀												► M46 0,05 (*) (P) ◀				

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																						
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprotalcarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide		
(ii) BULB VEGETABLES			► M70 0,02 (*) ◀					► M41 — ◀			► M70 0,05 (*) (P) ◀	► M70 0,02 (*) (P) ◀				► M48 — ◀		► M46 0,05 (*) (P) ◀					
Garlic						► M70 0,5 ◀																	
Onions						► M70 0,5 ◀		► M41 0,1 (9) ◀															
Shallots						► M70 0,5 ◀																	
Spring onions						► M70 0,2 ◀																	
Others						► M70 0,05 (*) ◀		► M41 0,05 (*) (9) ◀															
(iii) FRUITING VEGETABLES																► M48 — ◀		► M46 0,05 (*) (P) ◀					
(a) Solanacea																							

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitrole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide	
Tomatoes			► M70 1 ◀			► M70 0,2 ◀		► M41 1 ⁽⁹⁾ ◀			► M70 1 ^(P) ◀	► M70 1 ^(P) ◀										
Peppers						► M70 0,5 ◀					► M70 2 ^(P) ◀											
Aubergines			► M70 1 ◀								► M70 1 ^(P) ◀											
Others			► M70 0,02 (*) ◀			► M70 0,05 (*) ◀		► M41 0,05 (*) (9) ◀			► M70 0,05 (*) (P) ◀	► M70 0,02 (*) (P) ◀										
(b) Cucurbits — edible peel			► M70 0,2 ◀								► M70 1 ^(P) ◀	► M70 0,02 (*) (P) ◀										
Cucumbers						► M70 0,5 ◀		► M41 0,1 ⁽⁹⁾ ◀														
Gherkins								► M41 0,1 ⁽⁹⁾ ◀														

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► <u>C7</u> metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► <u>C7</u> cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphonate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
Courgettes							► <u>M41</u> 0,1 ⁽⁹⁾ ◀															
Others						► <u>M70</u> 0,05 (*) ◀		► <u>M41</u> 0,05 (*) (9) ◀														
(c) Cucurbits — inedible peel										► <u>M70</u> 0,05 (*) (P) ◀	► <u>M70</u> 0,02 (*) (P) ◀											
Melons			► <u>M70</u> 0,3 ◀			► <u>M70</u> 0,2 ◀		► <u>M41</u> 0,2 (9) ◀														
Squashes																						
Watermelons						► <u>M70</u> 0,2 ◀		► <u>M41</u> 0,2 (9) ◀														
Others			► <u>M70</u> 0,02 (*) ◀			► <u>M70</u> 0,05 (*) ◀		► <u>M41</u> 0,05 (*) (9) ◀														

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																						
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► <u>C7</u> metaxyl-M ◀	picolinafen	ipropalcarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► <u>C7</u> cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide		
(d) Sweet corn			► <u>M70</u> 0,02 (*) ◀			► <u>M70</u> 0,05 (*) ◀		► <u>M41</u> 0,05 (*) (9) ◀			► <u>M70</u> 0,05 (*) (p) ◀	► <u>M70</u> 0,02 (*) (p) ◀											
(iv) BRASSICA VEGETABLES			► <u>M70</u> 0,02 (*) ◀					► <u>M41</u> 0,05 (*) (9) ◀			► <u>M70</u> 0,05 (*) (p) ◀	► <u>M70</u> 0,02 (*) (p) ◀				► <u>M48</u> — ◀		► <u>M46</u> 0,05 (*) (p) ◀					
(a) Flowering brassica			► <u>M70</u> 0,1 ◀					► <u>M70</u> 0,2 ◀															
Broccoli																							
Cauliflower																							
Others																							
(b) Head brassica			► <u>M70</u> 0,02 (*) ◀																				
Brussels sprouts																							
Head cabbage						► <u>M70</u> 1 ◀																	

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																						
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► <u>M7</u> metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► <u>C7</u> cyclanilide ◀	pyraflufen-ethyl	amitrrole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide		
Others						► <u>M70</u> 0,05 (*) ◀																	
(c) Leafy brassica			► <u>M70</u> 0,02 (*) ◀																				
Chinese cabbage																							
Kale						► <u>M70</u> 0,2 ◀																	
Others						► <u>M70</u> 0,05 (*) ◀																	
(d) Kohlrabi			► <u>M70</u> 0,02 (*) ◀			► <u>M70</u> 0,05 (*) ◀																	
(v) LEAF VEGETABLES AND FRESH HERBS			► <u>M70</u> 0,02 (*) ◀													► <u>M48</u> ◀		► <u>M46</u> ◀					
(a) Lettuce and similar								► <u>M41</u> 1 ⁽⁹⁾ ◀			► <u>M70</u> 30 ^(P) ◀	► <u>M70</u> 0,02 ^(*) ^(P) ◀						► <u>M46</u> 0,05 ^(*) ^(P) ◀					

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► <u>M70</u> meta-laxyl-M ◀	picolinafen	ipropalcarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► <u>C7</u> cyclanilide ◀	pyraflufen-ethyl	amitrrole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide	
Cress						► <u>M70</u> 0,05 (*) ◀																
Lamb's lettuce						► <u>M70</u> 0,2 ◀																
Lettuce						► <u>M70</u> 2 ◀				► <u>M41</u> — ◀												
Scarole						► <u>M70</u> 1 ◀				► <u>M41</u> — ◀												
▼ M70																						
Ruccola						2																
Leaves and stems of brassica						2																
▼ M33																						
Others						► <u>M70</u> 0,05 (*) ◀				► <u>M41</u> — ◀												
(b) Spinach and similar						► <u>M70</u> 0,05 (*) ◀				► <u>M41</u> 0,05 (*) (9) ◀											► <u>M46</u> 0,05 (*) (P) ◀	

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
Spinach												► M70 0,3 (P) ◀										
Beet leaves (chard)																						
Others																						
(c) Water cress						► M70 0,05 (*) ◀		► M41 0,05 (*) (9) ◀			► M70 0,05 (*) (P) ◀	► M70 0,02 (*) (P) ◀										► M46 0,05 (*) (P) ◀
(d) Witloof						► M70 0,3 ◀		► M41 0,05 (*) (9) ◀			► M70 0,05 (*) (P) ◀	► M70 0,02 (*) (P) ◀										► M46 0,05 (*) (P) ◀
(e) Herbs						► M70 2 ◀		► M41 0,05 (*) (9) ◀			► M70 30 (P) ◀	► M70 0,3 (P) ◀										► M46 1 (P) ◀
Chervil																						

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► <u>C7</u> metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► <u>C7</u> cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
Chives																						
Parsley																						
Celery leaves																						
Others																						
(vi) LEGUME VEGETABLES (fresh)			► <u>M70</u> 0,02 (*) ◀			► <u>M70</u> 0,05 (*) ◀		► <u>M41</u> 0,05 (*) ◀ (9) ◀				► <u>M70</u> 0,02 (*) ◀ (P) ◀				► <u>M48</u> — ◀		► <u>M46</u> 0,05 (*) ◀ (P) ◀				
Beans (with pods)											► <u>M70</u> 2 (P) ◀											
Beans (without pods)																						
Peas (with pods)																						
Peas (without pods)																						
Others											► <u>M70</u> 0,05 (*) ◀ (P) ◀											

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► <u>C7</u> metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► <u>C7</u> cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide	
(vii) STEM VEGETABLES (fresh)							► <u>M41</u> 0,05 (*) (q) ◀				► <u>M70</u> 0,05 (*) (p) ◀	► <u>M70</u> 0,02 (*) (p) ◀				► <u>M48</u> ◀		► <u>M46</u> 0,05 (*) (p) ◀				
Asparagus																						
Cardoons																						
Celery																						
Fennel																						
Globe artichokes																						
Leek			► <u>M70</u> 2 ◀			► <u>M70</u> 0,2 ◀																
Rhubarb																						
Others			► <u>M70</u> 0,02 (*) ◀			► <u>M70</u> 0,05 (*) ◀																
(viii) FUNGI						► <u>M70</u> 0,05 (*) ◀	► <u>M41</u> 0,05 (*) (q) ◀				► <u>M70</u> 0,05 (*) (p) ◀	► <u>M70</u> 0,02 (*) (p) ◀				► <u>M48</u> ◀		► <u>M46</u> 0,05 (*) (p) ◀				
(a) Cultivated mushrooms																						

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► <u>C7</u> metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► <u>C7</u> cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
(b) Wild mushrooms																						
3. Pulses	0,05 (*)(p)	0,02 (*)(p)	► <u>M70</u> 0,02 (*)(p) ◀	0,01 (*)(p)	0,05 (*)(p)	► <u>M70</u> 0,05 (*)(p) ◀	0,05 (*)(p)	► <u>M41</u> 0,05 (*)(p) ◀	0,05 (*)(p)	0,05 (*)(p)	► <u>M70</u> 0,05 (*)(p) ◀	► <u>M70</u> 0,02 (*)(p) ◀	0,05 (*)(p)	0,02 (*)(p)	0,01 (*)(p)	► <u>M48</u> 0,2 (*)(p) ◀	0,05 (*)(p)	► <u>M46</u> 0,05 (*)(p) ◀	► <u>M63</u> 0,05 (*)(p) ◀	0,05 (*)(p)	0,05 (*)(p)	
Beans																► <u>M48</u> — ◀						
Lentils																						
Peas																► <u>M48</u> — ◀						
Others																► <u>M48</u> — ◀						
4. Oil seed	0,1 (*)(p)	0,05 (*)(p)	► <u>M70</u> 0,05 (*)(p) ◀	0,1 (*)(p)	0,1 (*)(p)		0,1 (*)(p)	► <u>M41</u> 0,1 (*)(p) ◀	0,1 (*)(p)	0,1 (*)(p)	► <u>M70</u> 0,1 (*)(p) ◀	► <u>M70</u> 0,05 (*)(p) ◀		0,05 (*)(p)	0,02 (*)(p)		0,1 (*)(p)	► <u>M46</u> 0,1 (*)(p) ◀	► <u>M63</u> 0,1 (*)(p) ◀	0,1 (*)(p)	0,1 (*)(p)	
Linseed						► <u>M70</u> 0,1 (*)(p) ◀										► <u>M48</u> 5 (*)(p) ◀						

▼ M33

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																				
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprovalicarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphamate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide
Peanuts																► M48 0,1 (*) (p) ◀					
Poppy seeds																► M48 0,1 (*) (p) ◀					
Sesame seeds																► M48 0,1 (*) (p) ◀					
Sunflower seed																► M48 1 (p) ◀					
Rape seed																► M48 2 (p) ◀					
Soya bean																► M48 0,2 (p) ◀					
Mustard seed																► M48 0,5 (p) ◀					
Cotton seed												0,2 (p)				► M48 0,1 (*) (p) ◀					

▼ **M33**

▼ **M48**

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																					
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprotalcarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitr-ole	diquat	isoproturon	etho-fumesate (sum of etho-fumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as etho-fumesate)	chlor-fenapyr	fentin acetate	fentin hydr-oxide	
Hemp seed																0,5 (p)						
Others													0,05 (*) (p)			► M48 0,1 (*) (p) ◀						
5. Potatoes	0,05 (*) (p)	0,02 (*) (p)	► M70 0,02 (*) ◀	0,01 (*) (p)	0,05 (*) (p)	► M70 0,05 (*) ◀	0,05 (*) (p)	► M41 0,05 (*) (q) ◀	0,02 (*) (p)	0,05 (*) (p)	► M70 0,05 (*) (p) ◀	► M70 0,02 (*) (p) ◀	0,05 (*) (p)	0,02 (*) (p)	0,01 (*) (p)	► M48 0,05 (*) (p) ◀	0,05 (*) (p)	► M46 0,05 (*) (p) ◀	► M63 0,05 (*) ◀	0,05 *	0,05 *	
Early potatoes																						
Ware potatoes																						
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of Camellia sinensis)	0,1 (*) (p)	0,05 (*) (p)	► M70 0,05 (*) ◀	0,1 (*) (p)	0,1 (*) (p)	► M70 0,1 (*) ◀	0,1 (*) (p)	► M41 0,1 (*) (q) ◀	0,1 (*) (p)	0,1 (*) (p)	► M70 0,1 (*) (p) ◀	► M70 0,05 (*) (p) ◀	0,1 (*) (p)	0,05 (*) (p)	0,02 (*) (p)	► M48 0,1 (*) (p) ◀	0,1 (*) (p)	► M46 0,1 (*) (p) ◀	► M63 50 ◀	0,1 (*)	0,1 (*)	
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) (p)	0,05 (*) (p)	► M70 0,05 (*) ◀	0,1 (*) (p)	0,1 (*) (p)	► M70 10 ◀	0,1 (*) (p)	► M41 0,1 (*) (q) ◀	0,1 (*) (p)	0,1 (*) (p)	► M70 0,1 (*) (p) ◀	► M70 0,05 (*) (p) ◀	0,1 (*) (p)	0,05 (*) (p)	0,02 (*) (p)	► M48 0,1 (*) (p) ◀	0,1 (*) (p)	► M46 0,1 (*) (p) ◀	► M63 0,1 (*) ◀	0,1 (*)	0,1 (*)	

▼ **M33**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)																				
	cinidon-ethyl (sum of cinidon-ethyl and its E-isomer)	cyhalofop butyl (sum of cyhalofop butyl and its free acids)	famoxadone	florasulam	flumioxazine	► C7 metaxyl-M ◀	picolinafen	iprotalcarb	pro-sulfuron	sulfosulfuron	fenhexamid	acibenzolar-S-methyl	► C7 cyclanilide ◀	pyraflufen-ethyl	amitrole	diquat	isoproturon	ethofumesate (sum of ethofumesate and the metabolite 2,3-dihydro-3,3-dimethyl-2-oxo-benzofuran-5-yl methane sulphinate expressed as ethofumesate)	chlorfenapyr	fentin acetate	fentin hydr-oxide
8. Spices Cumin seed Juniper seed Nutmeg Pepper, black and white Vanilla pods Others																		0,5 ^(p)			

▼ **M46**

▼ **M33**

(*) Indicates lower limit of analytical determination.

(p) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 14 July 2007.

► **M40** ^(p) Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC. ◀

► **M41** ^(*) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from [4 years from date of coming into force of the Directive introducing this amendment]. ◀

▼ **C10**

Groups and examples of individual products to which the MRLs apply	2,4-DB	Linuron	Pendimethalin	Imazamox	Oxasulfuron	Ethoxysulfuron	Foramsulfuron	Oxadiargyl	Cyazofamid
(iii) POME FRUIT									► M51 0,01 (*) (P) ◀
Apples									
Pears									
Quinces									
Others									
(iv) STONE FRUIT									► M51 0,01 (*) (P) ◀
Apricots									
Cherries									
Peaches (including nectarines and similar hybrids)									
Plums									
Others									
(v) BERRIES AND SMALL FRUIT									
(a) Table and wine grapes									► M51 0,5 (P) ◀
Table grapes									
Wine grapes									
(b) Strawberries (other than wild)									► M51 0,01 (*) (P) ◀
(c) Cane fruit (other than wild)									► M51 0,01 (*) (P) ◀
Blackberries									
Dewberries									
Loganberries									
Raspberries									
Others									

▼ **C10**

Groups and examples of individual products to which the MRLs apply	2,4-DB	Linuron	Pendimethalin	Imazamox	Oxasulfuron	Ethoxysulfuron	Foramsulfuron	Oxadiargyl	Cyazofamid
(d) Other small fruit and berries (other than wild)									► M51 0,01 (*) (P) ◀
Bilberries									
Cranberries									
Currants (red, black and white)									
Gooseberries									
Others									
(e) Wild berries and wild fruit									► M51 0,01 (*) (P) ◀
(vi) MISCELLANEOUS									► M51 0,01 (*) (P) ◀
Avocados									
Bananas									
Dates									
Figs									
Kiwi									
Kumquats									
Litchis									
Mangoes									
Olives									
Passion fruit									
Pineapples									
Papaya									
Others									
2. Vegetables, fresh or uncooked, frozen or dry	0,05 (*) (p)			0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)	0,01 (*) (P)	
(i) ROOT AND TUBER VEGETABLES									► M51 0,01 (*) (P) ◀
Beetroot									

▼ **C10**

Groups and examples of individual products to which the MRLs apply	2,4-DB	Linuron	Pendimethalin	Imazamox	Oxasulfuron	Ethoxysulfuron	Foramsulfuron	Oxadiargyl	Cyazofamid
Carrots		► <u>M57</u> 0,2 (P) ◄	► <u>M69</u> 0,2 ◄						
Celeriac		► <u>M57</u> 0,5 (P) ◄	► <u>M69</u> 0,1 ◄						
Horseradish			► <u>M69</u> 0,2 ◄						
Jerusalem artichokes									
Parsnips		► <u>M57</u> 0,2 (P) ◄	► <u>M69</u> 0,2 ◄						
Parsley root		► <u>M57</u> 0,2 (P) ◄	► <u>M69</u> 0,2 ◄						
Radishes									
Salsify									
Sweet potatoes									
Swedes									
Turnips									
Yam									
Others		► <u>M57</u> 0,05 (*) (P) ◄	► <u>M69</u> 0,05 (*) ◄						
(ii) BULB VEGETABLES		► <u>M57</u> 0,05 (*) (P) ◄	► <u>M69</u> 0,05 (*) ◄						► <u>M51</u> 0,01 (*) (P) ◄
Garlic									
Onions									
Shallots									
Spring onions									
Others									
(iii) FRUITING VEGETABLES		► <u>M57</u> 0,05 (*) (P) ◄	► <u>M69</u> 0,05 (*) ◄						
(a) Solanacea									
Tomatoes									► <u>M51</u> 0,2 (P) ◄

▼ **C10**

Groups and examples of individual products to which the MRLs apply	2,4-DB	Linuron	Pendimethalin	Imazamox	Oxasulfuron	Ethoxysulfuron	Foramsulfuron	Oxadiargyl	Cyazofamid
Peppers									
Aubergines									
Others									► M51 0,01 (*) (P) ◀
(b) Cucurbits — edible peel									► M51 0,1 (P) ◀
Cucumbers									
Gherkins									
Courgettes									
Others									
(c) Cucurbits — inedible peel									► M51 0,1 (P) ◀
Melons									
Squashes									
Watermelons									
Others									
(d) Sweet corn									► M51 0,01 (*) (P) ◀
(iv) BRASSICA VEGETABLES		► M57 0,05 (*) (P) ◀	► M69 0,05 (*) ◀						► M51 0,01 (*) (P) ◀
(a) Flowering brassica									
Broccoli									
Cauliflower									
Others									
(b) Head brassica									
Brussels sprouts									
Head cabbage									
Others									

▼ **C10**

Groups and examples of individual products to which the MRLs apply	2,4-DB	Linuron	Pendimethalin	Imazamox	Oxasulfuron	Ethoxysulfuron	Foramsulfuron	Oxadiargyl	Cyazofamid
(c) Leafy brassica									
Chinese cabbage									
Kale									
Others									
(d) Kohlrabi									
(v) LEAF VEGETABLES AND FRESH HERBS			► M69 0,05 (*) ◀						► M51 0,01 (*) (P) ◀
(a) Lettuce and similar		► M57 0,05 (*) (P) ◀							
Cress									
Lamb's lettuce									
Lettuce									
Scarole									
Others									
(b) Spinach and similar		► M57 0,05 (*) (P) ◀							
Spinach									
Beet leaves (chard)									
Others									
(c) Water cress		► M57 0,05 (*) (P) ◀							
(d) Witloof		► M57 0,05 (*) (P) ◀							
(e) Herbs		► M57 1 (P) ◀							
Chervil									
Chives									
Parsley									
Celery leaves									
Others									

▼ **C10**

Groups and examples of individual products to which the MRLs apply	2,4-DB	Linuron	Pendimethalin	Imazamox	Oxasulfuron	Ethoxysulfuron	Foramsulfuron	Oxadiargyl	Cyazofamid
(vi) LEGUME VEGETABLES (fresh)			► M69 0,2 ◀						► M51 0,01 (*) (P) ◀
Beans (with pods)									
Beans (without pods)		► M57 0,1 (P) ◀							
Peas (with pods)									
Peas (without pods)		► M57 0,1 (P) ◀							
Others		► M57 0,05 (*) (P) ◀							
(vii) STEM VEGETABLES (fresh)									► M51 0,01 (*) (P) ◀
Asparagus									
Cardoons									
Celery		► M57 0,1 (P) ◀	► M69 0,1 ◀						
Fennel		► M57 0,1 (P) ◀							
Globe artichokes									
Leek									
Rhubarb									
Others		► M57 0,05 (*) (P) ◀	► M69 0,05 (*) ◀						
(viii) FUNGI		► M57 0,05 (*) (P) ◀	► M69 0,05 (*) ◀						► M51 0,01 (*) (P) ◀
(a) Cultivated mushrooms									
(b) Wild mushrooms									

▼ C10

Groups and examples of individual products to which the MRLs apply	2,4-DB	Linuron	Pendimethalin	Imazamox	Oxasulfuron	Ethoxysulfuron	Foramsulfuron	Oxadiargyl	Cyazofamid
3. PULSES	0,05 (*) (P)	► M57 0,05 (*) (P) ◀	► M69 0,2 ◀	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)	0,01 (*) (P)	► M51 0,01 (*) (P) ◀
Beans									
Lentils									
Peas									
Others									
4. OIL SEED	0,05 (*) (P)	► M57 0,1 (*) (P) ◀	► M69 0,1 (*) ◀	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)	0,01 (*) (P)	► M51 0,02 (*) (P) ◀
Linseed									
Peanuts									
Poppy seeds									
Sesame seeds									
Sunflower seed									
Rape seed									
Soya bean									
Mustard seed									
Cotton seed									
Others									
5. POTATOES	0,05 (*) (P)	► M57 0,05 (*) (P) ◀	► M69 0,05 (*) ◀	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)	0,01 (*) (P)	► M51 0,01 (*) (P) ◀
Early potatoes									
Ware potatoes									
6. TEA (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (P) (*)	► M57 0,1 (*) (P) ◀	► M69 0,1 (*) ◀	0,1 (P) (*)	0,1 (P) (*)	0,1 (P) (*)	0,05 (*) (P)	0,05 (*) (P)	► M51 0,02 (*) (P) ◀
7. HOPS (dried), including hop pellets and unconcentrated powder	0,1 (P) (*)	► M57 0,1 (*) (P) ◀	► M69 0,1 (*) ◀	0,1 (P) (*)	0,1 (P) (*)	0,1 (P) (*)	0,05 (*) (P)	0,05 (*) (P)	► M51 0,02 (*) (P) ◀

(*) Indicates lower limit of analytical determination

► **M57** (P) Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 7 March 2011. ◀

▼ **M36**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)	
	Acephate	Parathion-methyl (sum of Parathion-methyl and para-oxon-methyl expressed as Parathion-methyl)
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,02 (*)	0,02 (*)
(i) CITRUS FRUIT		
Grapefruit		
Lemons		
Limes		
Mandarins (including clementines and other hybrids)		
Oranges		
Pomelos		
Others		
(ii) TREE NUTS (shelled or unshelled)		
Almonds		
Brazil nuts		
Cashew nuts		
Chestnuts		
Coconuts		
Hazelnuts		
Macadamia nuts		
Pecan nuts		
Pine nuts		
Pistachio nuts		
Walnuts		
Others		
(iii) POME FRUIT		
Apples		
Pears		
Quinces		
Others		
(iv) STONE FRUIT		
Apricots		
Cherries		
Peaches (including nectarines and similar hybrids)		
Plums		
Others		
(v) BERRIES AND SMALL FRUIT		
(a) Tables and wine grapes		
Table grapes		
Wine grapes		
(b) Strawberries (other than wild)		

▼ **M36**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)	
	Acephate	Parathion-methyl (sum of Parathion-methyl and para-oxon-methyl expressed as Parathion-methyl)
(c) Cane fruit (other than wild) Blackberries Dewberries Loganberries Raspberries Others		
(d) Other small fruit and berries (other than wild) Bilberries Cranberries Currants (red, black and white) Gooseberries Others		
(e) Wild berries and wild fruit		
(vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes Olives Passion fruit Pineapples Pomegranate Others		
2. Vegetables, fresh or uncooked, frozen or dry	0,02 (*)	0,02 (*)
(i) ROOT AND TUBER VEGETABLES Beetroot Carrots Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes		

▼ **M36**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)	
	Acephate	Parathion-methyl (sum of Parathion-methyl and para-oxon-methyl expressed as Parathion-methyl)
Swedes		
Turnips		
Yam		
Others		
(ii) BULB VEGETABLES		
Garlic		
Onions		
Shallots		
Spring onions		
Others		
(iii) FRUITING VEGETABLES		
(a) Solanacea		
Tomatoes		
Peppers		
Aubergines		
Others		
(b) Cucurbits — edible peel		
Cucumbers		
Gherkins		
Courgettes		
Others		
(c) Cucurbits — inedible peel		
Melons		
Squashes		
Watermelons		
Others		
(d) Sweetcorn		
(iv) BRASSICA VEGETABLES		
(a) Flowering brassica		
Broccoli		
Cauliflower		
Others		
(b) Head brassica		
Brussels sprouts		
Head cabbage		
Others		
(c) Leafy brassica		
Chinese cabbage		
Kale		
Others		
(d) Kohlrabi		

▼ **M36**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)	
	Acephate	Parathion-methyl (sum of Parathion-methyl and para-oxon-methyl expressed as Parathion-methyl)
(v) LEAF VEGETABLES AND FRESH HERBS		
(a) Lettuce and similar		
Cress		
Lamb's lettuce		
Lettuce		
Scarole		
Others		
(b) Spinach and similar		
Spinach		
Beet leaves (chard)		
Others		
(c) Water cress		
(d) Witloof		
(e) Herbs		
Chervil		
Chives		
Parsley		
Celery leaves		
Others		
(vi) LEGUME VEGETABLES (fresh)		
Beans (with pods)		
Beans (without pods)		
Peas (with pods)		
Peas (without pods)		
Others		
(vii) STEM VEGETABLES (fresh)		
Asparagus		
Cardoons		
Celery		
Fennel		
Globe artichokes		
Leek		
Rhubarb		
Others		
(viii) FUNGI		
(a) Cultivated mushrooms		
(b) Wild mushrooms		
3. PULSES	0,02 (*)	
Beans		
Lentils		
Peas		0,2
Others		0,02 (*)

▼ **M36**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue levels (mg/kg)	
	Acephate	Parathion-methyl (sum of Parathion-methyl and para-oxon-methyl expressed as Parathion-methyl)
4. OIL SEEDS	0,05 (*)	0,05 (*)
Linseed		
Peanuts		
Poppy seeds		
Sesame seeds		
Sunflower seed		
Rape seed		
Soya bean		
Mustard seed		
Cotton seed		
Others		
5. POTATOES	0,02 (*)	0,02 (*)
Early potatoes		
Ware potatoes		
6. TEA (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,05 (*)	0,05 (*)
7. HOPS (dried), including hop pellets and unconcentrated powder	0,05 (*)	0,05 (*)

(*) Indicates lower limit of analytical determination.

▼ **M37**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)
	Fenamiphos (sum of fenamiphos and its sulphoxide and sulphone expressed as fenamiphos)
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	
(i) CITRUS FRUIT	0,02 (*)
Grapefruit	
Lemons	
Limes	
Mandarins (including clementines and other hybrids)	
Oranges	
Pomelos	
Others	
(ii) TREE NUTS (shelled or unshelled)	0,02 (*)
Almonds	
Brazil nuts	
Cashew nuts	
Chestnuts	
Coconuts	
Hazelnuts	
Macadamia	
Pecans	
Pine nuts	
Pistachios	
Walnuts	
Others	
(iii) POME FRUIT	0,02 (*)
Apples	
Pears	
Quinces	
Others	
(iv) STONE FRUIT	0,02 (*)
Apricots	
Cherries	
Peaches (including nectarines and similar hybrids)	
Plums	
Others	
(v) BERRIES AND SMALL FRUIT	0,02 (*)
(a) Table and wine grapes	
Table grapes	
Wine grapes	
(b) Strawberries (other than wild)	

▼ **M37**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)
	Fenamiphos (sum of fenamiphos and its sulphoxide and sulphone expressed as fenamiphos)
(ii) BULB VEGETABLES	0,02 (*)
Garlic	
Onions	
Shallots	
Spring onions	
Others	
(iii) FRUITING VEGETABLES	
(a) Solanacea	
Tomatoes	0,05
Peppers	0,1
Aubergines	0,05
Others	0,02 (*)
(b) Cucurbits — edible peel	
Cucumbers	0,05
Gherkins	
Courgettes	0,05
Others	0,02 (*)
(c) Cucurbits — inedible peel	
Melons	0,05
Squashes	
Watermelons	0,05
Others	0,02 (*)
(d) Sweet corn	0,02 (*)
(iv) BRASSICA VEGETABLES	
(a) Flowering brassica	0,02 (*)
Broccoli (including Calabrese)	
Cauliflower	
Others	
(b) Head brassica	
Brussels sprouts	0,05
Head cabbage	0,05
Others	0,02 (*)
(c) Leafy brassica	0,02 (*)
Chinese cabbage	
Kale	
Others	
(d) Kohlrabi	0,02 (*)
(v) LEAF VEGETABLES AND FRESH HERBS	0,02 (*)
(a) Lettuce and similar	
Cress	
Lamb's lettuce	

▼ **M37**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)
	Fenamiphos (sum of fenamiphos and its sulphoxide and sulphone expressed as fenamiphos)
Lettuce Scarole (broad-leaf endive) Others (b) Spinach and similar Spinach Beet leaves (chard) Others (c) Water cress (d) Witloof (e) Herbs Chervil Chives Parsley Celery leaves Others	
(vi) LEGUME VEGETABLES (fresh)	0,02 (*)
Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others (vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek Rhubarb Others	0,02 (*)
(viii) FUNGI	0,02 (*)
(a) Cultivated mushrooms (b) Wild mushrooms 3. Pulses Beans Lentils Peas Others	0,02 (*)
4. Oilseeds Linseed Peanuts Poppy seed	0,05 (*)

▼ **M37**

Groups and examples of individual products to which the MRLs would apply	Pesticide residue and maximum residue level (mg/kg)
	Fenamiphos (sum of fenamiphos and its sulphoxide and sulphone expressed as fenamiphos)
Sesame seed Sunflower seed Rape seed Soya bean Mustard seed Cotton seed Others	
5. Potatoes Early potatoes Ware potatoes	0,02 (*)
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,05 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (*)

(*) Indicates lower limit of analytical determination.

▼ **M39**

Pesticide residue and maximum residue level (mg/kg)								
Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts i) CITRUS FRUIT Grapefruit Lemons Limes Mandarins (including clementines and other hybrids) Oranges Pomelos Others ii) TREE NUTS (shelled or unshelled) Almonds Brazil nuts Cashew nuts Chestnuts Coconuts Hazelnuts Macadamia Pecans Pine nuts Pistachios Walnuts Others	0,01 (*)	► M70 0,01 (*) ◀	0,01 (*)	0,01 (*)	► M66 0,01 (*) ◀	0,1 (*)	0,01 (*)	0,01 (*)

▼ M39

Pesticide residue and maximum residue level (mg/kg)								
Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
iii) POME FRUIT Apples Pears Quinces Others iv) STONE FRUIT Apricots Cherries Peaches (including nectarines and similar hybrids) Plums Others v) BERRIES AND SMALL FRUIT a) Table and wine grapes Table grapes Wine grapes b) Strawberries (other than wild) c) Cane fruit (other than wild) Blackberries Dewberries Loganberries Raspberries Others								

▼ M39

Pesticide residue and maximum residue level (mg/kg)								
Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
d) Other small fruit & berries (other than wild)								
Bilberries								
Cranberries								
Currants (red, black and white)								
Gooseberries								
Others								
e) Wild berries and wild fruit								
vi) MISCELLANEOUS								
Avocados								
Bananas								
Dates								
Figs								
Kiwi								
Kumquats								
Litchis								
Mangoes								
Olives								
Passion fruit								
Pineapples								
Pomegranate								
Others								

▼ M39

Pesticide residue and maximum residue level (mg/kg)

Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
2. Vegetables, fresh or uncooked, frozen or dry	0,01 (*)		0,01 (*)	0,01 (*)	► M66 0,01 (*) ◀	0,1 (*)	0,01 (*)	0,01 (*)
i) ROOT AND TUBER VEGETABLES								
Beetroot								
Carrots								
Celeriac								
Horseradish								
Jerusalem artichokes								
Parsnips		► M70 0,02 (*) ◀						
Parsley root								
Radishes								
Salsify								
Sweet potatoes								
Swedes								
Turnips								
Yam								
Others		► M70 0,01 (*) ◀						
ii) BULB VEGETABLES								
Garlic		► M70 0,01 (*) ◀						
Onions								
Shallots								

▼ M39

Pesticide residue and maximum residue level (mg/kg)								
Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
Spring onions								
Others								
iii) FRUITING VEGETABLES								
a) Solanacea		► <u>M70</u> 0,01 (*) ◀						
Tomatoes								
Peppers								
Aubergines								
Others								
b) Cucurbits — edible peel								
Cucumbers								
Gherkins								
Courgettes		► <u>M70</u> 0,05 ◀						
Others		► <u>M70</u> 0,02 (1) ◀						
c) Cucurbits-inedible peel								
Melons		► <u>M70</u> 0,03 (1) ◀						
Squashes								
Watermelons								
Others								
d) Sweet corn								

▼ M39

Pesticide residue and maximum residue level (mg/kg)								
Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
iv) BRASSICA VEGETABLES		► M70 0,01 (*) ◀						
a) Flowering brassica								
Broccoli (including Calabrese)								
Cauliflower								
Others								
b) Head brassica								
Brussels sprouts								
Head cabbage								
Others								
c) Leafy brassica								
Chinese cabbage								
Kale								
Others								
d) Kohlrabi								
v) LEAF VEGETABLES AND FRESH HERBS		► M70 0,01 (*) ◀						
a) Lettuce & similar								
Cress								
Lamb's lettuce								
Lettuce								
Scarole (broad-leaf endive)								
Others								

▼ M39

Pesticide residue and maximum residue level (mg/kg)								
Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
b) Spinach & similar Spinach Beet leaves (chard) Others								
c) Water cress								
d) Witloof								
e) Herbs Chervil Chives Parsley Celery leaves Others								
vi) LEGUME VEGETABLES (fresh)		► M70 0,01 (*) ◀						
Beans (with pods)								
Beans (without pods)								
Peas (with pods)								
Peas (without pods)								
Others								
vii) STEM VEGETABLES (fresh)		► M70 0,01 (*) ◀						
Asparagus								
Cardoons								
Celery								
Fennel								

▼ M39

Pesticide residue and maximum residue level (mg/kg)								
Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
Globe artichokes								
Leek								
Rhubarb								
Others								
viii) FUNGI		► M70 0,01 (*) ◀						
a) Cultivated mushrooms								
b) Wild mushrooms								
3. PULSES	0,01 (*)	► M70 0,01 (*) ◀	0,01 (*)	0,01 (*)	► M66 0,01 (*) ◀	0,1 (*)	0,01 (*)	0,01 (*)
Beans								
Lentils								
Peas								
Others								
4. OILSEEDS	0,02 (*)	► M70 0,02 (*) (2) ◀	0,02 (*)	0,02 (*)	0,02 (*)	0,2 (*)	0,02 (*)	0,02 (*)
Linseed								
Peanuts								
Poppy seed								
Sesame seed								
Sunflower seed								
Rape seed								
Soya bean								

▼ **M39**

Pesticide residue and maximum residue level (mg/kg)

Groups and examples of individual products to which the MRLs would apply	Sum of mercury compounds expressed as mercury	Aldrin and dieldrin combined expressed as dieldrin	Chlordane (sum of cis- and trans-chlordane)	HCH, Sum of isomers except the gamma isomer	Hexa-chloro-benzene	Ethylene oxyde (sum of ethylene oxyde and 2-chloro-ethanol expressed as ethylene oxyde)	Nitrofen	1,2-dichloro-ethane
Mustard seed								
Cotton seed								
▼ M66 Pumpkin seed		► M70 ⁽²⁾ ◄			0,05			
▼ M39 Others					► M66 0,02 (*) ◄			
5. POTATOES	0,01 (*)	► M70 0,01 (*) ◄	0,01 (*)	0,01 (*)	► M66 0,01 (*) ◄	0,1 (*)	0,01 (*)	0,01 (*)
Early potatoes								
Ware potatoes								
6. TEA (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,02 (*)	► M70 0,02 (*) ◄	0,02 (*)	0,02 (*)	► M66 0,02 (*) ◄	0,2 (*)	0,02 (*)	0,02 (*)
7. HOPS (dried), including hop pellets and unconcentrated powder	0,02 (*)	► M70 0,02 (*) ◄	0,02 (*)	0,02 (*)	► M66 0,02 (*) ◄	0,2 (*)	0,02 (*)	0,02 (*)

(*) Indicates lower limit of analytical determination.

(1) Based on background levels due to use of dieldrin and aldrin in the past.

(2) Monitoring data show that levels of up to 0,02 mg/kg of dieldrin can be found of pumpkin seeds used for oil extraction.

▼ **M1****Pesticide residues and maximum residue levels specifically in respect of tea (dried leaves and stalks, fermented or otherwise, *Camellia sinensis*)**

Pesticide residues	Maximum levels in mg/kg (ppm)	
1. Aldrin	► M8 0,02 ◀	
2. Dieldrin	} singly or combined expressed as dieldrin (HEOD)	
3. Endosulfan (sum of alpha- and beta-isomers and of endosulfan sulphate, expressed as endosulfan)		► M8 30 ◀
4. Hexachlorocyclohexane (HCH)		
4.1 alpha-isomer	} (sum)	
4.2 beta-isomer		► M8 0,2 ◀
4.3 gamma-isomer (lindane)		► M8 0,2 ◀
5. ► C1 Bifenthrin ◀	► M8 5 ◀	
6. Bromopropylate	► M8 0,1 (*) ◀	
7. Cartap	► M11 0,1 (*) ◀	
8. Chlordane (sum of cis- and trans-isomers)	► M8 0,02 (*) ◀	
9. Dichlorvos	► M8 0,1 (*) ◀	
10. Dicofol	► M8 20 ◀	
11. Dimethoate	► M8 0,2 ◀	
12. Omethoate	► M8 0,1 ◀	
13. Ethion	► M28 3 ◀	
14. Fenitrothion	► M8 0,5 ◀	
15. Flucythrinate (sum of isomers)	► M8 0,1 (*) ◀	
16. Hexachlorobenzene (HCB)	► M8 0,01 (*) ◀	
17. Malathion (sum of malathion and malaoxon expressed as malathion)	► M8 0,5 ◀	
18. Methidathion	► M8 0,1 (*) ◀	
19. Monocrotophos	► M8 0,1 (*) ◀	
20. Phoxim	► M8 0,1 (*) ◀	
21. Profenophos	► M8 0,1 (*) ◀	
22. Propargite	► M8 5 ◀	
23. Quinalphos	► M8 2 (*) ◀	
24. Phosmet (sum of phosmet and phosmet oxon expressed as phosmet)	► M8 0,1 (*) ◀	

(*) Indicates lower limit of analytical determination.

(a) (b) (c) (d) Should levels not be adopted by ► **M7** 31 October 1998 ◀, the following maximum levels shall apply as indicated thereafter:

- (a) 0,02 (*)
- (b) 0,01 (*)
- (c) 0,05 (*)
- (d) 0,1 (*).

► **M8** ► **C3** x Should this level not be confirmed or amended by a directive, with effect from 1 July 2000, the appropriate lower limit of analytical determination shall apply. ◀ ◀

▼ **M42**

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) (1)	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,05 (*) (a)		0,01 (*) (a)		0,05 (*) (a)
(i) CITRUS FRUIT		► M70 0,3 (p) ◀		0,02 (*) (a)	
Grapefruit					
Lemons					
Limes					
Mandarins (including clementines and other hybrids)					
Oranges					
Pomelos					
Others					
(ii) TREE NUTS (shelled or unshelled)		► M70 0,02 (*) (p) ◀		0,02 (*) (a)	
Almonds					
Brazil nuts					
Cashew nuts					
Chestnuts					
Coconuts					
Hazelnuts					
Macadamia					
Pecans					
Pine nuts					
Pistachios					
Walnuts					
Others					

▼ **M42**

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
(iii) POME FRUIT		► M70 0,5 ^(P) ◀		0,02 (*) ^(a)	
Apples					
Pears					
Quinces					
Others					
(iv) STONE FRUIT				0,02 (*) ^(a)	
Apricots		► M70 1 ^(P) ◀			
Cherries		► M70 1 ^(P) ◀			
Peaches (including nectarines and similar hybrids)		► M70 1 ^(P) ◀			
Plums		► M70 0,02 ^(P) ◀			
Others		► M70 0,02 (*) ^(P) ◀			
(v) BERRIES AND SMALL FRUIT					
(a) Table and wine grapes		► M70 5 ^(P) ◀		0,5 ^(a)	
Table grapes					
Wine grapes					
(b) Strawberries (other than wild)		► M70 0,5 ^(P) ◀		0,02 (*) ^(a)	
(c) Cane fruit (other than wild)		► M70 0,02 (*) ^(P) ◀		0,02 (*) ^(a)	
Blackberries					
Dewberries					
Loganberries					
Raspberries					
Others					

▼ M42

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
(d) Other small fruit and berries (other than wild)				0,02 (*) ^(a)	
Bilberries					
Cranberries					
Currants (red, black and white)		► <u>M70</u> 1 (P) ◀			
Gooseberries		► <u>M70</u> 1 (P) ◀			
Others		► <u>M70</u> 0,02 (*) ^(P) ◀			
(e) Wild berries and wild fruit		► <u>M70</u> 0,02 (*) ^(P) ◀		0,02 (*) ^(a)	
(vi) MISCELLANEOUS				0,02 (*) ^(a)	
Avocados					
Bananas		► <u>M70</u> 0,05 ^(P) ◀			
Dates					
Figs					
Kiwi					
Kumquats					
Litchis					
Mangoes		► <u>M70</u> 0,5 ^(P) ◀			
Olives (table consumption)		0,3 (P)			
Olives (oil extraction)		0,3 (P)			
Passion fruit					

▼ M70▼ M42

▼ M42

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
Pineapples					
Papaya		► <u>M70</u> 1 (P) ◀			
Others		► <u>M70</u> 0,02 (*) (P) ◀			
2. Vegetables, fresh or uncooked, frozen or dry	0,05 (*) (a)		0,01 (*) (a)		0,05 (*) (a)
(i) ROOT AND TUBER VEGETABLES				0,02 (*) (a)	
Beetroot					
Carrots		► <u>M70</u> 0,05 (P) ◀			
Celeriac					
Horseradish					
Jerusalem artichokes					
Parsnips					
Parsley root					
Radishes					
Salsify					
Sweet potatoes					
Swedes					
Turnips					
Yam					
Others		► <u>M70</u> 0,02 (*) (P) ◀			
(ii) BULB VEGETABLES		► <u>M70</u> 0,02 (*) (P) ◀		0,02 (*) (a)	
Garlic					
Onions					

▼ M42

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
Shallots					
Spring onions					
Others					
(iii) FRUITING VEGETABLES					
(a) Solanacea					
Tomatoes		▶ <u>M70</u> 0,5 ^(P) ◀		0,5 ^(a)	
Peppers		▶ <u>M70</u> 0,3 ^(P) ◀			
Aubergines					
Others		▶ <u>M70</u> 0,02 ^(*) ^(P) ◀		0,02 ^(*) ^(a)	
(b) Cucurbits — edible peel		▶ <u>M70</u> 0,2 ^(P) ◀		0,02 ^(*) ^(a)	
Cucumbers					
Gherkins					
Courgettes					
Others					
(c) Cucurbits — inedible peel					
Melons		▶ <u>M70</u> 0,3 ^(P) ◀		0,1 ^(a)	
Squashes		▶ <u>M70</u> 0,2 ^(P) ◀			
Watermelons		▶ <u>M70</u> 0,2 ◀			
Others		▶ <u>M70</u> 0,02 ^(*) ^(P) ◀		0,02 ^(*) ^(a)	
(d) Sweet corn		▶ <u>M70</u> 0,02 ^(*) ^(P) ◀		0,02 ^(*) ^(a)	

▼ **M42**

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
(iv) BRASSICA VEGETABLES				0,02 (*) ^(a)	
(a) Flowering brassica					
Broccoli		▶ M70 0,05 ^(p) ◀			
Cauliflower		▶ M70 0,05 ^(p) ◀			
Others		▶ M70 0,02 (*) ^(p) ◀			
(b) Head brassica					
Brussels sprouts					
Head cabbage					
Others					
(c) Leafy brassica					
Chinese cabbage		▶ M70 0,02 (*) ^(p) ◀			
Kale					
Others					
(d) Kohlrabi					
		▶ M70 0,02 (*) ^(p) ◀			
(v) LEAF VEGETABLES AND FRESH HERBS					
(a) Lettuce and similar					
Cress					
Lamb's lettuce					
Lettuce					
Scarole					
Others					
				2 ^(a)	

▼ M42

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
(b) Spinach and similar Spinach Beet leaves (chard) Others				0,02 (*) ^(a)	
(c) Watercress				0,02 (*) ^(a)	
(d) Witloof				0,02 (*) ^(a)	
(e) Herbs Chervil Chives Parsley Celery leaves Others				0,02 (*) ^(a)	
(vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others		► M70 0,5 ^(P) ◀		0,02 (*) ^(a)	
(vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel		► M70 0,02 (*) ^(P) ◀		0,02 (*) ^(a)	

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
Globe artichokes					
Leek		► M70 0,2 ^(P) ◀			
Rhubarb					
Others		► M70 0,02 ^(*) ^(P) ◀			
(viii) FUNGI				0,02 ^(*) ^(a)	
(a) Cultivated mushrooms		► M70 0,02 ^(*) ^(P) ◀			
(b) Wild mushrooms					
3. Pulses	0,05 ^(*) ^(a)	► M70 0,02 ^(*) ^(P) ◀	0,01 ^(*) ^(a)	0,02 ^(*) ^(a)	0,05 ^(*) ^(a)
Beans					
Lentils					
Peas					
Others					
4. Oil seed	0,1 ^(*) ^(a)	► M70 0,05 ^(*) ^(P) ◀	0,02 ^(*) ^(a)	0,05 ^(*) ^(a)	0,05 ^(*) ^(a)
Linseed					
Peanuts					
Poppy seeds					
Sesame seeds					
Sunflower seed					
Rape seed					
Soya bean					
Mustard seed					

▼ **M42**

Pesticide residues and maximum residue levels (mg/kg)					
Groups and examples of individual products to which the MRLs apply	Isoxaflutole (sum of isoxaflutole, RPA 202248 and RPA 203328, expressed as isoxaflutole) ⁽¹⁾	Trifloxystrobin	Carfentrazone-ethyl (determined as carfentrazone and expressed as carfentrazone-ethyl)	Fenamidone	Mecoprop (sum of mecoprop-p and mecoprop expressed as mecoprop)
Cotton seed					
Others		► M70 0,02 (*) ^(P) ◀			
5. Potatoes	0,05 (*) ^(a)	► M70 0,02 (*) ^(P) ◀	0,01 (*) ^(a)	0,02 (*) ^(a)	0,05 (*) ^(a)
Early potatoes					
Ware potatoes					
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*) ^(a)	► M70 0,05 (*) ^(P) ◀	0,02 (*) ^(a)	0,05 (*) ^(a)	0,1 (*) ^(a)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) ^(a)	► M70 30 ^(P) ◀	0,02 (*) ^(a)	0,05 (*) ^(a)	0,1 (*) ^(a)

⁽¹⁾ RPA 202248 is 2-cyano-3-cyclopropyl-1-(2-methylsulfonyl-4-trifluoromethylphenyl) propane-1,3-dione. RPA 203328 is 2-methanesulfonyl-4-trifluoromethylbenzoic acid.

(*) Indicates lower limit of analytical determination.

^(a) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 24 June 2009.

► **M70** ^(P) Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC. ◀

▼ **M44**

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,02 (*) (P)		0,05 (*) (P)
(i) CITRUS FRUIT						0,02 (*) (P)	
Grapefruit							
Lemons							
Limes							
Mandarins (including clementines and other hybrids)							
Oranges							
Pomelos							
Others							
(ii) TREE NUTS (shelled or unshelled)						0,02 (*) (P)	
Almonds							
Brazil nuts							
Cashew nuts							
Chestnuts							
Coconuts							
Hazelnuts							
Macadamia							
Pecans							
Pine nuts							
Pistachios							
Walnuts							
Others							

▼ M44

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
<p>(iii) POME FRUIT</p> <ul style="list-style-type: none"> Apples Pears Quinces Others <p>(iv) STONE FRUIT</p> <ul style="list-style-type: none"> Apricots Cherries Peaches (including nectarines and similar hybrids) Plums Others <p>(v) BERRIES AND SMALL FRUIT</p> <ul style="list-style-type: none"> (a) Table and wine grapes <ul style="list-style-type: none"> Table grapes Wine grapes (b) Strawberries (other than wild) (c) Cane fruit (other than wild) <ul style="list-style-type: none"> Blackberries Dewberries Loganberries Raspberries Others (d) Other small fruit and berries (other than wild) <ul style="list-style-type: none"> Bilberries 						<p>0,02 (*) ^(P)</p> <p>0,02 (*) ^(P)</p> <p>0,02 (*) ^(P)</p>	

▼ **M44**

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
Cranberries Currants (red, black and white) Gooseberries Others (e) Wild berries and wild fruit (vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes Olives Passion fruit Pineapples Papaya Others						0,05 (P)	
2. Vegetables, fresh or uncooked, frozen or dry (i) ROOT AND TUBER VEGETABLES Beetroot Carrots Celeriac Horseradish	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,02 (*) (P)	0,02 (*) (P)	0,05 (*) (P)

▼ **M44**

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips Yam Others (ii) BULB VEGETABLES Garlic Onions Shallots Spring onions Others (iii) FRUITING VEGETABLES (a) Solanacea Tomatoes Peppers Aubergines Others (b) Cucurbits — edible peel Cucumbers Gherkins Courgettes Others							

▼ **M44**

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
<p>(c) Cucurbits — inedible peel</p> <ul style="list-style-type: none"> Melons Squashes Watermelons Others <p>(d) Sweet corn</p> <p>(iv) BRASSICA VEGETABLES</p> <ul style="list-style-type: none"> (a) Flowering brassica <ul style="list-style-type: none"> Broccoli Cauliflower Others (b) Head brassica <ul style="list-style-type: none"> Brussels sprouts Head cabbage Others (c) Leafy brassica <ul style="list-style-type: none"> Chinese cabbage Kale Others (d) Kohlrabi <p>(v) LEAF VEGETABLES AND FRESH HERBS</p> <ul style="list-style-type: none"> (a) Lettuce & similar <ul style="list-style-type: none"> Cress Lamb's lettuce Lettuce Scarole Others 							

▼ **M44**

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
(b) Spinach & similar Spinach Beet leaves (chard) Others (c) Water cress (d) Witloof (e) Herbs Chervil Chives Parsley Celery leaves Others (vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others (vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek							

▼ **M44**

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
Rhubarb Others (viii) FUNGI (a) Cultivated mushrooms (b) Wild mushrooms							
3. Pulses Beans Lentils Peas Others	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,02 (*) (P)	0,02 (*) (P)	0,05 (*) (P)
4. Oil seed Linseed Peanuts Poppy seeds Sesame seeds Sunflower seed Rape seed Soya bean Mustard seed Cotton seed Others	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,02 (*) (P)	0,05 (*) (P)	0,05 (*) (P)
5. Potatoes Early potatoes Ware potatoes	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,1 (P)	0,02 (*) (P)	0,02 (*) (P)	0,05 (*) (P)

▼ **M44**

Groups and examples of individual products to which the MRLs apply	Mesotrione (Sum of mesotrione and MNBA (4-methylsulfonyl-2-nitro benzoic acid), expressed as mesotrione)	Silthiofam	Picoxystrobin	Flufenacet (Sum of all compounds containing the N fluorophenyl-N-isopropyl moiety expressed as flufenacet)	Iodosulfuron-methyl sodium (iodosulfuron-methyl including salts, expressed as iodosulfuron-methyl)	Fosthiazate	Molinate
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*) (P)	0,1 (*) (P)	0,1 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,1 (*) (P)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) (P)	0,1 (*) (P)	0,1 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,05 (*) (P)	0,1 (*) (P)

(*) Indicates lower limit of analytical determination.

(P) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 13 September 2009.

▼ **M45**

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)		0,02 (*) (P)	► M66 0,05 (*) (P) ◀		0,02 (*) (P)				
(i) CITRUS FRUIT				0,02 (P)			► M70 0,01 (*) (P) ◀		► M70 1 (P) ◀	► M66 0,02 (*) (P) ◀		0,02 (*) (P)
Grapefruit												
Lemons												
Limes												
Mandarins (including clementines and other hybrids)											► M53 0,5 (P) ◀	
Oranges											► M53 0,5 (P) ◀	
Pomelos											► M53 0,05 (*) (P) ◀	
Others											► M53 0,05 (*) (P) ◀	
(ii) TREE NUTS (shelled or unshelled)				0,01 (*) (P)			► M70 0,01 (*) (P) ◀			► M66 0,02 (*) (P) ◀	► M53 0,05 (*) (P) ◀	0,02 (*) (P)
Almonds												
Brazil nuts												
Cashew nuts												
Chestnuts												
Coconuts												

▼ **M45**

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Hazelnuts												
Macadamia												
Pecans												
Pine nuts												
Pistachios									► M70 1 (P) ◄			
Walnuts												
Others									► M70 0,02 (*) (P) ◄			
(iii) POME FRUIT				0,01 (*) (P)			► M70 0,01 (*) (P) ◄		► M70 0,3 (P) ◄		► M53 0,05 (*) (P) ◄	0,02 (*) (P)
Apples										► M66 0,05 (P) ◄		
Pears												
Quinces												
Others										► M66 0,02 (*) (P) ◄		
(iv) STONE FRUIT				0,01 (*) (P)			► M70 0,01 (*) (P) ◄				► M53 0,05 (*) (P) ◄	0,02 (*) (P)
Apricots									► M70 0,2 (P) ◄	► M66 0,05 (P) ◄		

▼ M45

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Cherries									► M70 0,3 (P) ◄	► M66 0,3 (P) ◄		
Peaches (including nectarines and similar hybrids)									► M70 0,2 (P) ◄	► M66 0,05 (P) ◄		
Plums									► M70 0,2 (P) ◄			
Others									► M70 0,02 (*) (P) ◄	► M66 0,02 (*) (P) ◄		
(v) BERRIES AND SMALL FRUIT											► M53 0,05 (*) (P) ◄	
(a) Table and wine grapes				0,02 (P)			► M70 3 (P) ◄			► M66 1 (P) ◄		5 (P)
Table grapes									► M70 1 (P) ◄			
Wine grapes									► M70 2 (P) ◄			
(b) Strawberries (other than wild)				0,01 (*) (P)			► M70 2 (P) ◄		► M70 0,5 (P) ◄	► M66 0,3 (P) ◄		0,02 (*) (P)
(c) Cane fruit (other than wild)				0,01 (*) (P)			► M70 0,01 (*) (P) ◄			► M66 0,02 (*) (P) ◄		0,02 (*) (P)
Blackberries									► M70 1 (P) ◄			
Dewberries												
Loganberries												

▼ M45

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxifen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Raspberries									► M70 1 (P) ◄			
Others									► M70 0,02 (*) (P) ◄			
(d) Other small fruit and berries (other than wild)				0,01 (*) (P)			► M70 0,01 (*) (P) ◄			► M66 2 (P) ◄		0,02 (*) (P)
Bilberries												
Cranberries												
Currants (red, black and white)									► M70 2 (P) ◄			
Gooseberries												
Others									► M70 0,5 (P) ◄			
(e) Wild berries and wild fruit				0,01 (*) (P)			► M70 0,01 (*) (P) ◄		► M70 0,02 (*) (P) ◄	► M66 0,02 (*) (P) ◄		0,02 (*) (P)
(vi) MISCELLANEOUS							► M70 0,01 (*) (P) ◄			► M66 0,02 (*) (P) ◄		0,02 (*) (P)
Avocados												
Bananas												
Dates												
Figs												
Kiwi												

▼ M45

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Kumquats												
Litchis												
Mangoes									► M70 0,05 (P) ◀			
Olives				0,02 (P)								
Olives (table consumption)	—	—	—	—	—			—			—	—
Olives (oil extraction)	—	—	—	—	—			—			► M53 1 (P) ◀	—
Passion fruit												
Pineapples												
Papaya									► M70 0,05 (P) ◀			
Others				0,01 (*) (P)					► M70 0,02 (*) (P) ◀		► M53 0,05 (*) (P) ◀	
2. Vegetables, fresh or uncooked, frozen or dry	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)	0,01 (*) (P)	0,02 (*) (P)			0,02 (*) (P)				
(i) ROOT AND TUBER VEGETABLES							► M70 0,01 (*) (P) ◀			► M66 0,02 (*) (P) ◀	► M53 0,05 (*) (P) ◀	0,02 (*) (P)
Beetroot												
Carrots						► M66 0,2 (P) ◀			► M70 0,1 (P) ◀			
Celeriac												
Horseradish									► M70 0,3 (P) ◀			
Jerusalem artichokes												

▼ **M45**

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Parsnips						► M66 0,2 (P) ◄			► M70 0,3 (P) ◄			
Parsley root									► M70 0,1 (P) ◄			
Radishes									► M70 0,2 (P) ◄			
Salsify									► M70 0,1 (P) ◄			
Sweet potatoes												
Swedes												
Turnips												
Yam												
Others						► M66 0,05 (*) (P) ◄			► M70 0,02 (*) (P) ◄			
(ii) BULB VEGETABLES							► M70 0,01 (*) (P) ◄			► M66 0,02 (*) (P) ◄	► M53 0,05 (*) (P) ◄	0,02 (*) (P)
Garlic						► M66 0,2 (P) ◄			► M70 0,2 (P) ◄			
Onions						► M66 0,2 (P) ◄			► M70 0,2 ◄			
Shallots						► M66 0,2 (P) ◄			► M70 0,2 (P) ◄			
Spring onions						► M66 3 (P) ◄						
Others						► M66 0,05 (*) (P) ◄			► M70 0,02 (*) (P) ◄			

▼ M45

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
(iii) FRUITING VEGETABLES						► M66 0,05 (*) (P) ◀					► M53 0,05 (*) (P) ◀	
(a) Solanacea										► M66 0,02 (*) (P) ◀		
Tomatoes							► M70 1 (P) ◀		► M70 0,2 (P) ◀			0,5 (P)
Peppers									► M70 0,5 (P) ◀			
Aubergines							► M70 1 (P) ◀		► M70 0,2 (P) ◀			
Others							► M70 0,01 (*) (P) ◀		► M70 0,02 (*) (P) ◀			0,02 (*) (P)
(b) Cucurbits — edible peel							► M70 0,01 (*) (P) ◀		► M70 0,02 (*) (P) ◀	► M66 0,02 (*) (P) ◀		0,02 (*) (P)
Cucumbers												
Gherkins												
Courgettes												
Others												
(c) Cucurbits — inedible peel							► M70 0,01 (*) (P) ◀		► M70 0,02 (*) (P) ◀	► M66 0,05 (P) ◀		0,02 (*) (P)
Melons												
Squashes												

▼ M45

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Watermelons												
Others												
(d) Sweet corn							► M70 0,01 (*) (P) ◀		► M70 0,02 (*) (P) ◀			0,02 (*) (P)
(iv) BRASSICA VEGETABLES						► M66 0,05 (*) (P) ◀	► M70 0,01 (*) (P) ◀			► M66 0,02 (*) (P) ◀	► M53 0,05 (*) (P) ◀	0,02 (*) (P)
(a) Flowering brassica									► M70 0,1 (P) ◀			
Broccoli												
Cauliflower												
Others												
(b) Head brassica												
Brussels sprouts									► M70 0,2 (P) ◀			
Head cabbage									► M70 0,2 (P) ◀			
Others									► M70 0,02 (*) (P) ◀			
(c) Leafy brassica									► M70 0,02 (*) (P) ◀			
Chinese cabbage												
Kale												

▼ **M45**

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxifen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Others												
(d) Kohlrabi									► M70 0,02 (*) (P) ◀			
(v) LEAF VEGETABLES AND FRESH HERBS						► M66 0,05 (*) (P) ◀	► M70 0,01 (*) (P) ◀			► M66 0,02 (*) (P) ◀	► M53 0,05 (*) (P) ◀	0,02 (*) (P)
(a) Lettuce & similar												
Cress												
Lamb's lettuce									► M70 10 (P) ◀			
Lettuce												
Scarole									► M70 2 (P) ◀			
Others									► M70 0,05 (P) ◀			
(b) Spinach & similar												
Spinach												
Beet leaves (chard)												
Others												
(c) Water cress									► M70 0,02 (*) (P) ◀			
(d) Witloof									► M70 0,02 (*) (P) ◀			

▼ **M45**

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
(e) Herbs									► M70 2 (P) ◀			
Chervil												
Chives												
Parsley												
Celery leaves												
Others												
(vi) LEGUME VEGETABLES (fresh)						► M66 0,05 (*) (P) ◀	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) (P) ◀	► M66 0,02 (*) (P) ◀	► M53 0,05 (*) (P) ◀	0,02 (*) (P)
Beans (with pods)												
Beans (without pods)												
Peas (with pods)												
Peas (without pods)												
Others												
(vii) STEM VEGETABLES (fresh)							► M70 0,01 (*) (P) ◀				► M53 0,05 (*) (P) ◀	0,02 (*) (P)
Asparagus												
Cardoons												
Celery												
Fennel												

▼ M45

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Globe artichokes										► M66 0,3 (P) ◀		
Leek						► M66 3 (P) ◀			► M70 0,5 (P) ◀			
Rhubarb												
Others						► M66 0,05 (*) (P) ◀			► M70 0,02 (*) (P) ◀	► M66 0,02 (*) (P) ◀		
(viii) FUNGI												0,02 (*) (P)
(a) Cultivated mushrooms											► M53 0,05 (*) (P) ◀	
(b) Wild mushrooms											► M53 20 (P) ◀	
3. Pulses	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)	0,01 (*) (P)	0,02 (P)	► M66 0,05 (*) (P) ◀	► M70 0,01 (*) (P) ◀	0,02 (*) (P)	► M70 0,3 (P) ◀	► M66 0,02 (*) (P) ◀	► M53 0,05 (*) (P) ◀	0,02 (*) (P)
Beans												
Lentils												
Peas												
Others												
4. Oil seed	0,1 (*) (P)	0,1 (*) (P)	0,02 (*) (P)	0,02 (*) (P)	0,05 (*) (P)	► M66 0,1 (*) (P) ◀	► M70 0,02 (*) (P) ◀	0,02 (*) (P)	► M70 0,05 (*) (P) ◀	► M66 0,05 (*) (P) ◀		0,05 (*) (P)
Linseed												

▼ **M45**

Groups and examples of individual products to which the MRLs apply	Bromoxynil including its esters expressed as bromoxynil	Chlorpropham (chlorpropham and 3-chloroaniline, expressed as chlorpropham) (**)	Dimethenamid-P including other mixtures of constituent isomers (sum of isomers)	Flazasulfuron	Flurtamone	Ioxynil, including its esters expressed as ioxynil	Mepanipyrim and its metabolite (2-anilino-4-(2-hydroxypropyl)-6-methylpyrimidine expressed as mepanipyrim	Propoxycarbazon, its salts and 2-hydroxypropoxycarbazon, calculated as propoxycarbazon	Pyraclostrobin	Quinoxyfen	Trimethylsulfonium cation resulting from the use of glyphosate	Zoxamide
Peanuts												
Poppy seeds												
Sesame seeds												
Sunflower seed												
Rape seed												
Soya bean											► M53 10 (P) ◄	
Mustard seed												
Cotton seed												
Others											► M53 0,05 (*) (P) ◄	
5. Potatoes	0,05 (*) (P)	10 (**) (P)	0,01 (*) (P)	0,01 (*) (P)	0,02 (*) (P)	► M66 0,05 (*) (P) ◄	► M70 0,01 (*) (P) ◄	0,02 (*) (P)	► M70 0,02 (*) (P) ◄	► M66 0,02 (*) ◄	► M53 0,05 (*) (P) ◄	0,02 (*) (P)
Early potatoes												
Ware potatoes												
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*) (P)	0,1 (*) (P)	0,02 (*) (P)	0,02 (*) (P)	0,05 (*) (P)	► M66 0,1 (*) (P) ◄	► M70 0,02 (*) (P) ◄	0,05 (*) (P)	► M70 0,05 (*) (P) ◄	► M66 0,05 (*) (P) ◄	► M53 0,05 (*) (P) ◄	0,05 (*) (P)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) (P)	0,1 (*) (P)	0,02 (*) (P)	0,02 (*) (P)	0,05 (*) (P)	► M66 0,1 (*) (P) ◄	► M70 0,02 (*) (P) ◄	0,05 (*) (P)	► M70 10 (P) ◄	► M66 0,5 (P) ◄	► M53 0,05 (*) (P) ◄	0,05 (*) (P)

(*) Indicates lower limit of analytical determination.

(**) Residue definition in potatoes is chlorpropham only.

► **M53** (P) Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC. ◄

▼ **M54**

Groups and examples of individual products to which the MRLs apply	Fenthion and its oxygen analogue, their sulfoxides and sulfones expressed as parent
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	
(i) CITRUS FRUIT Grapefruit Lemons Limes Mandarins (including clementines and other hybrids) Oranges Pomelos Others	3
(ii) TREE NUTS (shelled or unshelled) Almonds Brazil nuts Cashew nuts Chestnuts Coconuts Hazelnuts Macadamia Pecans Pine nuts Pistachios Walnuts Others	0,01 (*)
(iii) POME FRUIT Apples Pears Quinces Others	0,01 (*)
(iv) STONE FRUIT Apricots Cherries Peaches (including nectarines and similar hybrids) Plums Others	2 0,01 (*)
(v) BERRIES AND SMALL FRUIT (a) Table and wine grapes Table grapes Wine grapes (b) Strawberries (other than wild) (c) Cane fruit (other than wild) Blackberries Dewberries Loganberries Raspberries Others	0,01 (*)

▼ **M54**

Groups and examples of individual products to which the MRLs apply	Fenthion and its oxygen analogue, their sulfoxides and sulfones expressed as parent
(d) Other small fruit and berries (other than wild) Bilberries Cranberries Currants (red, black and white) Gooseberries Others	
(e) Wild berries and wild fruit	
(vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes olives (table consumption) olives (oil extraction) Papaya Passion fruit Pineapples Others	1 1 0,01 (*)
2. Vegetables, fresh or uncooked, frozen or dry	0,01 (*)
(i) ROOT AND TUBER VEGETABLES Beetroot Carrots Cassava Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips Yam Others	
(ii) BULB VEGETABLES Garlic Onions Shallots Spring onions Others	

▼ **M54**

Groups and examples of individual products to which the MRLs apply	Fenthion and its oxygen analogue, their sulfoxides and sulfones expressed as parent
<ul style="list-style-type: none"> (iii) FRUITING VEGETABLES <ul style="list-style-type: none"> (a) Solanacea <ul style="list-style-type: none"> Tomatoes Peppers Aubergines Okra Others (b) Cucurbits - edible peel <ul style="list-style-type: none"> Cucumbers Gherkins Courgettes Others (c) Cucurbits - inedible peel <ul style="list-style-type: none"> Melons Squashes Watermelons Others (d) Sweet corn (iv) BRASSICA VEGETABLES <ul style="list-style-type: none"> (a) Flowering brassica <ul style="list-style-type: none"> Broccoli Cauliflower Others (b) Head brassica <ul style="list-style-type: none"> Brussels sprouts Head cabbage Others (c) Leafy brassica <ul style="list-style-type: none"> Chinese cabbage Kale Others (d) Kohlrabi (v) LEAF VEGETABLES AND FRESH HERBS <ul style="list-style-type: none"> (a) Lettuce & similar <ul style="list-style-type: none"> Cress Lamb's lettuce Lettuce Scarole Ruccola Leaves and stems of brassica Others (b) Spinach & similar <ul style="list-style-type: none"> Spinach Beet leaves (chard) Others 	

▼ **M54**

Groups and examples of individual products to which the MRLs apply	Fenthion and its oxygen analogue, their sulfoxides and sulfones expressed as parent
<ul style="list-style-type: none"> (c) Water cress (d) Witloof (e) Herbs <ul style="list-style-type: none"> Chervil Chives Parsley Celery leaves Others (vi) LEGUME VEGETABLES (fresh) <ul style="list-style-type: none"> Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others (vii) STEM VEGETABLES (fresh) <ul style="list-style-type: none"> Asparagus Cardoons Celery Fennel Globe artichokes Leek Rhubarb Others (viii) FUNGI <ul style="list-style-type: none"> (a) Cultivated mushrooms (b) Wild mushrooms 	
<p>3. Pulses</p> <ul style="list-style-type: none"> Beans Lentils Peas Others 	0,01 (*)
<p>4. Oil seed</p> <ul style="list-style-type: none"> Linseed Peanuts Poppy seeds Sesame seeds Sunflower seed Rape seed Soya bean Mustard seed Cotton seed Hemp seed Others 	0,02 (*)

▼ **M54**

Groups and examples of individual products to which the MRLs apply	Fenthion and its oxygen analogue, their sulfoxides and sulfones expressed as parent
5. Potatoes Early potatoes Ware potatoes	0,01 (*)
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)

(*) Indicates lower limit of analytical determination.

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Captan	Folpet
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts		
(i) CITRUS FRUIT	0,02 (*)	► M63 0,02 (*) ◀
Grapefruit		
Lemons		
Limes		
Mandarins (including clementines and other hybrids)		
Oranges		
Pomelos		
Others		
(ii) TREE NUTS (shelled or unshelled)		► M63 0,02 (*) ◀
Almonds	0,3	
Brazil nuts		
Cashew nuts		
Chestnuts		
Coconuts		
Hazelnuts		
Macadamia		
Pecans		
Pine nuts		
Pistachios		
Walnuts		
Others	0,02 (*)	
(iii) POME FRUIT	3 ^(a)	► M63 3 ^(a) ◀
Apples		
Pears		
Quinces		
Others		
(iv) STONE FRUIT		
Apricots	3	
Cherries	5	► M63 2 ◀
Peaches (including nectarines and similar hybrids)		
Plums	1	
Others	0,02 (*)	► M63 0,02 (*) ◀
(v) BERRIES AND SMALL FRUIT		
(a) Table and wine grapes	0,02 (*)	
Table grapes		► M63 0,02 (*) ◀
Wine grapes		► M63 5 ◀

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Captan	Folpet
(b) Strawberries (other than wild)	3 (a)	► M63 3 (a) ◀
(c) Cane fruit (other than wild)		
Blackberries	3 (a)	► M63 3 (a) ◀
Dewberries		
Loganberries		
Raspberries	3 (a)	► M63 3 (a) ◀
Others	0,02 (*)	► M63 0,02 (*) ◀
(d) Other small fruit and berries (other than wild)		
Bilberries		
Cranberries		
Currants (red, black and white)	3 (a)	► M63 3 (a) ◀
Gooseberries	3 (a)	► M63 3 (a) ◀
Others	0,02 (*)	► M63 0,02 (*) ◀
(e) Wild berries and wild fruit	0,02 (*)	► M63 0,02 (*) ◀
(vi) MISCELLANEOUS		► M63 0,02 (*) ◀
Avocados		
Bananas		
Dates		
Figs		
Kiwi		
Kumquats		
Litchis		
Mangoes	2	
Olives (table consumption)		
Olives (oil extraction)		
Papaya		
Passion fruit		
Pineapples		
Pomegranate		
Others	0,02 (*)	
2. Vegetables, fresh or uncooked, frozen or dry		
(i) ROOT AND TUBER VEGETABLES		► M63 0,02 (*) ◀
Beetroot		
Carrots	0,1	
Cassava		
Celeriac	0,1	
Horseradish		

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Captan	Folpet
Jerusalem artichokes		
Parsnips		
Parsley root		
Radishes		
Salsify		
Sweet potatoes		
Swedes		
Turnips		
Yam		
Others	0,02 (*)	
(ii) BULB VEGETABLES	0,02 (*)	
Garlic		
Onions		► M63 0,1 ◀
Shallots		
Spring onions		
Others		► M63 0,02 (*) ◀
(iii) FRUITING VEGETABLES		
(a) Solanacea		
Tomatoes	2 (a)	► M63 2 (a) ◀
Peppers	0,1	
Aubergines		
Okra		
Others	0,02 (*)	► M63 0,02 (*) ◀
(b) Cucurbits — edible peel	0,02 (*)	► M63 0,02 (*) ◀
Cucumbers		
Gherkins		
Courgettes		
Others		
(c) Cucurbits — inedible peel		► M63 1 ◀
Melons	0,1	
Squashes		
Watermelons		
Others	0,02 (*)	
(d) Sweetcorn	0,02 (*)	► M63 0,02 (*) ◀
(iv) BRASSICA VEGETABLES	0,02 (*)	
(a) Flowering brassica		► M63 0,02 (*) ◀
Broccoli		
Cauliflower		

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Captan	Folpet
Others		
(b) Head brassica		► M63 0,02 (*) ◀
Brussels sprouts		
Head cabbage		
Others		
(c) Leafy brassica		► M63 0,02 (*) ◀
Chinese cabbage		
Kale		
Others		
(d) Kohlrabi		► M63 0,05 ◀
(v) LEAF VEGETABLES AND FRESH HERBS		
(a) Lettuce and similar		
Cress		
Lamb's lettuce		
Lettuce		► M63 2 ◀
Scarole	2	
Ruccola		
Leaves and stems of brassica		
Others	0,02 (*)	► M63 0,02 (*) ◀
(b) Spinach and similar		
Spinach	0,1	► M63 10 ◀
Beet leaves (chard)		
Others	0,02 (*)	► M63 0,02 (*) ◀
(c) Watercress	0,02 (*)	► M63 0,02 (*) ◀
(d) Witloof	0,02 (*)	► M63 0,02 (*) ◀
(e) Herbs		► M63 0,02 (*) ◀
Chervil		
Chives		
Parsley	0,1	
Celery leaves		
Others	0,02 (*)	
(vi) LEGUME VEGETABLES (fresh)		
Beans (with pods)	2 ^(a)	► M63 2 ^(a) ◀
Beans (without pods)	2 ^(a)	► M63 2 ^(a) ◀
Peas (with pods)		
Peas (without pods)		
Others	0,02 (*)	► M63 0,02 (*) ◀

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Captan	Folpet
(vii) STEM VEGETABLES (fresh)		► M63 0,02 (*) ◀
Asparagus		
Cardoons		
Celery	0,1	
Fennel		
Globe artichokes		
Leek	2	
Rhubarb		
Others	0,02 (*)	
(viii) FUNGI	0,02 (*)	► M63 0,02 (*) ◀
(a) Cultivated mushrooms		
(b) Wild mushrooms		
3. Pulses	0,02 (*)	► M63 0,02 (*) ◀
Beans		
Lentils		
Peas		
Lupines		
Others		
4. Oil seed	0,02 (*)	► M63 0,05 (*) ◀
Linseed		
Peanuts		
Poppy seeds		
Sesame seeds		
Sunflower seed		
Rapeseed		
Soya bean		
Mustard seed		
Cotton seed		
Hemp seed		
Others		
5. Potatoes	0,05	► M63 0,1 ◀
Early potatoes		
Ware potatoes		
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,05 (*)	► M63 0,05 (*) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (*)	► M63 150 ◀

(*) Sum of captan and folpet.

(*) Indicates lower limit of analytical determination.

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Dichlorvos	Ethion
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,01 (*)	0,01 (*)
(i) CITRUS FRUIT		
Grapefruit		
Lemons		
Limes		
Mandarins (including clementines and other hybrids)		
Oranges		
Pomelos		
Others		
(ii) TREE NUTS (shelled or unshelled)		
Almonds		
Brazil nuts		
Cashew nuts		
Chestnuts		
Coconuts		
Hazelnuts		
Macadamia		
Pecans		
Pine nuts		
Pistachios		
Walnuts		
Others		
(iii) POME FRUIT		
Apples		
Pears		
Quinces		
Others		
(iv) STONE FRUIT		
Apricots		
Cherries		
Peaches (including nectarines and similar hybrids)		
Plums		
Others		

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Dichlorvos	Ethion
(v) BERRIES AND SMALL FRUIT		
(a) Table and wine grapes		
Table grapes		
Wine grapes		
(b) Strawberries (other than wild)		
(c) Cane fruit (other than wild)		
Blackberries		
Dewberries		
Loganberries		
Raspberries		
Others		
(d) Other small fruit and berries (other than wild)		
Bilberries		
Cranberries		
Currants (red, black and white)		
Gooseberries		
Others		
(e) Wild berries and wild fruit		
(vi) MISCELLANEOUS		
Avocados		
Bananas		
Dates		
Figs		
Kiwi		
Kumquats		
Litchis		
Mangoes		
Olives (table consumption)		
Olives (oil extraction)		
Papaya		
Passion fruit		
Pineapples		
Pomegranate		
Others		

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Dichlorvos	Ethion
2. Vegetables, fresh or uncooked, frozen or dry	0,01 (*)	
(i) ROOT AND TUBER VEGETABLES		0,01 (*)
Beetroot		
Carrots		
Cassava		
Celeriac		
Horseradish		
Jerusalem artichokes		
Parsnips		
Parsley root		
Radishes		
Salsify		
Sweet potatoes		
Swedes		
Turnips		
Yam		
Others		
(ii) BULB VEGETABLES		0,01 (*)
Garlic		
Onions		
Shallots		
Spring onions		
Others		
(iii) FRUITING VEGETABLES		0,01 (*)
(a) Solanacea		
Tomatoes		
Peppers		
Aubergines		
Okra		
Others		
(b) Cucurbits — edible peel		
Cucumbers		
Gherkins		
Courgettes		
Others		
(c) Cucurbits — inedible peel		
Melons		
Squashes		

▼ M56

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Dichlorvos	Ethion
Watermelons		
Others		
(d) Sweetcorn		
(iv) BRASSICA VEGETABLES		0,01 (*)
(a) Flowering brassica		
Broccoli		
Cauliflower		
Others		
(b) Head brassica		
Brussels sprouts		
Head cabbage		
Others		
(c) Leafy brassica		
Chinese cabbage		
Kale		
Others		
(d) Kohlrabi		
(v) LEAF VEGETABLES AND FRESH HERBS		
(a) Lettuce and similar		0,01 (*)
Cress		
Lamb's lettuce		
Lettuce		
Scarole		
Ruccola		
Leaves and stems of brassica		
Others		
(b) Spinach and similar		0,01 (*)
Spinach		
Beet leaves (chard)		
Others		
(c) Watercress		0,01 (*)
(d) Witloof		0,01 (*)
(e) Herbs		
Chervil		
Chives		
Parsley		2
Celery leaves		
Others		0,01 (*)

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Dichlorvos	Ethion
(vi) LEGUME VEGETABLES (fresh)		0,01 (*)
Beans (with pods)		
Beans (without pods)		
Peas (with pods)		
Peas (without pods)		
Others		
(vii) STEM VEGETABLES (fresh)		
Asparagus		
Cardoons		
Celery		0,1
Fennel		
Globe artichokes		
Leek		
Rhubarb		
Others		0,01 (*)
(viii) FUNGI		0,01 (*)
(a) Cultivated mushrooms		
(b) Wild mushrooms		
3. Pulses	0,01 (*)	0,01 (*)
Beans		
Lentils		
Peas		
Lupines		
Others		
4. Oil seed	0,01 (*)	0,02 (*)
Linseed		
Peanuts		
Poppy seeds		
Sesame seeds		
Sunflower seed		
Rapeseed		
Soya bean		
Mustard seed		
Cotton seed		
Hemp seed		
Others		

▼ **M56**

Pesticide residues and maximum residue levels (mg/kg)		
Groups and examples of individual products to which the MRLs apply	Dichlorvos	Ethion
5. Potatoes Early potatoes Ware potatoes	0,01 (*)	0,01 (*)
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,02 (*)	3
7. Hops (dried), including hop pellets and unconcentrated powder	0,02 (*)	0,02 (*)

(*) Indicates lower limit of analytical determination.

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Carbaryl	Oxamyl ^(b)
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts		
(i) CITRUS FRUIT	0,05 (*)	
Grapefruit		
Lemons		
Limes		
Mandarins (including clementines and other hybrids)		▶ M66 0,02 (*) (P) ◀
Oranges		
Pomelos		
Others		▶ M66 0,01 (*) (P) ◀
(ii) TREE NUTS (shelled or unshelled)	0,05 (*)	▶ M66 0,01 (*) (P) ◀
Almonds		
Brazil nuts		
Cashew nuts		
Chestnuts		
Coconuts		
Hazelnuts		
Macadamia		
Pecans		
Pine nuts		
Pistachios		
Walnuts		
Others		
(iii) POME FRUIT	0,05 (*)	▶ M66 0,01 (*) (P) ◀
Apples		
Pears		
Quinces		
Others		
(iv) STONE FRUIT	0,05 (*)	▶ M66 0,01 (*) (P) ◀
Apricots		
Cherries		
Peaches (including nectarines and similar hybrids)		
Plums		
Others		
(v) BERRIES AND SMALL FRUIT	0,05 (*)	▶ M66 0,01 (*) (P) ◀
(a) Table and wine grapes		
Table grapes		
Wine grapes		
(b) Strawberries (other than wild)		
(c) Cane fruit (other than wild)		
Blackberries		
Dewberries		

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Carbaryl	Oxamyl (*)
Loganberries Raspberries Others (d) Other small fruit and berries (other than wild) Bilberries Cranberries Currants (red, black and white) Gooseberries Others (e) Wild berries and wild fruit		
(vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwis Kumquats Litchis Mangoes Olives (table consumption) Olives (oil extraction) Papayas Passion fruit Pineapples Pomegranate Others	5 5 0,05 (*)	► M66 0,01 (*) (P) ◀
2. Vegetables, fresh or uncooked, frozen or dry (i) ROOT AND TUBER VEGETABLES Beetroot Carrots Cassava Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips Yam Others	0,05 (*)	► M66 0,01 (*) (P) ◀

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Carbaryl	Oxamyl (b)
Ruccola		
Leaves and stems of brassica		
Others		
(b) Spinach & similar		
Spinach		
Beet leaves (chard)		
Others		
(c) Watercress		
(d) Witloof		
(e) Herbs		
Chervil		
Chives		
Parsley		
Celery leaves		
Others		
(vi) LEGUME VEGETABLES (fresh)	0,05 (*)	► M66 0,01 (*) (P) ◀
Beans (with pods)		
Beans (without pods)		
Peas (with pods)		
Peas (without pods)		
Others		
(vii) STEM VEGETABLES (fresh)	0,05 (*)	► M66 0,01 (*) (P) ◀
Asparagus		
Cardoons		
Celery		
Fennel		
Globe artichokes		
Leeks		
Rhubarb		
Others		
(viii) FUNGI	0,05 (*)	► M66 0,01 (*) (P) ◀
(a) Cultivated mushrooms		
(b) Wild mushrooms		
3. Pulses	0,05 (*)	► M66 0,01 (*) (P) ◀
Beans		
Lentils		
Peas		
Others		
4. Oil seed	0,05 (*)	► M66 0,02 (*) (P) ◀
Linseed		
Peanuts		
Poppy seeds		
Sesame seeds		
Sunflower seeds		

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Carbaryl	Oxamyl ^(b)
Rape-seed Soya beans Mustard seeds Cotton seed Hemp seed Others		
5. Potatoes Early potatoes Ware potatoes	0,05 (*)	► M66 0,01 (*) ^(p) ◀
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,1 (*)	► M66 0,02 (*) ^(p) ◀
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)	► M66 0,02 (*) ^(p) ◀

(*) Indicates lower limit of analytical determination.

^(p) ► **M66** Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC. ◀

^(b) Temporary MRL valid until 1 January 2008, pending submission of trial data.

▼ **M55**

Pesticide residues and maximum residue levels (mg/kg)			
Groups and examples of individual products to which the MRLs would apply	Desmedipham	Phenmedipham	Chlorfenvinphos (sum of E- and Z-isomers)
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,05 (*) (P)		0,02 (*)
(i) CITRUS FRUIT		► M57 0,05 (*) (P) ◀	
Grapefruit			
Lemons			
Limes			
Mandarins (including clementines and other hybrids)			
Oranges			
Pomelos			
Others			
(ii) TREE NUTS (shelled or unshelled)		► M57 0,05 (*) (P) ◀	
Almonds			
Brazil nuts			
Cashew nuts			
Chestnuts			
Coconuts			
Hazelnuts			
Macadamia			
Pecans			
Pine nuts			
Pistachios			
Walnuts			
Others			
(iii) POME FRUIT		► M57 0,05 (*) (P) ◀	
Apples			
Pears			
Quinces			
Others			
(iv) STONE FRUIT		► M57 0,05 (*) (P) ◀	
Apricots			
Cherries			
Peaches (including nectarines and similar hybrids)			
Plums			
Others			
(v) BERRIES & SMALL FRUIT			
(a) Table and wine grapes		► M57 0,05 (*) (P) ◀	
Table grapes			
Wine grapes			

▼ **M55**

Pesticide residues and maximum residue levels (mg/kg)			
Groups and examples of individual products to which the MRLs would apply	Desmedipham	Phenmedipham	Chlorfenvinphos (sum of E- and Z-isomers)
(b) Strawberries (other than wild)		► M57 0,1 (P) ◀	
(c) Cane fruit (other than wild)		► M57 0,05 (*) (P) ◀	
Blackberries			
Dewberries			
Loganberries			
Raspberries			
Others			
(d) Other small fruit & berries (other than wild)		► M57 0,05 (*) (P) ◀	
Bilberries			
Cranberries			
Currants (red, black and white)			
Gooseberries			
Others			
(e) Wild berries and wild fruit		► M57 0,05 (*) (P) ◀	
(vi) MISCELLANEOUS		► M57 0,05 (*) (P) ◀	
Avocados			
Bananas			
Dates			
Figs			
Kiwi			
Kumquats			
Litchis			
Mangoes			
Olives (table consumption)			
Olives (oil extraction)			
Papaya			
Passion fruit			
Pineapples			
Pomegranate			
Others			
2. Vegetables, fresh or uncooked, frozen or dry	0,05 (*) (P)		
(i) ROOT AND TUBER VEGETABLES			
Beetroot		► M57 0,1 (P) ◀	
Carrots			0,5
Cassava			
Celeriac			
Horseradish			
Jerusalem artichokes			

▼ **M55**

Pesticide residues and maximum residue levels (mg/kg)			
Groups and examples of individual products to which the MRLs would apply	Desmedipham	Phenmedipham	Chlorfenvinphos (sum of E- and Z-isomers)
Parsnips			0,5
Parsley root			
Radishes			0,5
Salsify			
Sweet potatoes			
Swedes			0,5
Turnips			0,5
Yam			
Others		► M57 0,05 (*) (P) ◀	0,02 (*)
(ii) BULB VEGETABLES		► M57 0,05 (*) (P) ◀	
Garlic			0,5
Onions			
Shallots			0,5
Spring onions			
Others			0,02 (*)
(iii) FRUITING VEGETABLES		► M57 0,05 (*) (P) ◀	
(a) Solanacea			0,02 (*)
Tomatoes			
Peppers			
Aubergines			
Okra			
Others			
(b) Cucurbits - edible peel			
Cucumbers			
Gherkins			
Courgettes			0,1
Others			0,02 (*)
(c) Cucurbits-inedible peel			0,02 (*)
Melons			
Squashes			
Watermelons			
Others			
(d) Sweet corn			0,02 (*)
(iv) BRASSICA VEGETABLES		► M57 0,05 (*) (P) ◀	
(a) Flowering brassica			0,02 (*)
Broccoli (including Calabrese)			
Cauliflower			
Others			
(b) Head brassica			
Brussels sprouts			0,1

▼ **M55**

Pesticide residues and maximum residue levels (mg/kg)			
Groups and examples of individual products to which the MRLs would apply	Desmedipham	Phenmedipham	Chlorfenvinphos (sum of E- and Z-isomers)
Head cabbage			0,5
Others			0,02 (*)
(c) Leafy brassica			0,02 (*)
Chinese cabbage			
Kale			
Others			
(d) Kohlrabi			0,3
(v) LEAF VEGETABLES & FRESH HERBS			
(a) Lettuce & similar		► M57 0,05 (*) (P) ◀	
Cress			0,1
Lamb's lettuce			0,1
Lettuce			
Scarole (broad-leaf endive)			
Ruccola			
Leaves and stems of brassica			
Others			0,02 (*)
(b) Spinach & similar		► M57 0,5 (P) ◀	
Spinach			0,1
Beet leaves (chard)			
Others			0,02 (*)
(c) Water cress		► M57 0,05 (*) (P) ◀	0,02 (*)
(d) Witloof		► M57 0,05 (*) (P) ◀	0,02 (*)
(e) Herbs		► M57 7 ◀	
Chervil			
Chives			
Parsley			0,5
Celery leaves			
Others			0,02 (*)
(vi) LEGUME VEGETABLES (fresh)		► M57 0,05 (*) (P) ◀	0,02 (*)
Beans (with pods)			
Beans (without pods)			
Peas (with pods)			
Peas (without pods)			
Others			
(vii) STEM VEGETABLES (fresh)			
Asparagus			0,1
Cardoons			
Celery			0,5

▼ **M55**

Pesticide residues and maximum residue levels (mg/kg)			
Groups and examples of individual products to which the MRLs would apply	Desmedipham	Phenmedipham	Chlorfenvinphos (sum of E- and Z-isomers)
Fennel			
Globe artichokes		► M57 0,2 (P) ◀	
Leek			0,1
Rhubarb			
Others		► M57 0,05 (*) (P) ◀	0,02 (*)
(viii) FUNGI		► M57 0,05 (*) (P) ◀	
(a) Cultivated mushrooms			0,05
(b) Wild mushrooms			0,02 (*)
3. Pulses	0,05 (*) (P)	► M57 0,05 (*) (P) ◀	0,02 (*)
Beans			
Lentils			
Peas			
Others			
4. Oilseeds	0,1 (*) (P)	► M57 0,1 (*) (P) ◀	0,02 (*)
Linseed			
Peanuts			
Poppy seed			
Sesame seed			
Sunflower seed			
Rape seed			
Soya bean			
Mustard seed			
Cotton seed			
Hemp seed			
Others			
5. Potatoes	0,05 (*) (P)	► M57 0,05 (*) (P) ◀	0,02 (*)
Early potatoes			
Ware potatoes			
6. Tea (dried leaves and stalks, fermented or other-wise, <i>Camellia sinensis</i>)	0,1 (*) (P)	► M57 0,1 (*) (P) ◀	0,05 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*) (P)	► M57 0,1 (*) (P) ◀	0,05 (*)

(*) Indicates lower limit of analytical determination.

► **M57** (P) Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 7 March 2011. ◀

▼ **M59**

Pesticide residue and maximum residue level (mg/kg)		
Groups and examples of individual products to which the MRLs would apply	Phosphamidon	Mevinphos, sum of E- and Z-isomers
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts	0,01 (*)	0,01 (*)
(i) CITRUS FRUIT Grapefruit Lemons Limes Mandarins (including clementines and other hybrids) Oranges Pomelos Others		
(ii) TREE NUTS (shelled or unshelled) Almonds Brazil nuts Cashew nuts Chestnuts Coconuts Hazelnuts Macadamia Pecans Pine nuts Pistachios Walnuts Others		
(iii) POME FRUIT Apples Pears Quinces Others		
(iv) STONE FRUIT Apricots Cherries Peaches (including nectarines and similar hybrids) Plums Others		
(v) BERRIES AND SMALL FRUIT (a) Table and wine grapes Table grapes Wine grapes (b) Strawberries (other than wild)		

▼ **M59**

Pesticide residue and maximum residue level (mg/kg)		
Groups and examples of individual products to which the MRLs would apply	Phosphamidon	Mevinphos, sum of E- and Z-isomers
(c) Cane fruit (other than wild)		
Blackberries		
Dewberries		
Loganberries		
Raspberries		
Others		
(d) Other small fruit and berries (other than wild)		
Bilberries		
Cranberries		
Currants (red, black and white)		
Gooseberries		
Others		
(e) Wild berries and wild fruit		
(vi) MISCELLANEOUS		
Avocados		
Bananas		
Dates		
Figs		
Kiwi		
Kumquats		
Litchis		
Mangoes		
Olives (table consumption)		
Olives (oil extraction)		
Papaya		
Passion fruit		
Pineapples		
Pomegranate		
Others		
2. Vegetables, fresh or uncooked, frozen or dry	0,01 (*)	0,01 (*)
(i) ROOT AND TUBER VEGETABLES		
Beetroot		
Carrots		
Cassava		
Celeriac		
Horseradish		
Jerusalem artichokes		
Parsnips		
Parsley root		

▼ **M59**

Pesticide residue and maximum residue level (mg/kg)		
Groups and examples of individual products to which the MRLs would apply	Phosphamidon	Mevinphos, sum of E- and Z-isomers
Radishes		
Salsify		
Sweet potatoes		
Swedes		
Turnips		
Yam		
Others		
(ii) BULB VEGETABLES		
Garlic		
Onions		
Shallots		
Spring onions		
Others		
(iii) FRUITING VEGETABLES		
(a) Solanacea		
Tomatoes		
Peppers		
Aubergines		
Okra		
Others		
(b) Cucurbits — edible peel		
Cucumbers		
Gherkins		
Courgettes		
Others		
(c) Cucurbits-inedible peel		
Melons		
Squashes		
Watermelons		
Others		
(d) Sweet corn		
(iv) BRASSICA VEGETABLES		
(a) Flowering brassica		
Broccoli (including Calabrese)		
Cauliflower		
Others		
(b) Head brassica		
Brussels sprouts		
Head cabbage		
Others		

▼ **M59**

Pesticide residue and maximum residue level (mg/kg)		
Groups and examples of individual products to which the MRLs would apply	Phosphamidon	Mevinphos, sum of E- and Z-isomers
(c) Leafy brassica Chinese cabbage Kale Others		
(d) Kohlrabi		
(v) LEAF VEGETABLES AND FRESH HERBS		
(a) Lettuce and similar Cress Lamb's lettuce Lettuce Scarole (broad-leaf endive) Ruccola Leaves and stems of brassica Others		
(b) Spinach and similar Spinach Beet leaves (chard) Others		
(c) Water cress		
(d) Witloof		
(e) Herbs Chervil Chives Parsley Celery leaves Others		
(vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others		
(vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek		

▼ **M59**

Pesticide residue and maximum residue level (mg/kg)		
Groups and examples of individual products to which the MRLs would apply	Phosphamidon	Mevinphos, sum of E- and Z-isomers
Rhubarb		
Others		
(viii) FUNGI		
(a) Cultivated mushrooms		
(b) Wild mushrooms		
3. Pulses	0,01 (*)	0,01 (*)
Beans		
Lentils		
Peas		
Lupines		
Others		
4. Oilseeds	0,01 (*)	0,01 (*)
Linseed		
Peanuts		
Poppy seed		
Sesame seed		
Sunflower seed		
Rape seed		
Soya bean		
Mustard seed		
Cotton seed		
Hemp seed		
Others		
5. Potatoes	0,01 (*)	0,01 (*)
Early potatoes		
Ware potatoes		
6. Tea (dried leaves and stalks, fermented or other-wise, <i>Camellia sinensis</i>)	0,02 (*)	0,02 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,02 (*)	0,02 (*)

(*) Indicates lower limit of analytical determination.

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts		0,01 (*) (P)			0,05 (*) (P)		0,01 (*) (P)
(i) CITRUS FRUIT	► M70 1 (P) ◀		► M70 1 ◀	0,05 (*) (P)		► M70 0,02 (*) (P) ◀	
Grapefruit							
Lemons							
Limes							
Mandarins (including clementines and other hybrids)							
Oranges							
Pomelos							
Others							
(ii) TREE NUTS (shelled or unshelled)	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀	0,1 (*) (P)		► M70 0,02 (*) (P) ◀	
Almonds							
Brazil nuts							
Cashew nuts							
Chestnuts							
Coconuts							
Hazelnuts							
Macadamia							
Pecans							

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamidrid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
Pine nuts							
Pistachios							
Walnuts							
Others							
(iii) POME FRUIT	► M70 0,1 (P) ◀		► M70 2 ◀	0,05 (*) (P)		► M70 0,3 (P) ◀	
Apples							
Pears							
Quinces							
Others							
(iv) STONE FRUIT				0,05 (*) (P)			
Apricots	► M70 0,1 (P) ◀		► M70 0,3 ◀			► M70 0,3 (P) ◀	
Cherries	► M70 0,2 (P) ◀					► M70 0,3 (P) ◀	
Peaches (including nectarines and similar hybrids)	► M70 0,1 (P) ◀		► M70 0,3 ◀			► M70 0,3 (P) ◀	
Plums	► M70 0,02 (P) ◀					► M70 0,1 (P) ◀	
Others	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀			► M70 0,02 (*) (P) ◀	

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamidrid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
(v) BERRIES AND SMALL FRUIT	► M70 0,01 (*) (P) ◀			0,05 (*) (P)			
(a) Table and wine grapes			► M70 1 ◀			► M70 0,02 (*) (P) ◀	
Table grapes			► M70 1 ◀				
Wine grapes			► M70 1 ◀				
(b) Strawberries (other than wild)			► M70 0,02 (*) ◀			► M70 0,5 (P) ◀	
(c) Cane fruit (other than wild)			► M70 0,02 (*) ◀				
Blackberries						► M70 3 (P) ◀	
Dewberries							
Loganberries							
Raspberries						► M70 3 (P) ◀	
Others						► M70 1 (P) ◀	
(d) Other small fruit and berries (other than wild)			► M70 0,02 (*) ◀			► M70 1 (P) ◀	
Bilberries							
Cranberries							
Currants (red, black and white)							
Gooseberries							
Others							

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
(e) Wild berries and wild fruit						► M70 0,02 (*) (P) ◀	
(vi) MISCELLANEOUS	► M70 0,01 (*) (P) ◀			0,05 (*) (P)			
Avocados							
Bananas							
Dates							
Figs							
Kiwi			► M70 1 ◀				
Kumquats							
Litchis							
Mangoes							
olives (table consumption)							
olives (oil extraction)							
Papaya						► M70 0,5 (P) ◀	
Passion fruit							
Pineapples							
Pomegranate							
Others			► M70 0,02 (*) (P) ◀			► M70 0,02 (*) (P) ◀	

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamidid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
2. Vegetables, fresh or uncooked, frozen or dry		0,01 (*) (P)		0,05 (*) (P)	0,05 (*) (P)		0,01 (*) (P)
(i) ROOT AND TUBER VEGETABLES	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀				
Beetroot							
Carrots							
Cassava							
Celeriac						► M70 0,1 (P) ◀	
Horseradish							
Jerusalem artichokes							
Parsnips							
Parsley root							
Radishes							
Salsify							
Sweet potatoes							
Swedes							
Turnips							
Yam							
Others							

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
(ii) BULB VEGETABLES	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀			► M70 0,02 (*) (P) ◀	
Garlic							
Onions							
Shallots							
Spring onions							
Others							
(iii) FRUITING VEGETABLES							
(a) Solanacea							
Tomatoes	► M70 0,1 (P) ◀		► M70 2 ◀			► M70 0,5 (P) ◀	
Peppers	► M70 0,3 (P) ◀		► M70 1 ◀			► M70 1 (P) ◀	
Aubergines	► M70 0,1 (P) ◀		► M70 0,5 ◀			► M70 0,5 (P) ◀	
Okra							
Others	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀			► M70 0,02 (*) (P) ◀	
(b) Cucurbits — edible peel	► M70 0,3 (P) ◀		► M70 0,02 (*) ◀			► M70 0,3 (P) ◀	
Cucumbers							
Gherkins							
Courgettes							
Others							

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
(c) Cucurbits — inedible peel	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀				
Melons						► M70 0,2 (P) ◀	
Squashes							
Watermelons						► M70 0,2 (P) ◀	
Others						► M70 0,02 (*) (P) ◀	
(d) Sweet corn	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀			► M70 0,1 (P) ◀	
(iv) BRASSICA VEGETABLES			► M70 0,02 (*) ◀				
(a) Flowering brassica	► M70 0,01 (*) ◀					► M70 0,1 (P) ◀	
Broccoli							
Cauliflower							
Others							
(b) Head brassica							
Brussels sprouts	► M70 0,05 ◀					► M70 0,05 (P) ◀	
Head cabbage						► M70 0,2 (P) ◀	
Others	► M70 0,01 (*) ◀					► M70 0,02 (*) (P) ◀	

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
(c) Leafy brassica	► M70 0,01 (*) ◀					► M70 1 (P) ◀	
Chinese cabbage							
Kale							
Others							
(d) Kohlrabi	► M70 0,01 (*) ◀					► M70 0,05 (P) ◀	
(v) LEAF VEGETABLES AND FRESH HERBS				► M70 0,02 (*) ◀			
(a) Lettuce and similar							
Cress							
Lamb's lettuce	► M70 5 ◀						
Lettuce	► M70 5 ◀						
Scarole	► M70 5 ◀						
Ruccola	► M70 5 ◀					► M70 3 (P) ◀	
Leaves and stems of brassica	► M70 5 ◀						
Others	► M70 0,01 (*) (P) ◀					► M70 2 (P) ◀	
(b) Spinach and similar	► M70 0,01 (*) (P) ◀					► M70 0,02 (*) (P) ◀	
Spinach							
Beet leaves (chard)							
Others							

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
(c) Water cress	► M70 0,01 (*) (P) ◀					► M70 0,02 (*) (P) ◀	
(d) Witloof	► M70 0,01 (*) (P) ◀					► M70 0,02 (*) (P) ◀	
(e) Herbs						► M70 3 (P) ◀	
Chervil							
Chives							
Parsley	► M70 5 ◀						
Celery leaves							
Others	► M70 0,01 (*) (P) ◀						
(vi) LEGUME VEGETABLES (fresh)	► M70 0,01 (*) (P) ◀						
Beans (with pods)			► M70 0,2 ◀			► M70 1 (P) ◀	
Beans (without pods)							
Peas (with pods)							
Peas (without pods)						► M70 0,2 (P) ◀	
Others			► M70 0,02 (*) ◀			► M70 0,02 (*) (P) ◀	
(vii) STEM VEGETABLES (fresh)	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀				
Asparagus							

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
Cardoons							
Celery						► M70 0,3 (*) ◀	
Fennel							
Globe artichokes							
Leek							
Rhubarb							
Others			► M70 0,02 (*) ◀			► M70 0,02 (*) (P) ◀	
(viii) FUNGI	► M70 0,01 (*) (P) ◀		► M70 0,02 (*) ◀			► M70 0,02 (*) (P) ◀	
(a) Cultivated mushrooms							
(b) Wild mushrooms							
3. Pulses	► M70 0,01 (*) (P) ◀	0,01 (*) (P)	► M70 0,02 (*) ◀	0,05 (*) (P)	0,05 (*) (P)	► M70 0,1 (P) ◀	0,01 (*) (P)
Beans							
Lentils							
Peas							
Lupines							
Others							
4. Oilseed		0,01 (*) (P)		0,1 (*) (P)	0,1 (*) (P)		0,01 (*) (P)
Linseed							
Peanuts							

▼ **M61**

Pesticide residue and maximum residue level (mg/kg)							
Groups and examples of individual products to which the MRLs apply	Acetamiprid	Imazosulfuron	Methoxyfenozide	Sum of MA4 + 8,9Z-MA4, expressed as milbemectin	Metholachlor including other mixtures of constituent isomers including S-metolachlor (Sum of isomers)	Thiacloprid	Tribenuron — methyl
Poppy seeds							
Sesame seeds							
Sunflower seed							
Rape seed						► M70 0,3 (*) ◀	
Soya bean			► M70 2 ◀				
Mustard seed						► M70 0,2 (*) ◀	
Cotton seed	► M70 0,02 (*) ◀		► M70 2 ◀				
Hemp seed							
Others	► M70 0,01 (*) (*) ◀		► M70 0,05 (*) ◀			► M70 0,05 (*) (*) ◀	
5. Potatoes	► M70 0,01 (*) (*) ◀	0,01 (*) (*)	► M70 0,02 (*) ◀	0,05 (*) (*)	0,05 (*) (*)	► M70 0,02 (*) (*) ◀	0,01 (*) (*)
Early potatoes							
Ware potatoes							
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	► M70 0,1 (*) (*) ◀	0,02 (*)	► M70 0,05 (*) ◀	0,1 (*) (*)	0,1 (*) (*)	► M70 0,05 (*) (*) ◀	0,02 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	► M70 0,1 (*) (*) ◀	0,02 (*)	► M70 0,05 (*) ◀	0,1 (*) (*)	0,1 (*) (*)	► M70 0,1 ◀	0,02 (*)

(*) Indicates lower limit of analytical determination.

(P) ► **M70** Indicates that the maximum residue level has been established provisionally in accordance with Article 4(1)(f) of Directive 91/414/EEC. ◀

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as mesosulfuron	Triticonazole	1-methylcyclopropene
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts			0,05 (*) (P)		0,01 (*) (P)	0,01 (*) (P)	0,01 (*) (P)
(i) CITRUS FRUIT	0,1 (P)	► M70 0,02 (*) (P) ◀		0,05 (*) (P)			
Grapefruit							
Lemons							
Limes							
Mandarins (including clementines and other hybrids)							
Oranges							
Pomelos							
Others							
(ii) TREE NUTS (shelled or unshelled)	0,02 (*) (P)	► M70 0,05 (*) (P) ◀		0,05 (*) (P)			
Almonds							
Brazil nuts							
Cashew nuts							
Chestnuts							
Coconuts							
Hazelnuts							
Macadamia							
Pecans							
Pine nuts							
Pistachios							

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as mesosulfuron	Triticonazole	1-methylcyclopropene
Walnuts							
Others							
(iii) POME FRUIT	0,02 (*) (P)			3 (P)			
Apples		► M70 0,5 (P) ◀					
Pears							
Quinces							
Others		► M70 0,3 (P) ◀					
(iv) STONE FRUIT							
Apricots	0,1 (P)	► M70 0,3 (P) ◀					
Cherries				1 (P)			
Peaches (including nectarines and similar hybrids)	0,1 (P)	► M70 0,3 (P) ◀					
Plums				0,5 (P)			
Others	0,02 (*) (P)	0,02 (*) (P)		0,05 (*) (P)			
(v) BERRIES AND SMALL FRUIT							
(a) Table and wine grapes	0,02 (*) (P)	► M70 2 (P) ◀		5 (P)			
Table grapes							
Wine grapes							

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as meso-sulfuron	Triticonazole	1-methylcyclopropene
(b) Strawberries (other than wild)	0,2 (P)	► M70 0,02 (*) (P) ◀		5 (P)			
(c) Cane fruit (other than wild)	0,02 (*) (P)	► M70 0,02 (*) (P) ◀		5 (P)			
Blackberries							
Dewberries							
Loganberries							
Raspberries							
Others							
(d) Other small fruit and berries (other than wild)	0,02 (*) (P)			5 (P)			
Bilberries							
Cranberries							
Currants (red, black and white)		► M70 1 (P) ◀					
Gooseberries		► M70 1 (P) ◀					
Others		► M70 0,02 (*) (P) ◀					
(e) Wild berries and wild fruit	► C13 0,02 (*) (P) ◀	► M70 0,02 (*) (P) ◀		0,05 (*) (P)			
(vi) MISCELLANEOUS	0,02 (*) (P)			0,05 (*) (P)			
Avocados							
Bananas		► M70 0,2 (P) ◀					
Dates							

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolylfluanid and dimethylaminosulfotoluidide expressed as tolylfluanid)	Mesosulfuron-methyl expressed as meso-sulfuron	Triticonazole	1-methylcyclopropene
Figs							
Kiwi							
Kumquats							
Litchis							
Mangoes							
Olives (table consumption)							
Olives (oil extraction)							
Papaya							
Passion fruit							
Pineapples							
Pomegranate							
Others		▶ M70 0,02 (*) (P) ◀					
2. Vegetables, fresh or uncooked, frozen or dry					0,01 (*) (P)	0,01 (*) (P)	0,01 (*) (P)
(i) ROOT AND TUBER VEGETABLES	0,02 (*) (P)		0,05 (*) (P)	0,05 (*) (P)			
Beetroot							
Carrots							
Cassava							
Celeriac							
Horseradish							
Jerusalem artichokes							
Parsnips							

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as mesosulfuron	Triticonazole	1-methylcyclopropene
Parsley root							
Radishes		► M70 0,2 (P) ◀					
Salsify							
Sweet potatoes							
Swedes							
Turnips							
Yam							
Others		► M70 0,02 (*) (P) ◀					
(ii) BULB VEGETABLES	0,02 (*) (P)	► M70 0,02 (*) (P) ◀	0,05 (*) (P)				
Garlic				0,5 (P)			
Onions				0,5 (P)			
Shallots				0,5 (P)			
Spring onions							
Others				0,05 (*) (P)			
(iii) FRUITING VEGETABLES							
(a) Solanacea			0,05 (*) (P)				
Tomatoes	0,1 (P)	► M70 0,5 (P) ◀		3 (P)			
Peppers		► M70 0,3 (P) ◀		2 (P)			

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as mesosulfuron	Triticonazole	1-methylcyclopropene
Aubergines	0,1 (P)	► M70 0,5 (P) ◄		3 (P)			
Okra							
Others	0,02 (*) (P)	► M70 0,02 (*) (P) ◄		0,05 (*) (P)			
(b) Cucurbits — edible peel	0,02 (*) (P)	► M70 0,2 (P) ◄	0,05 (*) (P)	2 (P)			
Cucumbers							
Gherkins							
Courgettes							
Others							
(c) Cucurbits — inedible peel	0,05 (P)	► M70 0,1 (P) ◄	0,05 (*) (P)	0,3 (P)			
Melons							
Squashes							
Watermelons							
Others							
(d) Sweet corn	0,02 (*) (P)	► M70 0,02 (*) (P) ◄	0,05 (*) (P)	0,05 (*) (P)			
(iv) BRASSICA VEGETABLES	0,02 (*) (P)		0,05 (*) (P)				
(a) Flowering brassica		► M70 0,3 (P) ◄					
Broccoli (including Calabrese)				1 (P)			
Cauliflower							
Others				0,05 (*) (P)			

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolylfluanid and dimethylaminosulfotoluidide expressed as tolylfluanid)	Mesosulfuron-methyl expressed as mesosulfuron	Triticonazole	1-methylcyclopropene
(b) Head brassica				0,05 (*) (P)			
Brussels sprouts							
Head cabbage		► M70 3 (P) ◀					
Others		► M70 0,02 (*) (P) ◀					
(c) Leafy brassica		► M70 0,2 (P) ◀		0,05 (*) (P)			
Chinese cabbage							
Kale		► M70 0,2 (P) ◀					
Others		► M70 0,02 (*) (P) ◀					
(d) Kohlrabi		► M70 0,02 (*) (P) ◀		0,05 (*) (P)			
(v) LEAF VEGETABLES AND FRESH HERBS	0,02 (*) (P)		0,05 (*) (P)				
(a) Lettuce and similar				20 (P)			
Cress							
Lamb's lettuce		► M70 1 (P) ◀					
Lettuce		► M70 2 (P) ◀					
Scarole (broad-leaf endive)		► M70 2 (P) ◀					
Ruccola							
Leaves and stems of brassica, including turnip greens		► M70 1 (P) ◀					
Others		► M70 0,02 (*) (P) ◀					

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as meso-sulfuron	Triticonazole	1-methylcyclopropene
(b) Spinach and similar		► M70 0,02 (*) (P) ◀		0,05 (*) (P)			
Spinach		► M70 2 (P) ◀					
Beet leaves (chard)							
Others							
(c) Water cress		► M70 0,02 (*) (P) ◀		0,05 (*) (P)			
(d) Witloof		► M70 0,02 (*) (P) ◀		0,05 (*) (P)			
(e) Herbs		► M70 2 (P) ◀		0,05 (*) (P)			
Chervil							
Chives							
Parsley							
Celery leaves							
Others							
(vi) LEGUME VEGETABLES (fresh)	0,02 (*) (P)	► M70 0,02 (*) (P) ◀					
Beans (with pods)				3 (P)			
Beans (without pods)			0,1 (P)				
Peas (with pods)			0,1 (P)	3 (P)			
Peas (without pods)			0,1 (P)				
Others			0,05 (*) (P)	0,05 (*) (P)			

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as meso-sulfuron	Triticonazole	1-methylcyclopropene
(vii) STEM VEGETABLES (fresh)	0,02 (*) (P)		0,05 (*) (P)				
Asparagus							
Cardoons							
Celery		► M70 2 (P) ◀					
Fennel							
Globe artichokes		► M70 0,1 (P) ◀					
Leek				3 (P)			
Rhubarb							
Others		► M70 0,02 (*) (P) ◀		0,05 (*) (P)			
(viii) FUNGI	0,02 (*) (P)	► M70 0,02 (*) (P) ◀	0,05 (*) (P)	0,05 (*) (P)			
(a) Cultivated mushrooms							
(b) Wild mushrooms							
3. Pulses	0,02 (*) (P)	► M70 0,02 (*) (P) ◀		0,05 (*) (P)	0,01 (*)	0,01 (*) (P)	► C13 0,01 (*) (P) ◀
Beans			0,1 (P)				
Lentils							
Peas			0,1 (P)				
Lupines							
Others			0,05 (*) (P)				

▼ **M62**

Groups and examples of individual products to which the MRLs apply	Pesticide residue and maximum residue level (mg/kg)						
	Etoxazole	Indoxacarb as sum of the isomers S and R	MCPA, MCPB including their salts, esters and conjugates expressed as MCPA	Tolyfluanid (sum of tolyfluanid and dimethylaminosulfotoluidide expressed as tolyfluanid)	Mesosulfuron-methyl expressed as meso-sulfuron	Triticonazole	1-methylcyclopropene
4. Oilseeds	0,05 (*) (P)		0,1 (*) (P)	0,1 (*) (P)	0,02 (*) (P)	0,02 (*) (P)	0,02 (*) (P)
Linseed							
Peanuts							
Poppy seed							
Sesame seed							
Sunflower seed							
Rape seed							
Soya bean		► M70 0,5 (P) ◀					
Mustard seed							
Cotton seed							
Hemp seed							
Others		► M70 0,05 (*) (P) ◀					
5. Potatoes	0,02 (*) (P)	► M70 0,02 (*) (P) ◀	0,05 (*) (P)	0,05 (*) (P)	0,01 (*) (P)	0,01 (*) (P)	0,01 (*) (P)
Early potatoes							
Ware potatoes							
6. Tea (dried leaves and stalks, fermented or other-wise, <i>Camellia sinensis</i>)	0,05 (*) (P)	► M70 0,05 (*) (P) ◀	0,1 (*) (P)	0,1 (*) (P)	0,02 (*) (P)	0,02 (*) (P)	0,02 (*) (P)
7. Hops (dried), including hop pellets and unconcentrated powder	0,05 (*) (P)	► M70 0,05 (*) (P) ◀	0,1 (*) (P)	50 (P)	0,02 (*) (P)	0,02 (*) (P)	0,02 (*) (P)

(*) Indicates lower limit of analytical determination.

(P) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 5 June 2011.

▼ **M65**

Groups and examples of individual products to which the MRLs apply	Azinphos-methyl
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar, nuts	
(i) CITRUS FRUIT	0,05 (*)
Grapefruit	
Lemons	
Limes	
Mandarins (including clementines and other hybrids)	
Oranges	
Pomelos	
Others	
(ii) TREE NUTS (shelled or unshelled)	0,5
Almonds	
Brazil nuts	
Cashew nuts	
Chestnuts	
Coconuts	
Hazelnuts	
Macadamia	
Pecans	
Pine nuts	
Pistachios	
Walnuts	
Others	
(iii) POME FRUIT	0,5 (†)
Apples	
Pears	
Quinces	
Others	
(iv) STONE FRUIT	0,5 (†)
Apricots	
Cherries	
Peaches (including nectarines and similar hybrids)	
Plums	
Others	
(v) BERRIES AND SMALL FRUIT	
(a) Table and wine grapes	0,05 (*)
Table grapes	
Wine grapes	
(b) Strawberries (other than wild)	0,5 (†)
(c) Cane fruit (other than wild)	0,5 (†)
Blackberries	
Dewberries	
Loganberries	
Raspberries	
Others	

▼ **M65**

Groups and examples of individual products to which the MRLs apply	Azinphos-methyl
(d) Other small fruit and berries (other than wild)	
Bilberries	
Cranberries	0,1
Currants (red, black and white)	0,5 (!)
Gooseberries	0,5 (!)
Others	0,05 (*)
(e) Wild berries and wild fruit	0,05 (*)
(vi) MISCELLANEOUS	0,05 (*)
Avocados	
Bananas	
Dates	
Figs	
Kiwi	
Kumquats	
Litchis	
Mangoes	
Olives (table consumption)	
Olives (oil extraction)	
Papaya	
Passion fruit	
Pineapples	
Pomegranate	
Others	
2. Vegetables, fresh or uncooked, frozen or dry	
(i) ROOT AND TUBER VEGETABLES	0,05 (*)
Beetroot	
Carrots	
Cassava	
Celeriac	
Horseradish	
Jerusalem artichokes	
Parsnips	
Parsley root	
Radishes	
Salsify	
Sweet potatoes	
Swedes	
Turnips	
Yam	
Others	
(ii) BULB VEGETABLES	0,05 (*)
Garlic	
Onions	
Shallots	
Spring onions	
Others	

▼ **M65**

Groups and examples of individual products to which the MRLs apply	Azinphos-methyl
(iii) FRUITING VEGETABLES	
(a) Solanacea Tomatoes Peppers Aubergines Okra Others	0,05 (*)
(b) Cucurbits — edible peel Cucumbers Gherkins Courgettes Others	0,2 0,05 (*)
(c) Cucurbits — inedible peel Melons Squashes Watermelons Others	0,05 (*)
(d) Sweetcorn	0,05 (*)
(iv) BRASSICA VEGETABLES	0,05 (*)
(a) Flowering brassica Broccoli (including Calabrese) Cauliflower Others	
(b) Head brassica Brussels sprouts Head cabbage Others	
(c) Leafy brassica Chinese cabbage Kale Others	
(d) Kohlrabi	
(v) LEAF VEGETABLES AND FRESH HERBS	0,05 (*)
(a) Lettuce and similar Cress Lamb's lettuce Lettuce Scarole (broad-leaf endive) Rocket Leaves and stems of brassica, including turnip greens Others	
(b) Spinach and similar Spinach Beet leaves (chard) Others	

▼ **M65**

Groups and examples of individual products to which the MRLs apply	Azinphos-methyl
(c) Watercress	
(d) Witloof	
(e) Herbs	
Chervil	
Chives	
Parsley	
Celery leaves	
Others	
(vi) LEGUME VEGETABLES (fresh)	0,05 (*)
Beans (with pods)	
Beans (without pods)	
Peas (with pods)	
Peas (without pods)	
Others	
(vii) STEM VEGETABLES (fresh)	0,05 (*)
Asparagus	
Cardoons	
Celery	
Fennel	
Globe artichokes	
Leek	
Rhubarb	
Others	
(viii) FUNGI	0,05 (*)
(a) Cultivated mushrooms	
(b) Wild mushrooms	
3. Pulses	0,05 (*)
Beans	
Lentils	
Peas	
Lupines	
Others	
4. Oilseed	
Linseed	
Peanuts	
Poppy seeds	
Sesame seeds	
Sunflower seed	
Rapeseed	
Soya bean	
Mustard seed	
Cotton seed	0,2
Hemp seed	
Others	0,05 (*)

▼ **M65**

Groups and examples of individual products to which the MRLs apply	Azinphos-methyl
5. Potatoes Early potatoes Ware potatoes	0,05 (*)
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,1 (*)
7. Hops (dried), including hop pellets and unconcentrated powder	0,1 (*)

(*) Indicates lower limit of analytical determination.

(†) Temporary MRL until 18 September 2008. After this date the MRL will be 0,05 (*) mg/kg unless modified by a Directive or a Regulation.

▼ **M68**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Bifenazate	Pethoxamid	Pyrimethanil	Rimsulfuron
1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts		0,01 (*) (P)		0,05 (*) (P)
(i) CITRUS FRUIT	0,01 (*) (P)		► M70 10 (P) ◀	
Grapefruit				
Lemons				
Limes				
Mandarins (including clementines and other hybrids)				
Oranges				
Pomelos				
Others				
(ii) TREE NUTS (shelled or unshelled)	0,01 (*) (P)		► M70 0,2 (P) ◀	
Almonds				
Brazil nuts				
Cashew nuts				
Chestnuts				
Coconuts				
Hazelnuts				
Macadamia				
Pecans				
Pine nuts				
Pistachios			► M70 0,2 (P) ◀	
Walnuts				
Others			► M70 0,05 (*) (P) ◀	
(iii) POME FRUIT	0,01 (*) (P)		► M70 5 (P) ◀	
Apples				
Pears				
Quinces				
Others				
(iv) STONE FRUIT	0,01 (*) (P)		► M70 3 (P) ◀	
Apricots				
Cherries				
Peaches (including nectarines and similar hybrids)			► M70 10 (P) ◀	
Plums			► M70 3 (P) ◀	

▼ **M68**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Bifenazate	Pethoxamid	Pyrimethanil	Rimsulfuron
Others			► M70 0,05 (*) (P) ◀	
(v) BERRIES AND SMALL FRUIT				
(a) Table and wine grapes	0,01 (*) (P)		► M70 5 (P) ◀	
Table grapes				
Wine grapes				
(b) Strawberries (other than wild)	2 (P)		► M70 5 (P) ◀	
(c) Cane fruit (other than wild)	0,01 (*) (P)			
Blackberries			► M70 10 (P) ◀	
Dewberries				
Loganberries				
Raspberries			► M70 10 (P) ◀	
Others			► M70 0,05 (*) (P) ◀	
(d) Other small fruit & berries (other than wild)	0,01 (*) (P)		► M70 5 (P) ◀	
Bilberries				
Cranberries				
Currants (red, black and white)				
Gooseberries				
Others				
(e) Wild berries and wild fruit	0,01 (*) (P)		► M70 0,05 (*) (P) ◀	
(vi) MISCELLANEOUS	0,01 (*) (P)			
Avocados				
Bananas			► M70 0,1 (P) ◀	
Dates				
Figs				
Kiwi				
Kumquats				
Litchis				
Mangoes				
Olives (table consumption)				
Olives (oil extraction)				
Papaya				
Passion fruit				
Pineapples				

▼ **M68**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Bifenazate	Pethoxamid	Pyrimethanil	Rimsulfuron
Pomegranate				
Others			► M70 0,05 (*) (P) ◀	
2. Vegetables, fresh or uncooked, frozen or dry		0,01 (*) (P)		0,05 (*) (P)
(i) ROOT AND TUBER VEGETABLES	0,01 (*) (P)			
Beetroot				
Carrots			► M70 1 (P) ◀	
Cassava				
Celeriac				
Horseradish				
Jerusalem artichokes				
Parsnips				
Parsley root				
Radishes				
Salsify				
Sweet potatoes				
Swedes				
Turnips				
Yam				
Others			► M70 0,05 (*) (P) ◀	
(ii) BULB VEGETABLES	0,01 (*) (P)			
Garlic				
Onions			► M70 0,1 (P) ◀	
Shallots				
Spring onions				
Others			► M70 0,05 (*) (P) ◀	
(iii) FRUITING VEGETABLES				
(a) Solanacea				
Tomatoes	0,5 (P)		► M70 1 (P) ◀	
Peppers	2 (P)		► M70 2 (P) ◀	
Aubergines	0,5 (P)		► M70 1 (P) ◀	
Okra				
Others	0,01 (*) (P)		► M70 0,05 (*) (P) ◀	

▼ **M68**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Bifenazate	Pethoxamid	Pyrimethanil	Rimsulfuron
(b) Cucurbits — edible peel	0,3 (P)		► M70 1 (P) ◀	
Cucumbers				
Gherkins				
Courgettes				
Others				
(c) Cucurbits — inedible peel	0,01 (*) (P)		► M70 0,05 (*) (P) ◀	
Melons				
Squashes				
Watermelons				
Others				
(d) Sweetcorn	0,01 (*) (P)		► M70 0,05 (*) (P) ◀	
(iv) BRASSICA VEGETABLES	0,01 (*) (P)		► M70 0,05 (*) (P) ◀	
(a) Flowering brassica				
Broccoli (including Calabrese)				
Cauliflower				
Others				
(b) Head brassica				
Brussels sprouts				
Head cabbage				
Others				
(c) Leafy brassica				
Chinese cabbage				
Kale				
Others				
(d) Kohlrabi				
(v) LEAF VEGETABLES AND FRESH HERBS	0,01 (*) (P)			
(a) Lettuce and similar				
Cress				
Lamb's lettuce				
Lettuce			► M70 10 (P) ◀	
Scarole (broad-leaf endive)			► M70 10 (P) ◀	
Rocket				
Leaves and stems of brassica, including turnip greens				
Others			► M70 0,05 (*) (P) ◀	

▼ **M68**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Bifenazate	Pethoxamid	Pyrimethanil	Rimsulfuron
(b) Spinach and similar				
Spinach				
Beet leaves (chard)				
Others				
(c) Watercress			► M70 0,05 (*) (P) ◀	
(d) Witloof			► M70 0,05 (*) (P) ◀	
(e) Herbs			► M70 3 (P) ◀	
Chervil				
Chives				
Parsley				
Celery leaves				
Others				
(vi) LEGUME VEGETABLES (fresh)	0,01 (*) (P)			
Beans (with pods)			► M70 2 (P) ◀	
Beans (without pods)				
Peas (with pods)				
Peas (without pods)			► M70 0,2 (P) ◀	
Others			► M70 0,05 (*) (P) ◀	
(vii) STEM VEGETABLES (fresh)	0,01 (*) (P)			
Asparagus				
Cardoons				
Celery				
Fennel				
Globe artichokes				
Leeks			► M70 1 (P) ◀	
Rhubarb				
Others			► M70 0,05 (*) (P) ◀	
(viii) FUNGI	0,01 (*) (P)		► M70 0,05 (*) (P) ◀	
(a) Cultivated mushrooms				
(b) Wild mushrooms				
3. Pulses	0,01 (*) (P)	0,01 (*) (P)	► M70 0,5 (P) ◀	0,05 (*) (P)
Beans				
Lentils				
Peas				

▼ **M68**

Groups and examples of individual products to which the MRLs apply	Pesticide residues and maximum residue levels (mg/kg)			
	Bifenazate	Pethoxamid	Pyrimethanil	Rimsulfuron
Lupines				
Others				
4. Oilseeds	0,02 (*) (P)	0,01 (*) (P)	► M70 0,1 (*) (P) ◀	0,05 (*) (P)
Linseed				
Peanuts				
Poppy seed				
Sesame seed				
Sunflower seed				
Rape seed				
Soya bean				
Mustard seed				
Cotton seed				
Hemp seed				
Pumpkin seed				
Others				
5. Potatoes	0,01 (*) (P)	0,01 (*) (P)	► M70 0,05 (*) (P) ◀	0,05 (*) (P)
Early potatoes				
Ware potatoes				
6. Tea (dried leaves and stalks, fermented or otherwise, <i>Camellia sinensis</i>)	0,02 (*) (P)	0,02 (*) (P)	► M70 0,1 (*) (P) ◀	0,1 (P)
7. Hops (dried), including hop pellets and unconcentrated powder	0,02 (*) (P)	0,02 (*) (P)	► M70 0,1 (*) (P) ◀	0,1 (*) (P)

(*) Indicates lower limit of analytical determination.

(P) Indicates provisional maximum residue level in accordance with Article 4(1)(f) of Directive 91/414/EEC: unless amended, this level will become definitive with effect from 25 October 2011.

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Fenitrothion
<p>1. Fruit, fresh, dried or uncooked, preserved by freezing, not containing added sugar; nuts</p> <p>(i) CITRUS FRUIT</p> <p>Grapefruit</p> <p>Lemons</p> <p>Limes</p> <p>Mandarins (including clementines and other hybrids)</p> <p>Oranges</p> <p>Pomelos</p> <p>Others</p> <p>(ii) TREE NUTS (shelled or unshelled)</p> <p>Almonds</p> <p>Brazil nuts</p> <p>Cashew nuts</p> <p>Chestnuts</p> <p>Coconuts</p> <p>Hazelnuts</p> <p>Macadamia</p> <p>Pecans</p> <p>Pine nuts</p> <p>Pistachios</p> <p>Walnuts</p> <p>Others</p> <p>(iii) POME FRUIT</p> <p>Apples</p> <p>Pears</p> <p>Quinces</p> <p>Others</p> <p>(iv) STONE FRUIT</p> <p>Apricots</p> <p>Cherries</p> <p>Peaches (including nectarines and similar hybrids)</p> <p>Plums</p> <p>Others</p> <p>(v) BERRIES AND SMALL FRUIT</p> <p>(a) Table and wine grapes</p> <p>Table grapes</p> <p>Wine grapes</p> <p>(b) Strawberries (other than wild)</p> <p>(c) Cane fruit (other than wild)</p> <p>Blackberries</p> <p>Dewberries</p>	0,01 (*)

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Fenitrothion
Loganberries Raspberries Others (d) Other small fruit and berries (other than wild) Bilberries Cranberries Currants (red, black and white) Gooseberries Others (e) Wild berries and wild fruit	
(vi) MISCELLANEOUS Avocados Bananas Dates Figs Kiwi Kumquats Litchis Mangoes Olives (table consumption) Olives (oil extraction) Papaya Passion fruit Pineapples Pomegranate Others	
2. Vegetables, fresh or uncooked, frozen or dry	0,01 (*)
(i) ROOT AND TUBER VEGETABLES Beetroot Carrots Cassava Celeriac Horseradish Jerusalem artichokes Parsnips Parsley root Radishes Salsify Sweet potatoes Swedes Turnips Yam Others	

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Fenitrothion
<p>(ii) BULB VEGETABLES</p> <ul style="list-style-type: none"> Garlic Onions Shallots Spring onions Others <p>(iii) FRUITING VEGETABLES</p> <ul style="list-style-type: none"> (a) Solanacea <ul style="list-style-type: none"> Tomatoes Peppers Aubergines Okra Others (b) Cucurbits — edible peel <ul style="list-style-type: none"> Cucumbers Gherkins Courgettes Others (c) Cucurbits — inedible peel <ul style="list-style-type: none"> Melons Squashes Watermelons Others (d) Sweet corn <p>(iv) BRASSICA VEGETABLES</p> <ul style="list-style-type: none"> (a) Flowering brassica <ul style="list-style-type: none"> Broccoli Cauliflower Others (b) Head brassica <ul style="list-style-type: none"> Brussels sprouts Head cabbage Others (c) Leafy brassica <ul style="list-style-type: none"> Chinese cabbage Kale Others (d) Kohlrabi <p>(v) LEAF VEGETABLES AND FRESH HERBS</p> <ul style="list-style-type: none"> (a) Lettuce & similar <ul style="list-style-type: none"> Cress Lamb's lettuce Lettuce 	

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Fenitrothion
Scarole Ruccola Leaves and stems of brassica Others (b) Spinach & similar Spinach Beet leaves (chard) Others (c) Water cress (d) Witloof (e) Herbs Chervil Chives Parsley Celery leaves Others	
(vi) LEGUME VEGETABLES (fresh) Beans (with pods) Beans (without pods) Peas (with pods) Peas (without pods) Others	
(vii) STEM VEGETABLES (fresh) Asparagus Cardoons Celery Fennel Globe artichokes Leek Rhubarb Others	
(viii) FUNGI (a) Cultivated mushrooms (b) Wild mushrooms	
3. Pulses Beans Lentils Peas Others	0,01 (*)
4. Oil seed Linseed Peanuts Poppy seeds	0,01 (*)

▼ **M52**

Groups and examples of individual products to which the MRLs apply	Fenitrothion
Sesame seeds	
Sunflower seed	
Rape seed	
Soya bean	
Mustard seed	
Cotton seed	
Hemp seed	
Others	
5. Potatoes	0,01 (*)
Early potatoes	
Ware potatoes	
6. Tea (leaves and stems, dried, fermented or otherwise, from the leaves of <i>Camellia sinensis</i>)	0,5
7. Hops (dried), including hop pellets and unconcentrated powder	0,02 (*)

(*) Indicates lower limit of analytical determination.