

ANNEX V

QUALITY AND SAFETY REQUIREMENTS FOR BLOOD AND BLOOD COMPONENTS (as referred to in Article 6)

2. QUALITY CONTROL REQUIREMENTS FOR BLOOD AND BLOOD COMPONENTS
 - 2.1. Blood and blood components must comply with the following technical quality measurements and meet the acceptable results.
 - 2.2. Appropriate bacteriological control of the collection and manufacturing process must be performed.
 - 2.3. Member States must take all necessary measures to ensure that all imports of blood and blood components from third countries, including those used as starting material/raw material for the manufacture of medicinal products derived from human blood or human plasma, shall meet equivalent standards of quality and safety to the ones laid down in this Directive.
 - 2.4. For autologous donations, the measures marked with an asterisk (*) are recommendations only.

Component	Quality measurements required <i>The required frequency of sampling for all measurements shall be determined using statistical process control</i>	Acceptable results for quality measurements
Red cells	Volume	Valid for storage characteristics to maintain product within specifications for haemoglobin and haemolysis
	Haemoglobin (*)	Not less than 45 g per unit
	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Red cells, buffy coat removed	Volume	Valid for storage characteristics to maintain product within specifications for haemoglobin and haemolysis
	Haemoglobin (*)	Not less than 43 g per unit
	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Red cells, leucocyte-depleted	Volume	Valid for storage characteristics to maintain product within specifications

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

		for haemoglobin and haemolysis
	Haemoglobin (*)	Not less than 40 g per unit
	Leucocyte content	Less than 1×10^6 per unit
	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Red cells, in additive solution	Volume	Valid for storage characteristics to maintain product within specifications for haemoglobin and haemolysis
	Haemoglobin (*)	Not less than 45 g per unit
	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Red cells, buffy coat removed, in additive solution	Volume	Valid for storage characteristics to maintain product within specifications for haemoglobin and haemolysis
	Haemoglobin (*)	Not less than 43 g per unit
	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Red cells, leucocyte-depleted, in additive solution	Volume	Valid for storage characteristics to maintain product within specifications for haemoglobin and haemolysis
	Haemoglobin (*)	Not less than 40 g per unit
	Leucocyte content	Less than 1×10^6 per unit
	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Red cells, apheresis	Volume	Valid for storage characteristics to maintain product within specifications for haemoglobin and haemolysis
	Haemoglobin (*)	Not less than 40 g per unit

	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Whole blood	Volume	Valid for storage characteristics to maintain product within specifications for haemoglobin and haemolysis 450 ml +/- 50ml For paediatric autologous whole blood collections — not to exceed 10,5 ml per kg body weight
	Haemoglobin (*)	Not less than 45 g per unit
	Haemolysis	Less than 0,8 % of red cell mass at the end of the shelf life
Platelets, apheresis	Volume	Valid for storage characteristics to maintain product within specifications for pH
	Platelet content	Variations in platelet content per single donation are permitted within limits that comply with validated preparation and preservation conditions
	pH	[^{F1} Minimum 6,4 corrected for 22 °C, at the end of the shelf life]
Platelets, apheresis, leucocyte-depleted	Volume	Valid for storage characteristics to maintain product within specifications for pH
	Platelet content	Variations in platelet content per single donation are permitted within limits that comply with validated preparation and preservation conditions
	Leucocyte content	Less than 1×10^6 per unit
	pH	[^{F1} Minimum 6,4 corrected for 22 °C, at the end of the shelf life]
Platelets, recovered, pooled	Volume	Valid for storage characteristics to maintain

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

		product within specifications for pH
	Platelet content	Variations in platelet content per pool are permitted within limits that comply with validated preparation and preservation conditions
	Leucocyte content	Less than $0,2 \times 10^9$ per single unit (platelet-rich plasma method) Less than $0,05 \times 10^9$ per single unit (buffy coat method)
	pH	[^{F1} Minimum 6,4 corrected for 22 °C, at the end of the shelf life]
Platelets, recovered, pooled, leucocyte-depleted	Volume	Valid for storage characteristics to maintain product within specifications for pH
	Platelet content	Variations in platelet content per pool are permitted within limits that comply with validated preparation and preservation conditions
	Leucocyte content	Less than 1×10^6 per pool
	pH	[^{F1} Minimum 6,4 corrected for 22 °C, at the end of the shelf life]
Platelets, recovered, single unit	Volume	Valid for storage characteristics to maintain product within specifications for pH
	Platelet content	Variations in platelet content per single unit are permitted within limits that comply with validated preparation and preservation conditions
	Leucocyte content	Less than $0,2 \times 10^9$ per single unit (platelet-rich plasma method) Less than $0,05 \times 10^9$ per single unit (buffy coat method)

	pH	[^{F1} Minimum 6,4 corrected for 22 °C, at the end of the shelf life]
Platelets, recovered, single unit, leucocyte-depleted	Volume	Valid for storage characteristics to maintain product within specifications for pH
	Platelet content	Variations in platelet content per single unit are permitted within limits that comply with validated preparation and preservation conditions
	Leukocyte content	Less than 1×10^6 per unit
	pH	[^{F1} Minimum 6,4 corrected for 22 °C, at the end of the shelf life]
Plasma, fresh-frozen	Volume	Stated volume +/- 10 %
	Factor VIIIc (*)	Average (after freezing and thawing): 70 % or more of the value of the freshly collected plasma unit
	Total protein (*)	Not less than 50 g/l
	Residual cellular content (*)	Red cells: less than $6,0 \times 10^9/l$ Leucocytes: less than $0,1 \times 10^9/l$ Platelets: less than $50 \times 10^9/l$
Plasma, fresh-frozen, cryoprecipitate-depleted	Volume	Stated volume: +/- 10 %
	Residual cellular content (*)	Red cells: less than $6,0 \times 10^9/l$ Leucocytes: less than $0,1 \times 10^9/l$ Platelets: less than $50 \times 10^9/l$
Cryoprecipitate	Fibrinogen content (*)	Greater than or equal to 140 mg per unit
	Factor VIIIc content (*)	Greater than or equal to 70 international units per unit
Granulocytes, apheresis	Volume	Less than 500 ml
	Granulocyte content	Greater than 1×10^{10} granulocytes per unit

Status: EU Directives are being published on this site to aid cross referencing from UK legislation. After
IP completion day (31 December 2020 11pm) no further amendments will be applied to this version.

Textual Amendments

- F1** Substituted by [Commission Implementing Directive 2011/38/EU](#) of 11 April 2011 amending Annex V to Directive 2004/33/EC with regards to maximum pH values for platelets concentrates at the end of the shelf life (Text with EEA relevance).