[^{F1}ANNEX I

Microbiological criteria for foodstuffs

Textual Amendments

F1 Substituted by Commission Regulation (EC) No 1441/2007 of 5 December 2007 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs (Text with EEA relevance).

Chapter 1.

Food safety criteria

Food	Micro-	Sampling plan ^a		Limits ^b		Analytica	
category	organism their toxins, metabolit	^{s/} n	c	Vm	Μ	reference method ^c	the criterion applies
	<i>Listeria</i> Read/ <i>iccytog</i> eat foods intended for infants and ready- to- eat foods for special medical purposes ^d	10 enes	0	Absence	in 25 g	EN/ISO 11290-1	Products placed on the market during their shelf-life
1.2 F t f a t	<i>Listeria</i> Read <i>inohocytog</i> to- eat foods able to support	5 enes	0	100 cfu/g	g°	EN/ISO 11290-2 ^r	Products placed on the market during their shelf-life
	the growth of <i>L. monocytogen</i> other than	5 es,	0	Absence	in 25 g ^g	EN/ISO 11290-1	Before the food has left the immediate control of the food business

	those intended for infants and for special medical purposes					operator, who has produced it
1.3	Listeria Ready of to- eat foods unable to support the growth of L monocytogen other than those intended for infants and for special medical purposes ^{dh}		0	100 cfu/g	EN/ISO 11290-2 ^f	Products placed on the market during their shelf-life
1.4	Salmonella Minced meat and meat preparations intended to be eaten raw	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
[^{F3} 1.5	Salmonella Minced meat and meat preparations made from	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life]

	poultry meat intended to be eaten cooked					
1.6	Salmonella Minced meat and meat preparations made from other species than poultry intended to be eaten cooked	5	0	Absence in 10 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.7	Salmonella Mechanically separated meat (MSM) ⁱ	5	0	Absence in 10 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.8	Salmonella Meat products intended to be eaten raw, excluding products where the manufacturin process or the composition of the product will		0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life

	eliminate the salmonella risk					
[^{F3} 1.9	Salmonella Meat products made from poultry meat intended to be eaten cooked	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life]
1.10	Salmonella Gelatine and collagen	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.11	Salmonella Cheeses, butter and cream made from raw milk or milk that has undergone a lower heat treatment than pasteurisation	l	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.12	Salmonella Milk powder and whey powder		0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life

1.13	Salmonella Ice cream ^k , excluding products where the manufacturin process or the composition of the product will eliminate the salmonella risk	g	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.14	Salmonella Egg products, excluding products where the manufacturin process or the composition of the product will eliminate the salmonella risk		0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.15	Salmonella Ready- to- eat foods containing raw egg, excluding products where the	5	0	Absence in 25 g or ml	EN/ISO 6579	Products placed on the market during their shelf-life

	manufacturin process or the composition of the product will eliminate the salmonella risk					
1.16	Salmonella Cooked crustaceans and molluscan shellfish	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.17	Salmonella Live bivalve molluscs and live echinoderms, tunicates and gastropods		0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.18	Salmonella Sprouted seeds (ready- to- eat) ^w	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.19	Salmonella Precut fruit and vegetables (ready- to- eat)	5	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.20	Salmonella Unpasteurised fruit and vegetable juices	5 d	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during

	(ready- to- eat)					their shelf-life
1.21	Staphyloco Chemerotoxin milk powder and whey powder, as referred to in the coagulase- positive staphylococc criteria in Chapter 2 2 of this Annex	15	0	Not detected in 25 g	European screening method of the CRL for coagulase positive staphyloco	Products placed on the market during their shelf-life cci ^m
1.22	Salmonella Dried infant formulae and dried dietary foods for special medical purposes intended for infants below six months of age	30	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life
1.23	Salmonella Dried follow- on formulae	30	0	Absence in 25 g	EN/ISO 6579	Products placed on the market during their shelf-life

Status: Point in time view as at 01/06/2014.

Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2073/2005. (See end of Document for details)

[^{F3} 1.24	<i>Cronobacte</i> Dreft in <i>fashterobace</i> formulae <i>kii</i>) and dried dietary foods for special medical purposes intended for infants below 6 months of age"		0	Absence in 10 g		ISO/TS 22964	Products placed on the market during their shelf-life]
1.25	Live coli ^o bivalve molluscs and live echinoderms, tunicates and gastropods	1 ^p	0	230 MPN/100 g of flesh and intra- valvular liquid		ISO TS 16649-3	Products placed on the market during their shelf-life
1.26	Histamine Fishery products from fish species associated with a high amount of histidine ⁴	9 ^r	2	100 mg/ kg	200 mg/ kg	HPLC ^s	Products placed on the market during their shelf-life
[^{F2} 1.27	Histamine Fishery products, except those in food category 1.27a,	9 ^r	2	200 mg/ kg	400 mg/ kg	HPLC	Products placed on the market during their shelf-life]

vhich ave indergone						
nzyme naturation reatment n orine, nanufactured rom ish pecies issociated vith igh ingh						
	1	0	400 mg/kg		HPLC*	Products placed on the market during their shelf- life]]
Salmonella Tesphimuriu Oultry enteritidis neat	5 m ^u Salmonel	0 la	Absence in 25 g		Le Minor scheme (for	Products placed on the market during their shelf-life
Shiga producing <i>E. coli</i> (STEC) 0157, 026, 0111, 0103, 0145 and 0104:H4	5	0	Absence in 25 grams		CEN/ ISO TS 13136 ^v	Products placed on the market during their shelf-life
	shiga Shiga	orine, manufactured rom ish pecies ssociated vith ligh mount of istidine ⁴ Histamine 1 rish auce produced by ermentation of ishery products <i>Salmonella</i> 5 <i>Typhimurium</i> ⁴ <i>Salmonel</i> <i>enferitidis</i> neat <i>Shiga</i> 5 <i>Shiga</i> 5 <i>Shi</i>	rine, manufactured form ish pecies issociated with igh mount of istidine ^q Histamine auce produced by ermentation of ishery products Salmonella 5 0 resphimurium"Salmonella enteritidis neat Shiga 5 0 Shiga 5 0 Shiga 5 0 Shiga 5 0 Shiga 5 0 Shiga 10 Shiga 5 0 Shiga 5 0 Shiga 10 Shiga 10 Shig	rine, manufactured from ish pecies ssociated with ingh mount of istidine ⁴ Histamine 1 0 400 mg/kg auce froduced by ermentation of ishery products Salmonella 5 0 frshimurium ⁴ Salmonella enteritidis neat Shiga 5 0 Koxtiff producing E. coli (STEC) 0157, 026, 0111, 0103, 0145 and 0104:H4	rrine, manufactured form ish pecies ssociated with ingh mount of istidine ⁴ Histamine auce produced yy ermentation of ishery products Salmonella 5 Shiga Shiga Shiga Shiga Shiga Shiga Shiga O Shiga O Shiga Shig	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$

с The most recent edition of the standard shall be used. d Regular testing against the criterion is not required in normal circumstances for the following ready-to-eat foods: those which have received heat treatment or other processing effective to eliminate L. monocytogenes, when recontamination is not possible after this treatment (for example, products heat treated in their final package), fresh, uncut and unprocessed vegetables and fruits, excluding sprouted seeds, bread, biscuits and similar products, bottled or packed waters, soft drinks, beer, cider, wine, spirits and similar products, sugar, honey and confectionery, including cocoa and chocolate products, live bivalve molluscs[^{F3},] [^{F4}food grade salt.] This criterion shall apply if the manufacturer is able to demonstrate, to the satisfaction of the competent authority, that the product will not exceed the limit 100 cfu/g throughout the shelf-life. The operator may fix intermediate limits during the process that must be low enough to guarantee that the limit of 100 cfu/g is not exceeded at the end of shelf-life. f 1 ml of inoculum is plated on a Petri dish of 140 mm diameter or on three Petri dishes of 90 mm diameter. This criterion shall apply to products before they have left the immediate control of the producing food business operator, g when he is not able to demonstrate, to the satisfaction of the competent authority, that the product will not exceed the limit of 100 cfu/g throughout the shelf-life. h Products with pH \leq 4,4 or $a_w \leq$ 0,92, products with pH \leq 5,0 and $a_w \leq$ 0,94, products with a shelf-life of less than five days shall be automatically considered to belong to this category. Other categories of products can also belong to this category, subject to scientific justification. i This criterion shall apply to mechanically separated meat (MSM) produced with the techniques referred to in paragraph 3 of Chapter III of Section V of Annex III to Regulation (EC) No 853/2004 of the European Parliament and of the Council. j Excluding products when the manufacturer can demonstrate to the satisfaction of the competent authorities that, due to the ripening time and a_w of the product where appropriate, there is no salmonella risk. k Only ice creams containing milk ingredients. 1 [^{F5}] m Reference: Community reference laboratory for coagulase positive staphylococci. European screening method for the detection of staphylococcal enterotoxins in milk and milk products. Parallel testing for Enterobacteriaceae and E. sakazakii shall be conducted, unless a correlation between these micron organisms has been established at an individual plant level. If Enterobacteriaceae are detected in any of the product samples tested in such a plant, the batch must be tested for *E. sakazakii*. It shall be the responsibility of the manufacturer to demonstrate to the satisfaction of the competent authority whether such a correlation exists between Enterobacteriaceae and E. sakazakii. E. coli is used here as an indicator of faecal contamination. 0 A pooled sample comprising a minimum of 10 individual animals. р Particularly fish species of the families: Scombridae, Clupeidae, Engraulidae, Coryfenidae, Pomatomidae, q Scombresosidae. I^{F2}Single samples may be taken at retail level. In such a case the presumption laid down in Article 14(6) of Regulation r (EC) No 178/2002, according to which the whole batch should be deemed unsafe, shall not apply, unless the result is above M.] References: 1. Malle P., Valle M., Bouquelet S. Assay of biogenic amines involved in fish decomposition. J. AOAC S Internat. 1996, 79, 43-49. 2. Duflos G., Dervin C., Malle P., Bouquelet S. Relevance of matrix effect in determination of biogenic amines in plaice (Pleuronectes platessa) and whiting (Merlangus merlangus. J. AOAC Internat. 1999, 82, 1097-1101. t I^{F6}This criterion shall apply to fresh meat from breeding flocks of *Gallus gallus*, laying hens, broilers and breeding and fattening flocks of turkeys. As regards monophasic Salmonella typhimurium only [X11,4,[5],12:i:-] is included.] u [^{F7}Taking into account the most recent adaptation by the European Union reference laboratory for Escherichia coli, including Verotoxigenic E. coli (VTEC), for the detection of STEC 0104:H4. Excluding sprouts that have received a treatment effective to eliminate *Salmonella* spp. and STEC.] w

Editorial Information

- X1 Substituted by Corrigendum to Commission Regulation (EU) No 1086/2011 of 27 October 2011 amending Annex II to Regulation (EC) No 2160/2003 of the European Parliament and of the Council and Annex I to Commission Regulation (EC) No 2073/2005 as regards salmonella in fresh poultry meat (Official Journal of the European Union L 281 of 28 October 2011).
- X2 Substituted by Corrigendum to Commission Regulation (EU) No 1019/2013 of 23 October 2013 amending Annex I to Regulation (EC) No 2073/2005 as regards histamines in fishery products (Official Journal of the European Union L 282 of 24 October 2013).

Textual Amendments

- F2 Substituted by Commission Regulation (EU) No 1019/2013 of 23 October 2013 amending Annex I to Regulation (EC) No 2073/2005 as regards histamine in fishery products (Text with EEA relevance).
- **F3** Substituted by Commission Regulation (EU) No 365/2010 of 28 April 2010 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs as regards Enterobacteriaceae in pasteurised milk and other pasteurised liquid dairy products and Listeria monocytogenes in food grade salt (Text with EEA relevance).
- F4 Inserted by Commission Regulation (EU) No 365/2010 of 28 April 2010 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs as regards Enterobacteriaceae in pasteurised milk and other pasteurised liquid dairy products and Listeria monocytogenes in food grade salt (Text with EEA relevance).
- **F5** Deleted by Commission Regulation (EU) No 209/2013 of 11 March 2013 amending Regulation (EC) No 2073/2005 as regards microbiological criteria for sprouts and the sampling rules for poultry carcases and fresh poultry meat (Text with EEA relevance).
- **F6** Inserted by Commission Regulation (EU) No 1086/2011 of 27 October 2011 amending Annex II to Regulation (EC) No 2160/2003 of the European Parliament and of the Council and Annex I to Commission Regulation (EC) No 2073/2005 as regards salmonella in fresh poultry meat (Text with EEA relevance).
- F7 Inserted by Commission Regulation (EU) No 209/2013 of 11 March 2013 amending Regulation (EC) No 2073/2005 as regards microbiological criteria for sprouts and the sampling rules for poultry carcases and fresh poultry meat (Text with EEA relevance).

Interpretation of the test results

The limits given refer to each sample unit tested, excluding live bivalve molluscs and live echinoderms, tunicates and gastropods in relation to testing *E. coli*, where the limit refers to a pooled sample.

The test results demonstrate the microbiological quality of the batch tested⁽¹⁾.

L. monocytogenes in ready-to-eat foods intended for infants and for special medical purposes:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

L. monocytogenes in ready-to-eat foods able to support the growth of *L. monocytogenes* before the food has left the immediate control of the producing food business operator when he is not able to demonstrate that the product will not exceed the limit of 100 cfu/g throughout the shelf-life:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

L. monocytogenes in other ready-to-eat foods and E. coli in live bivalve molluscs:

satisfactory, if all the values observed are \leq the limit,

— unsatisfactory, if any of the values are > the limit.

Salmonella in different food categories:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

Staphylococcal enterotoxins in dairy products:

- satisfactory, if in all the sample units the enterotoxins are not detected,
- unsatisfactory, if the enterotoxins are detected in any of the sample units.

Enterobacter sakazakii in dried infant formulae and dried dietary foods for special medical purposes intended for infants below 6 months of age:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

[^{F2}Histamine in fishery products:

- Histamine in fishery products from fish species associated with a high amount of histidine except fish sauce produced by fermentation of fishery products:
 - satisfactory, if the following requirements are fulfilled:
 - 1. the mean value observed is $\leq m$
 - 2. a maximum of c/n values observed are between m and M
 - 3. no values observed excess the limit of M.
 - unsatisfactory, if the mean value observed exceeds m or more than c/n values are between m and M or one or more of the values observed are > M.

Histamine in fish sauce produced by fermentation of fishery products:

- satisfactory, if the value observed is \leq the limit,
- unsatisfactory, if the value observed is > the limit.]

Chapter 2.

Process hygiene criteria

Food	Micro- organisn	Sampling plan ^a		Limits ^b		Analytic		Action
category		ns _h	c	m	М	cr	the	in case of n unsatisfactory results
	Aerobic Carrases ofount cattle, sheep, goats and horses ^d			3,5 log cfu/cm ² daily mean log	5,0 log cfu/cm ² daily mean log	ISO 4833	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls

2.1 *Meat and products thereof*

	Enterobac	teriaceae		1,5 log cfu/cm ² daily mean log	2,5 log cfu/cm ² daily mean log	ISO 21528-2	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls
2.1.2	Aerobic Carcases of ount pigs ^d			4,0 log cfu/cm ² daily mean log	5,0 log cfu/cm ² daily mean log	ISO 4833	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls
	Enterobac	teriaceae		2,0 log cfu/cm ² daily mean log	3,0 log cfu/cm ² daily mean log	ISO 21528-2	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls
2.1.3	Salmonell Carcases of cattle, sheep, goats and horses	€£0°	2 ^f	Absence in the area tested per carcase		EN/ISO 6579	Carcases after dressing but before chilling	Improvements in slaughter hygiene, review of process controls and of origin of animals
[^{F8} 2.1.4	Salmonell Carcases of pigs	₫0°	3 ^f	Absence : area teste carcase		EN/ISO 6579	Carcases after dressing but before chilling	Improvements in slaughter hygiene and review of process controls, origin of animals and

Status: Point in time view as at 01/06/2014.

Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2073/2005. (See end of Document for details)

[^{F9} 2.1.5	Salmonel. Poultry spo carcases of broilers and turkeys	⁷ 4 50 (⁵)	7 (⁶) From 1.1.2012 c = 5 for broilers From 1.1.2013 c = 5 for turkeys		in 25 g of sample of	EN/ISO 6579 (for detection)	Carcases after chilling	of the biosecurity measures in the farms of origin] Improvement in slaughter hygiene and review of process controls, origin of animals and biosecurity measures in the farms of
2.1.6	Aerobic Milisfiy meat count ^g	5	2	5×10^5 cfu/g	5×10^6 cfu/g	ISO 4833	End of the manufactu process	origin] Improvements in upingluction hygiene and improvements in selection and/or origin of raw materials
	E. coli ^h	5	2	50 cfu/g	500 cfu/ g	ISO 16649-1 or 2	End of the manufactu process	Improvements in apingluction hygiene and improvements in selection and/or origin of raw materials
2.1.7	Aerobic Mechanica separated	5 Illy	2	5×10^5 cfu/g	5×10^{6} cfu/g	ISO 4833	End of the manufactu process	Improvements in upingluction hygiene

	meat (MSM) ⁱ							and improvements in selection and/or origin of raw materials
	E. coli ^h	5	2	50 cfu/g	500 cfu/ g	ISO 16649-1 or 2	End of the manufactu process	Improvements in upingluction hygiene and improvements in selection and/or origin of raw materials
2.1.8	<i>E. coli</i> ^h Meat preparation	5 ns	2	500 cfu/ g or cm ²	5 000 cfu/g or cm ²	ISO 16649-1 or 2	End of the manufactu process	Improvements in upingluction hygiene and improvements in selection and/or origin of raw materials
a n = nur	nber of units co	omprising the s	sample; c = nur	mber of sample	e units giving	values between	m and M.	
b For por	ints 2.1.3-2.1.5	m = M.						
c The me	ost recent editio	n of the standa	ard shall be use	ed.				
	nits (m and M) taking a log va							be calculated
	samples shall b ncies laid down			ve sampling se	ssions in acco	rdance with the	e sampling rule	es and
into ac	mber of sample count the progr ence may use lo	ess made in red	ducing the saln	nonella prevale				
g This cr hours.	iterion shall no	t apply to mind	ced meat produ	iced at retail le	vel when the s	helf-life of the	product is less	then 24
h E. coli	is used here as	an indicator of	f faecal contam	nination.				
	criteria apply to r III of Section							
	re Salmonella s idis in order to							

Textual Amendments

- **F8** Substituted by Commission Regulation (EU) No 217/2014 of 7 March 2014 amending Regulation (EC) No 2073/2005 as regards Salmonella in pig carcases (Text with EEA relevance).
- **F9** Substituted by Commission Regulation (EU) No 1086/2011 of 27 October 2011 amending Annex II to Regulation (EC) No 2160/2003 of the European Parliament and of the Council and Annex I to Commission Regulation (EC) No 2073/2005 as regards salmonella in fresh poultry meat (Text with EEA relevance).

Interpretation of the test results

The limits given refer to each sample unit tested, excluding testing of carcases where the limits refer to pooled samples.

The test results demonstrate the microbiological quality of the process tested.

Enterobacteriaceae and aerobic colony count in carcases of cattle, sheep, goats, horses and pigs:

- satisfactory, if the daily mean log is $\leq m$,
- acceptable, if the daily mean log is between m and M,
- unsatisfactory, if the daily mean $\log i s > M$.

Salmonella in carcases:

- satisfactory, if the presence of *Salmonella* is detected in a maximum of c/n samples,
- unsatisfactory, if the presence of *Salmonella* is detected in more than c/n samples.

After each sampling session, the results of the last ten sampling sessions shall be assessed in order to obtain the n number of samples.

E. coli and aerobic colony count in minced meat, meat preparations and mechanically separated meat (MSM):

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.
- 2.2 Milk and dairy products

Food	Micro-	Samplin	g plan ^a	Limits ^b		Analytic		Action
category	organisr	ns _h	c	m	Μ	referenc method ^c	_	in case of unsatisfactory results
[^{F3} 2.2.1	Entero- Pasteurises milk and other pasteurised liquid dairy products ^d		0	10 cfu/ml		ISO 21528-2	End of the manufacto process	Check on the definition of heat- treatment and prevention of recontamination as well

_								as the quality of raw materials]
2.2.2	<i>E. coli</i> ^e Cheeses made from milk or whey that has undergone heat treatment	5	2	100 cfu/ g	1 000 cfu/g	ISO 16649-1 or 2	At the time during the manufactu process when the <i>E.</i> <i>coli</i> count is expected to be highest ^f	Improvements in production hygiene uning selection of raw materials
2.2.3	Coagulase Possiese Waaphyloc from raw milk	occi	2	10 ⁴ cfu/ g	10 ⁵ cfu/ g	EN/ISO 6888-2	At the time during the manufactu process	selection
2.2.4	Coagulase positive maaphyloc from milk that has undergone a lower heat treatment than pasteurisat and ripened cheeses made from milk or whey that has undergone pasteurisat or a stronger	occi ion ^g	2	100 cfu/ g	1 000 cfu/g	EN/ISO 6888-1 or 2	when the number of staphyloc is expected to be highest	of raw materials. If values > 10 ⁵ cfu/ Seire detected, the cheese batch has to be tested for staphylococcal enterotoxins.

	heat treatment ^g							
2.2.5	Coagulase Unrinened soft softaphyloc cheeses (fresh cheeses) made from milk or whey that has undergone pasteurisat or a stronger heat treatment ^g	occi	2	10 cfu/g	100 cfu/ g	EN/ISO 6888-1 or 2	End of the manufacta process	Improvements in upingluction hygiene. If values $> 10^5$ cfu/g are detected, the cheese batch has to be tested for staphylococcal enterotoxins.
2.2.6	<i>E. coli</i> ^e Butter and cream made from raw milk or milk that has undergone a lower heat treatment than pasteurisat	5 ion	2	10 cfu/g	100 cfu/ g	ISO 16649-1 or 2	End of the manufactu process	Improvements in upingluction hygiene and selection of raw materials
2.2.7	Enterobac Milk powder and whey powder ^d	teriaceae	0	10 cfu/g	·	ISO 21528-2	End of the manufactu process	Check on the deffig eiency of heat treatment and prevention of recontamination

	Coagulase positive staphyloco		2	10 cfu/g	100 cfu/ g	EN/ISO 6888-1 or 2	End of the manufactu process	Improvements in apingluction hygiene. If values $> 10^5$ cfu/g are detected, the batch has to be tested for staphylococcal enterotoxins.
2.2.8	Enterobac Ice cream ^h and frozen dairy desserts		2	10 cfu/g	100 cfu/ g	ISO 21528-2	End of the manufactu process	Improvements in pingluction hygiene
2.2.9	Enterobac Dried infant formulae and dried dietary foods for special medical purposes intended for infants below six months of age		0	Absence in 10 g		ISO 21528-1	End of the manufacta process	Improvements in upingluction hygiene to minimise contamination ⁱ
2.2.10	Enterobac Dried follow- on formulae	teriaceae	0	Absence i	n 10 g	ISO 21528-1	End of the manufactu process	Improvements in apingluction hygiene to minimise contamination

Status: Point in time view as at 01/06/2014.

Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2073/2005. (See end of Document for details)

Presumptive 1 $50 \, \text{cfu/g}$ 500 cfu/ EN/ISO End Improvements 2.2.11 Bucillus of the 7932^j in g ntant manufactupinguction formulae process hygiene. and Prevention dried of dietary recontamination. foods Selection for of raw special material. medical purposes intended for infants below six months bf age n = number of units comprising the sample; c = number of sample units giving values between m and M. a b [F3For points 2.2.1, 2.2.7, 2.2.9 and 2.2.10 m=M.] с The most recent edition of the standard shall be used. d The criterion shall not apply to products intended for further processing in the food industry. e E. coli is used here as an indicator for the level of hygiene. f For cheeses which are not able to support the growth of E. coli, the E. coli count is usually the highest at the beginning of the ripening period, and for cheeses which are able to support the growth of E. coli, it is normally at the end of the ripening period. Excluding cheeses where the manufacturer can demonstrate, to the satisfaction of the competent authorities, that the g product does not pose a risk of staphylococcal enterotoxins. h Only ice creams containing milk ingredients. i Parallel testing for Enterobacteriaceae and E. sakazakii shall be conducted, unless a correlation between these microorganisms has been established at an individual plant level. If Enterobacteriaceae are detected in any of the product samples tested in such a plant, the batch has to be tested for E. sakazakii. It shall be the responsibility of the manufacturer to demonstrate to the satisfaction of the competent authority whether such a correlation exists between Enterobacteriaceae and E_sakazakii 1 ml of inoculum is plated on a Petri dish of 140 mm diameter or on three Petri dishes of 90 mm diameter. i Interpretation of the test results The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

Enterobacteriaceae in dried infant formulae, dried dietary foods for special medical purposes intended for infants below six months of age and dried follow-on formulae:

- satisfactory, if all the values observed indicate the absence of the bacterium,
- unsatisfactory, if the presence of the bacterium is detected in any of the sample units.

E. coli, Enterobacteriaceae (other food categories) and coagulase-positive staphylococci:

- satisfactory, if all the values observed are \leq m,

Status: Point in time view as at 01/06/2014.
Changes to legislation: There are currently no known outstanding effects for the
Commission Regulation (EC) No 2073/2005. (See end of Document for details)
Commission Regulation (EC) No 20/3/2005. (See end of Document for details)

- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Presumptive *Bacillus cereus* in dried infant formulae and dried dietary foods for special medical purposes intended for infants below six months of age:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- -- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Food	Micro- organisn	Sampling plan ^a		Limits		AnalyticaStage		Action
category		ns _h	c	m	М	method ^b the of criterion unsati		
	Enterobac Egg products	teriaceae	2	10 cfu/g or ml	100 cfu/ g or ml	ISO 21528-2	End of the manufacto process	Checks on the ueffigiency of the heat treatment and prevention of recontamination

2.3 *Egg products*

Interpretation of the test results

The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

Enterobacteriaceae in egg products:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.
- 2.4 Fishery products

	³ Food Mic ntegory orga	ro- Sampling plan ^a inisms	Limits			
a	n = number of u	nits comprising the sample; $c = r$	number of sample units giving	values betweer	n m and M.	
b	The most recent	edition of the standard shall be u	ised.]			

Status: Point in time view as at 01/06/2014.

Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2073/2005. (See end of Document for details)

		n	c	m	Μ	Analytic referenc method ^b	e where the	Action in case of unsatisfactory results
2.4.1.	E. coli Shelled and shucked products	5	2	1 MPN/ g	10 MPN/g	ISO TS 16649-3	End of the manufactu process	Improvements in apirogluction hygiene
	ocoagulas cpokiet/e cstapavelaa and molluscan shellfish	occi	2	100 cfu/ g	1 000 cfu/g	EN/ISO 6888-1 or 2	End of the manufactu process	Improvements in apingluction hygiene

a n = number of units comprising the sample; c = number of sample units giving values between m and M.

b The most recent edition of the standard shall be used.]

Editorial Information

X3 Substituted by Corrigendum to Commission Regulation (EC) No 1441/2007 of 5 December 2007 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs (Official Journal of the European Union L 322 of 7 December 2007).

Interpretation of the test results

The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

E. coli in shelled and shucked products of cooked crustaceans and molluscan shellfish:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Coagulase-positive staphylococci in shelled and cooked crustaceans and molluscan shellfish:

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- -- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

	bod	Micro-	Samplin	g plan ^a	Limits		AnalyticaStage reference where method ^b the		Action	
ca	itegory	organisn	nş _h	c	m	М			in case of	
a	a n = number of units comprising the sample; c = number of sample units giving values between m and M.									
b	The mos	st recent editio	on of the standa	ard shall be use	ed.					

2.5 *Vegetables, fruits and products thereof*

							criterion applies	unsatisfactory results
2.5.1	<i>E. coli</i> Precut fruit and vegetables (ready- to- eat)	5	2	100 cfu/ g	1 000 cfu/g	ISO 16649-1 or 2	Manufact process	uningrovements in production hygiene, selection of raw materials
2.5.2	<i>E. coli</i> Unpasteuri fruit and vegetable juices (ready- to- eat)	5 sed	2	100 cfu/ g	1 000 cfu/g	ISO 16649-1 or 2	Manufact process	uhing rovements in production hygiene, selection of raw materials
a n = nu	umber of units co	omprising the s	sample; c = nur	mber of sample	e units giving	values between	m and M.	
b The m	ost recent editio	on of the standa	ard shall be use	ed.				

Interpretation of the test results

The limits given refer to each sample unit tested.

The test results demonstrate the microbiological quality of the process tested.

E. coli in precut fruit and vegetables (ready-to-eat) and in unpasteurised fruit and vegetable juices (ready-to-eat):

- satisfactory, if all the values observed are $\leq m$,
- acceptable, if a maximum of c/n values are between m and M, and the rest of the values observed are \leq m,
- -- unsatisfactory, if one or more of the values observed are > M or more than c/n values are between m and M.

Chapter 3.

Rules for sampling and preparation of test samples

3.1 *General rules for sampling and preparation of test samples*

In the absence of more specific rules on sampling and preparation of test samples, the relevant standards of the ISO (International Organisation for Standardisation) and the guidelines of the Codex Alimentarius shall be used as reference methods.

[^{F9}3.2 Bacteriological sampling in slaughterhouses and at premises producing minced meat, meat preparations, mechanically separated meat and fresh meat Sampling rules for carcases of cattle, pigs, sheep, goats and horses

The destructive and non-destructive sampling methods, the selection of the sampling sites and the rules for storage and transport of samples to be used are set out in standard ISO 17604.

Five carcases shall be sampled at random during each sampling session. Sample sites must be selected taking into account the slaughter technology used in each plant.

When sampling for analyses of Enterobacteriaceae and aerobic colony counts, four sites of each carcase shall be sampled. Four tissue samples representing a total of 20 cm^2 shall be obtained by the destructive method. When using the non-destructive method for this purpose, the sampling area shall cover a minimum of 100 cm^2 (50 cm² for small ruminant carcases) per sampling site.

When sampling for salmonella analyses, an abrasive sponge sampling method shall be used. Areas most likely to be contaminated shall be selected. The total sampling area shall cover a minimum of 400 cm^2 .

When samples are taken from the different sampling sites on the carcase, they shall be pooled before examination.

Sampling rules for poultry carcases and fresh poultry meat

[^{F10}Slaughterhouses shall sample whole poultry carcases with neck skin for *Salmonella* analyses. Cutting and processing establishments other than those adjacent to a slaughterhouse cutting and processing meat received only from this slaughterhouse, shall also take samples for *Salmonella* analysis. When doing so, they shall give priority to whole poultry carcases with neck skin, if available, but ensuring that also poultry portions with skin and/or poultry portions without skin or with only a small amount of skin are covered, and that choice shall be risk-based.]

Textual Amendments

F10 Substituted by Commission Regulation (EU) No 209/2013 of 11 March 2013 amending Regulation (EC) No 2073/2005 as regards microbiological criteria for sprouts and the sampling rules for poultry carcases and fresh poultry meat (Text with EEA relevance).

Slaughterhouses shall include in their sampling plans poultry carcases from flocks with an unknown salmonella status or with a status known to be positive for *Salmonella enteritidis* or *Salmonella typhimurium*.

When testing against the process hygiene criterion set out in Row 2.1.5 of Chapter 2 for salmonella in poultry carcases in slaughterhouses, neck skins from a minimum of 15 poultry carcases shall be sampled at random after chilling during each sampling session. A piece of approximately 10 g from neck skin shall be obtained from each poultry carcase. On each occasion the neck skin samples from three poultry carcases from the same flock of origin shall be pooled before examination in order to form 5 x 25 g final samples. These samples shall also be used to verify the compliance with the food safety criterion set out in Row 1.28 of Chapter 1.

[^{F10}For the *Salmonella* analyses for fresh poultry meat other than poultry carcases, five samples of at least 25 g of the same batch shall be collected. The sample taken from poultry portions with skin shall contain skin and a thin surface muscle slice in case the amount of skin is not sufficient to form a sample unit. The sample taken from poultry portions without skin or with only a small amount of skin shall contain a thin surface muscle slice or slices added to any skin present to make a sufficient sample unit. The slices of meat shall be taken in a way that includes as much as possible of the surface of the meat.]

Guidelines for sampling

More detailed guidelines on the sampling of carcases, in particular concerning the sampling sites, may be included in the guides to good practice referred to in Article 7 of Regulation (EC) No 852/2004.

Sampling frequencies for carcases, minced meat, meat preparations, mechanically separated meat and fresh poultry meat

The food business operators of slaughterhouses or establishments producing minced meat, meat preparations, mechanically separated meat or fresh poultry meat shall take samples for microbiological analysis at least once a week. The day of sampling shall be changed each week to ensure that each day of the week is covered.

As regards the sampling of minced meat and meat preparations for *E. coli* and aerobic colony count analyses and the sampling of carcases for Enterobacteriaceae and aerobic colony count analyses, the frequency may be reduced to fortnightly testing if satisfactory results are obtained for six consecutive weeks.

In the case of sampling for salmonella analyses of minced meat, meat preparations, carcases and fresh poultry meat, the frequency may be reduced to fortnightly if satisfactory results have been obtained for 30 consecutive weeks. The salmonella sampling frequency may also be reduced if there is a national or regional salmonella control programme in place and if this programme includes testing that replaces the sampling laid down in this paragraph. The sampling frequency may be further reduced if the national or regional salmonella control programme demonstrates that the salmonella prevalence is low in animals purchased by the slaughterhouse.

However, when justified on the basis of a risk analysis and consequently authorised by the competent authority, small slaughterhouses and establishments producing minced meat, meat preparations and fresh poultry meat in small quantities may be exempted from these sampling frequencies.

[^{F7}3.3 *Sampling rules for sprouts*

For the purposes of this Section, the definition of batch in Article 2(b) of Implementing Regulation (EU) No 208/2013 will apply.

A. *General rules for sampling and testing*

1. Preliminary testing of the batch of seeds

Food business operators producing sprouts shall carry out a preliminary testing of a representative sample of all batches of seeds. A representative sample shall include at least 0,5 % of the weight of the batch of seeds in sub samples of 50 g or be selected based on a structured statistically equivalent sampling strategy verified by the competent authority.

For the purposes of performing the preliminary testing, the food business operator must sprout the seeds in the representative sample under the same conditions as the rest of the batch of seeds to be sprouted.

2. Sampling and testing of the sprouts and the spent irrigation water

Food business operators producing sprouts shall take samples for microbiological testing at the stage where the probability of finding Shiga toxin producing *E. coli* (STEC) and *Salmonella* spp. is the highest, in any case not before 48 hours after the start of the sprouting process.

Samples of sprouts shall be analysed according to the requirements in rows 1.18 and 1.29 of Chapter 1.

However, if a food business operator producing sprouts has a sampling plan, including sampling procedures and sampling points of the spent irrigation water, they may replace the sampling requirement under the sampling plans set out in rows 1.18 and 1.29 of Chapter 1 with the analysis of 5 samples of 200 ml of the water that was used for the irrigation of the sprouts.

Status: Point in time view as at 01/06/2014.
Changes to legislation: There are currently no known outstanding effects for the
Commission Regulation (EC) No 2073/2005. (See end of Document for details)

In that case requirements set out in rows 1.18 and 1.29 of Chapter 1 shall apply to the analysis of the water that was used for the irrigation of the sprouts, with the limit of absence in 200 ml.

When testing a batch of seeds for the first time, food business operators may only place sprouts on the market if the results of the microbiological analysis comply with rows 1.18 and 1.29 of Chapter 1, or the limit of absence in 200 ml if they analyse spent irrigation water.

3. Sampling frequency

Food business operators producing sprouts shall take samples for microbiological analysis at least once a month at the stage where the probability of finding Shiga toxin producing *E. coli* (STEC) and *Salmonella* spp. is the highest, in any case not before 48 hours after the start of the sprouting process.

B. Derogation from the preliminary testing of all batches of seeds set out in point A.1 of this Section

When justified on the basis of the following conditions and authorised by the competent authority, food business operators producing sprouts may be exempted from the sampling set out in point A.1 of this Section:

- (a) the competent authority is satisfied that the food business operator implements a food safety management system in that establishment, which may include steps in the production process, which reduces the microbiological risk; and,
- (b) historical data confirms that during at least 6 consecutive months prior to granting the authorisation, all batches of the different types of sprouts produced in the establishment comply with the food safety criteria set out in rows 1.18 and 1.29 of Chapter 1.]]]

ANNEX II

The studies referred to in Article 3(2) shall include:

- specifications for physico-chemical characteristics of the product, such as pH, a_w, salt content, concentration of preservatives and the type of packaging system, taking into account the storage and processing conditions, the possibilities for contamination and the foreseen shelf-life, and
- consultation of available scientific literature and research data regarding the growth and survival characteristics of the micro-organisms of concern.

When necessary on the basis of the abovementioned studies, the food business operator shall conduct additional studies, which may include:

- predictive mathematical modelling established for the food in question, using critical growth or survival factors for the micro-organisms of concern in the product,
- tests to investigate the ability of the appropriately inoculated micro-organism of concern to grow or survive in the product under different reasonably foreseeable storage conditions,
- -- studies to evaluate the growth or survival of the micro-organisms of concern that may be present in the product during the shelf-life under reasonably foreseeable conditions of distribution, storage and use.

The above mentioned studies shall take into account the inherent variability linked to the product, the micro-organisms in question and the processing and storage conditions.

(1) [^{F1}The test results may be used also for demonstrating the effectiveness of the hazard analysis and critical control point principles or good hygiene procedure of the process.]

Textual Amendments

F1 Substituted by Commission Regulation (EC) No 1441/2007 of 5 December 2007 amending Regulation (EC) No 2073/2005 on microbiological criteria for foodstuffs (Text with EEA relevance).

Status:

Point in time view as at 01/06/2014.

Changes to legislation:

There are currently no known outstanding effects for the Commission Regulation (EC) No 2073/2005.