

Commission Regulation (EC) No 2075/2005 of 5 December  
2005 laying down specific rules on official controls for  
*Trichinella* in meat (Text with EEA relevance) (repealed)

COMMISSION REGULATION (EC) No 2075/2005  
of 5 December 2005

laying down specific rules on official controls for *Trichinella* in meat  
(Text with EEA relevance) (repealed)

THE COMMISSION OF THE EUROPEAN COMMUNITIES,

Having regard to the Treaty establishing the European Community,

Having regard to Regulation (EC) No 854/2004 of the European Parliament and of the Council of 29 April laying down specific rules for the organisation of official controls on products of animal origin intended for human consumption<sup>(1)</sup>, and in particular points 9 and 10 of Article 18 thereof,

Whereas:

- (1) Regulation (EC) No 853/2004 of the European Parliament and of the Council of 29 April 2004 laying down specific hygiene rules for food of animal origin<sup>(2)</sup>, Regulations (EC) No 854/2004 and (EC) No 882/2004 of the European Parliament and of the Council of 29 April 2004 on the official controls performed to ensure the verification of compliance with feed and food law, animal health and animal welfare rules<sup>(3)</sup> lay down the health rules and requirements regarding food of animal origin and the official controls required.
- (2) In addition to those rules, more specific requirements should be laid down for *Trichinella*. Meat of domestic swine, wild boar, horses and other animal species may be infested with nematodes of the genus *Trichinella*. Consumption of meat infested with *Trichinella* can cause serious disease in humans. Measures should be put in place to prevent human disease caused by the consumption of meat infested with *Trichinella*.
- (3) On 22 November 2001, the Scientific Committee on Veterinary Measures relating to Public Health adopted an opinion on trichinellosis, epidemiology, methods of detection and *Trichinella*-free pig production. On 1 December 2004, the Scientific Panel on biological hazards (Biohaz) of the European Food Safety Authority adopted an opinion on the suitability and details of freezing methods to allow human consumption of meat infected with *Trichinella* or *Cysticercus*. On 9 and 10 March 2005, Biohaz adopted an opinion on risk assessment of a revised inspection of slaughter animals in areas with low prevalence of *Trichinella*.
- (4) Council Directive 77/96/EEC of 21 December 1976 on the examination for trichinae (*trichinella spiralis*) upon importation from third countries of fresh meat derived from domestic swine<sup>(4)</sup> was repealed by Directive 2004/41/EC of the European Parliament

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

and of the Council of 21 April 2004 repealing certain Directives concerning food hygiene and health conditions for the production and placing on the market of certain products of animal origin intended for human consumption and amending Council Directives 89/662/EEC and 92/118/EEC and Council Decision 95/408/EC<sup>(5)</sup>.

- (5) Various laboratory methods have been approved for the detection of *Trichinella* in fresh meat. The magnetic stirrer method for pooled-sample digestion is recommended as a reliable method for routine use. Sample size for parasitic analysis should be increased if the sample cannot be collected at the predilection site and if the type or species of animal is at higher risk of being infected. Trichinoscopic examination fails to detect non-encapsulated *Trichinella* species infecting domestic and sylvatic animals and humans and is no longer suitable as a detection method for standard use. The trichinoscopic method should only be used under exceptional circumstances for the examination of a small number of animals slaughtered per week, provided that measures are taken by the food business operator to process the meat in such a way that it is completely safe for consumption. However, the method should be replaced by a more reliable detection method within a transitional period. Other methods, such as serological tests, can be useful for monitoring purposes once the tests have been validated by a Community reference laboratory as soon as such a laboratory has been appointed by the Commission. Serological tests are not suitable for detecting *Trichinella* infestation in individual animals intended for human consumption.
- (6) Freezing meat under specified conditions can kill any parasites present but certain *Trichinella* species occurring in game and horses are resistant when freezing is carried out using the recommended temperature and time combinations.
- (7) Holdings should be officially recognised by the competent authority as *Trichinella*-free, provided specific conditions are met. Fattening pigs coming from such holdings should be exempted from inspection for *Trichinella*. Categories of holdings should be officially recognised by the competent authority as *Trichinella*-free, provided specific conditions are met. Such recognition should reduce the number of on-site inspections to be carried out by the competent authority, but is only feasible in Member States with a history of very low disease prevalence.
- (8) Regular monitoring of domestic swine, wild boar, horses and foxes or other indicator animals is an important tool for assessing changes in disease prevalence. The results of such monitoring should be communicated in an annual report in accordance with Directive 2003/99/EC of the European Parliament and of the Council of 17 November 2003 on the monitoring of zoonoses and zoonotic agents<sup>(6)</sup>.
- (9) Regulation (EC) No 853/2004 does not apply to wild game or wild game meat directly supplied to the final consumer or to local retail establishments directly supplying the final consumer. It should therefore be the responsibility of the Member States to adopt national measures to mitigate the risk of *Trichinella*-infested wild boar meat reaching the final consumer.
- (10) The measures provided in this Regulation are in accordance with the opinion of the Standing Committee on the Food Chain and Animal Health,

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

HAS ADOPTED THIS REGULATION:

## CHAPTER I

### GENERAL PROVISION

#### *[<sup>F1</sup>Article 1*

##### **Definitions**

For the purposes of this Regulation, the following definitions shall apply:

- (1) ‘*Trichinella*’ means any nematode belonging to the species of the genus *Trichinella*;
- (2) ‘controlled housing conditions’ means a type of animal husbandry where swine are kept at all times under conditions controlled by the food business operator with regard to feeding and housing;
- (3) ‘compartment’ means a group of holdings which apply controlled housing conditions. All holdings applying controlled housing conditions in a Member States, may be considered as one compartment.]

#### **Textual Amendments**

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

## CHAPTER II

### **OBLIGATIONS OF COMPETENT AUTHORITIES AND OF FOOD BUSINESS OPERATORS**

#### *[<sup>F1</sup>Article 2*

##### **Sampling of carcasses**

1 Carcasses of domestic swine shall be sampled in slaughterhouses as part of the post-mortem examination as follows:

- a all carcasses of breeding sows and boars or at least 10 % of carcasses of animals sent in for slaughter each year from each holding being officially recognised as applying controlled housing conditions, shall be examined for *Trichinella*;
- b all carcasses from holdings not being officially recognised as applying controlled housing conditions shall be systematically examined for *Trichinella*.

A sample shall be collected from each carcase and the sample shall be examined for *Trichinella*, in a laboratory designated by the competent authority, using one of the following methods of detection:

- a the reference method of detection set out in Chapter I of Annex I; or

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

b an equivalent method of detection set out in Chapter II of Annex I.

2 Pending the results of the *Trichinella* examination and provided full traceability is guaranteed by the food business operator, such carcasses may be cut up into a maximum of six parts in a slaughterhouse or in a cutting plant on the same premises as the slaughterhouse (the premises).

By way of derogation from the first subparagraph and following approval by the competent authority, such carcasses may be cut up at a cutting plant attached to or separate from the slaughterhouse provided that:

- a the procedure is under supervision by the competent authority;
- b a carcass or the parts thereof have not more than one cutting plant as its destination;
- c the cutting plant is situated within the territory of the Member State; and
- d in case of a positive result all the parts are declared unfit for human consumption.

3 Carcasses of horses, wild boar and other farmed and wild animal species susceptible to *Trichinella* infestation shall be systematically sampled in slaughterhouses or game-handling establishments as part of the post-mortem examination.

A sample shall be collected from each carcass and the sample shall be examined in accordance with Annexes I and III in a laboratory designated by the competent authority.

#### Textual Amendments

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

### Article 3

#### Derogations

1 By way of derogation from Article 2(1), meat of domestic swine that has undergone a freezing treatment in accordance with Annex II under the supervision of the competent authority shall be exempt from *Trichinella* examination.

2 By way of derogation from Article 2(1), carcasses and meat of not weaned domestic swine less than five weeks of age shall be exempt from *Trichinella* examination.

3 By way of derogation from Article 2(1), carcasses and meat of domestic swine may be exempt from *Trichinella* examination where the animals come from a holding or a compartment officially recognised as applying controlled housing conditions in accordance with Annex IV, if:

- a no autochthonous *Trichinella* infestations in domestic swine kept in holdings officially recognised as applying controlled housing conditions have been detected in the Member State in the past three years, during which time continuous testing has been conducted in accordance with Article 2; or
- b historical data on continuous testing carried out on slaughtered swine population provide at least 95 % confidence that the prevalence of *Trichinella* does not exceed 1 per million in that population; or
- c the holdings applying controlled housing conditions are located in Belgium or Denmark.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

4 Where a Member State implements the derogation provided for in paragraph 3, the Member State concerned shall inform the Commission and the other Member States at the Standing Committee of the Food Chain and Animal Health and submit an annual report to the Commission containing the information referred to in Chapter II of Annex IV. The Commission shall publish on its website the list of Member States implementing the derogation.

Where a Member State fails to submit that annual report or the annual report is unsatisfactory for the purposes of this Article, then the derogation shall cease to apply to that Member State.]

---

**Textual Amendments**

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

*Article 4*

**Trichinella examination and application of health mark**

1 Carcasses as referred to in Article 2 or parts thereof, except for those referred to in Article 2(2)(b), may not leave the premises, before the result of the *Trichinella* examination is found to be negative.

Similarly, other parts of an animal intended for human or animal consumption which contain striated muscle tissue may not leave the premises before the result of the *Trichinella* examination is found to be negative.

2 Animal waste and animal by-products not intended for human consumption and not containing striated muscle may leave the premises before the results of the *Trichinella* examination are available.

However, the competent authority may require a *Trichinella* examination or prior treatment of animal by-products to be carried out before permitting them to leave the premises.

[<sup>F23</sup> Where a procedure is in place in the slaughterhouse to ensure that no part of carcasses examined leaves the premises until the result of the *Trichinella* examination is found to be negative and the procedure is formally approved by the competent authority or where the derogation provided for in Article 2(2)(b) applies, the health mark provided for in Article 5(2) of Regulation (EC) No 854/2004 may be applied before the results of the *Trichinella* examination are available.]

---

**Textual Amendments**

- F2** Substituted by [Commission Regulation \(EC\) No 1665/2006 of 6 November 2006 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

## Article 5

### Training

The competent authority shall ensure that all personnel involved in the examination of samples to detect *Trichinella* shall be properly trained and participate in:

- (a) a quality control programme of the tests used to detect *Trichinella*; and
- (b) a regular assessment of the testing, recording and analysis procedures used in the laboratory.

## Article 6

### Methods of detection

1 The methods of detection set out in Chapters I and II of Annex I shall be used for examining samples as referred to in Article 2:

- a where they provide grounds for suspecting *Trichinella* infestation; or
- b when samples coming from the same holding were previously found to be positive using the trichoscopic method referred to in Article 16(1).

2 All positive samples shall be forwarded to the national reference laboratory or the Community reference laboratory for determination of the *Trichinella* species involved.

## Article 7

### Contingency plans

The competent authorities of the Member States shall prepare a contingency plan by 31 December 2006 outlining all action to be taken where samples as referred to in Articles 2 and 16 test positive to *Trichinella*. That plan shall include details covering:

- (a) traceability of infested carcass(es) and parts thereof containing muscle tissue;
- (b) measures for dealing with infested carcass(es) and parts thereof;
- (c) investigation of the source of infestation and any spreading among wildlife;
- (d) any measures to be taken at the retail or consumer level;
- (e) measures to be taken where the infested carcass cannot be identified at the slaughterhouse;
- (f) determination of the *Trichinella* species involved.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

## *F1* Article 8

### **Official recognition of holdings applying controlled housing conditions**

1 For the purpose of this Regulation, the competent authority may officially recognise a holding or a compartment applying controlled housing conditions where the requirements laid down in Annex IV are complied with.

2 Holdings or a compartment applying controlled housing conditions in Denmark or Belgium in accordance with Article 3, paragraph 3(c) at the date of application of this Regulation are considered to be officially recognised applying controlled housing conditions as listed in Annex IV to this Regulation.

---

#### **Textual Amendments**

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

## Article 9

### **Obligation on food business operators to inform**

Food business operators of holdings officially recognised as applying controlled housing conditions shall inform the competent authority of any requirement as laid down in Annex IV that is no longer fulfilled or of any other change that might affect holdings' *Trichinella* status.

---

#### **Textual Amendments**

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

## Article 10

### **Audits of holdings officially recognised as applying controlled housing conditions**

The competent authority shall ensure that audits are carried out periodically of holdings officially recognised as applying controlled housing conditions.

The frequency of audits shall be risk-based, taking account of disease history and prevalence, previous findings, the geographical area, local susceptible wildlife, animal husbandry practices, veterinary supervision and farmers' compliance.

The competent authority shall verify that domestic swine coming from those holdings are examined in accordance with Article 2(1).

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

#### Textual Amendments

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

### Article 11

#### Monitoring programmes

The competent authority may implement a monitoring programme covering the population of domestic swine coming from a holding or a compartment officially recognised as applying controlled housing conditions, in order to verify that *Trichinella* is effectively absent in that population.

The frequency of testing, the number of animals to be tested and the sampling plan shall be laid down in the monitoring programme. To that end, meat samples shall be collected and examined for presence of *Trichinella* parasites in accordance with Chapter I or II of Annex I.

The monitoring programme may include serological methods as an additional tool once a suitable test is validated by the EU reference laboratory.

#### Textual Amendments

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

### Article 12

#### Withdrawal of official recognition of holdings as applying controlled housing conditions

1 Where the results of the audits carried out in accordance with Article 10 show that the requirements of Annex IV are no longer fulfilled, the competent authority shall withdraw the holdings official recognition without delay.

2 Where domestic swine from a holding officially recognised as applying controlled housing conditions test positive to *Trichinella*, the competent authority shall without delay:

- a withdraw the holding's official recognition;
- b examine all domestic swine of that holding at the time of slaughter;
- c trace and test all breeding animals that arrived on the holding and, as far as possible, all those that left the holding in at least the six months preceding the positive finding; to that end, meat samples shall be collected and examined for presence of *Trichinella* parasites using the detection methods laid down in Chapters I and II of Annex I;
- d when relevant, as far as is feasible, investigate the spread of parasite infestation due to the distribution of meat from domestic swine slaughtered in the period preceding the positive finding;
- e inform the Commission and the other Member States;



---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- f When relevant, initiate an epidemiological investigation to elucidate the cause of infestation;
- g take appropriate measures where any infested carcass cannot be identified at the slaughterhouse, including:
- (i) increasing the size of each meat sample collected for testing of the suspect carcasses; or
  - (ii) declaring the carcasses unfit for human consumption;
  - (iii) taking appropriate measures for the disposal of suspect carcasses or parts thereof and those testing positive.
- 3 Following withdrawal of recognition, holdings may be officially recognised again once the problems identified have been solved and the requirements laid down in Annex IV are fulfilled to the satisfaction of the competent authority.
- 4 If the inspection identified a lack of compliance with Article 9 or positive testing in a holding of a compartment, the holding concerned should be removed from the compartment until compliance is re-established.]

---

**Textual Amendments**

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

CHAPTER III

**IMPORTS**

*[<sup>F1</sup>Article 13*

**Import health requirements**

Meat of animal species that may be carriers of *Trichinella*, containing striated muscles and coming from a third country may only be imported into the Union if examination for *Trichinella* has been performed in accordance with Articles 2 and 3 in that third country before export.]

---

**Textual Amendments**

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

*<sup>F3</sup>Article 14*

**[<sup>F3</sup>Derogations from Article 13]**

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

#### Textual Amendments

- F3** Deleted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

### *[<sup>F1</sup>Article 15*

#### Documents

The health certificate accompanying imports of meat as referred to in Article 13 shall be endorsed with a statement by the official veterinarian to the effect that the examination for *Trichinella* in the third country of origin has been performed in accordance with Article 13.

That document shall accompany the meat in the original unless an exemption has been granted in accordance with Article 14(4) of Regulation (EC) No 854/2004.]

#### Textual Amendments

- F1** Substituted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

## CHAPTER IV

### TRANSITIONAL AND FINAL PROVISIONS

#### *Article 16*

#### Transitional provisions

1 The Member State may allow the trichinoscopic method set out in Chapter III of Annex I to be used for domestic swine and wild boar in exceptional cases until 31 December 2009, where:

- a single carcasses as referred to in Article 2 need to be examined individually in an establishment that does not slaughter more than 15 domestic swine per day or 75 domestic swine per week or prepare for placing on the market more than 10 wild boar per day; and
- b the detection methods set out in Chapters I and II of Annex I are not available.

2 Where the trichinoscopic method is used, the competent authority shall ensure that:

- a the meat is marked with a health mark that is clearly different from the health mark provided for in Article 5(1)(a) of Regulation (EC) No 853/2004, and the meat is supplied directly to the final consumer or to retail establishments directly supplying the final consumer; and
- b the meat is not used for the production of products where the production process does not kill *Trichinella*.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

### *Article 17*

#### **Entry into force**

This Regulation shall enter into force on the 20th day following its publication in the *Official Journal of the European Union*.

It shall apply from 1 January 2006.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

## ANNEX I

### Detection methods

#### CHAPTER I

##### REFERENCE METHOD OF DETECTION

###### Magnetic stirrer method for pooled sample digestion

1. Apparatus and reagents
  - (a) Knife or scissors and tweezers for cutting specimens
  - (b) Trays marked off into 50 squares, each of which can hold samples of approximately 2 g of meat, or other tools giving equivalent guarantees as regards the traceability of the samples
  - (c) A blender with a sharp chopping blade. Where the samples are larger than 3 g, a meat mincer with openings of 2 to 4 mm or scissors must be used. In the case of frozen meat or tongue (after removal of the superficial layer, which cannot be digested), a meat mincer is necessary and the sample size will need to be increased considerably
  - (d) Magnetic stirrers with thermostatically controlled heating plate and teflon-coated stirring rods approximately 5 cm long
  - (e) Conical glass separation funnels, capacity of at least 2 litres, preferably fitted with teflon safety plugs
  - (f) Stands, rings and clamps
  - (g) Sieves, mesh size 180 microns, external diameter 11 cm, with stainless steel mesh
  - (h) Funnels, internal diameter not less than 12 cm, to support the sieves
  - (i) Glass beakers, capacity 3 litres
  - (j) Glass measuring cylinders, capacity 50 to 100 ml, or centrifuge tubes
  - (k) A trichinoscope with a horizontal table or a stereo-microscope, with a substage transmitted light source of adjustable intensity
  - (l) A number of 9 cm diameter petri dishes (for use with a stereo-microscope), marked on their undersides into 10 × 10 mm square examination areas using a pointed instrument
  - (m) A larval counting basin (for use with a trichinoscope), made of 3 mm thick acrylic plates as follows:
    - (i) the bottom of the basin to be 180 × 40 mm, marked off into squares,
    - (ii) the sides to be 230 × 20 mm,
    - (iii) the end to be 40 × 20 mm. The bottom and the ends must be inserted between the sides, to form two small handles at the ends. The upper side of the bottom must be raised 7 to 9 mm from the base of the frame formed by the sides and the ends. The components must be stuck together with glue suitable for the material
  - (n) Aluminium foil

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

- (o) 25 % hydrochloric acid
- (p) [<sup>F4</sup>Pepsin, strength: 1: 10 000 NF (US National Formulary) corresponding to 1: 12 500 BP (British Pharmacopoeia) and to 2 000 FIP (Fédération internationale de pharmacie), or stabilized liquid pepsin with minimum 660 European Pharmacopoeia units/ml]
- (q) Tap water heated to 46 to 48 °C
- (r) A balance accurate to at least 0,1 g
- (s) Metal trays, capacity 10 to 15 litres, to collect the remaining digestive juice
- (t) Pipettes of different sizes (1, 10 and 25 ml) and pipette holders
- (u) A thermometer accurate to 0,5 °C within the range 1 to 100 °C
- (v) Siphon for tap water.

#### Textual Amendments

- F4** Substituted by [Commission Regulation \(EC\) No 1245/2007 of 24 October 2007 amending Annex I to Regulation \(EC\) No 2075/2005](#), as regards the use of liquid pepsin for the detection of *Trichinella* in meat (Text with EEA relevance).

2. Collecting of specimens and quantity to be digested
  - (a) In the case of whole carcasses of domestic swine, a specimen weighing at least 1 g is to be taken from a pillar of the diaphragm at the transition to the sinewy part. Special trichinae forceps can be used provided an accuracy of between 1,0 and 1,15 g can be guaranteed.

In the case of breeding sows and boars, a larger sample weighing at least 2 g is to be taken from a pillar of the diaphragm at the transition to the sinewy part.

In the absence of diaphragm pillars, a specimen of twice the size 2 g (or 4 g in the case of breeding sows and boars) is to be taken from the rib part or the breastbone part of the diaphragm, or from the jaw muscle, tongue or abdominal muscles.
  - (b) For cuts of meat, a sample weighing at least 5 g of striated muscle, containing little fat is to be taken, where possible from close to bones or tendons. A sample of the same size is to be collected from meat that is not intended to be cooked thoroughly or other types of post-slaughter processing.
  - (c) For frozen samples, a sample weighing at least 5 g of striated muscle tissue is to be taken for analysis.

The weight of meat specimens relates to a sample of meat that is free of all fat and fascia. Special attention must be paid when collecting muscle samples from the tongue in order to avoid contamination with the superficial layer of the tongue, which is indigestible and can prevent reading of the sediment.
3. Procedure
  - I. Complete pools (100 g of samples at a time)

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- (a)  $16 \pm 0,5$  ml of hydrochloric acid is added to a 3 litre beaker containing 2,0 litre of tap water, preheated to 46 to 48 °C; a stirring rod is placed in the beaker, the beaker is placed on the preheated plate and the stirring is started.
- (b) [<sup>F4</sup>10  $\pm 0,2$  g of pepsin or 30  $\pm 0,5$  ml liquid pepsin is added.]
- (c) 100 g of samples collected in accordance with point 2 is chopped in the blender.
- (d) The chopped meat is transferred to the 3 litre beaker containing the water, pepsin and hydrochloric acid.
- (e) The mincing insert of the blender is immersed repeatedly in the digestion fluid in the beaker and the blender bowl is rinsed with a small quantity of digestion fluid to remove any meat still adhering.
- (f) The beaker is covered with aluminium foil.
- (g) The magnetic stirrer must be adjusted so that it maintains a constant temperature of 44 to 46 °C throughout the operation. During stirring, the digestion fluid must rotate at a sufficiently high speed to create a deep whirl without splashing.
- (h) The digestion fluid is stirred until the meat particles disappear (approximately 30 minutes). The stirrer is then switched off and the digestion fluid is poured through the sieve into the sedimentation funnel. Longer digestion times may be necessary (not exceeding 60 minutes) in the processing of certain types of meat (tongue, game meat, etc.).
- (i) The digestion process is considered satisfactory if not more than 5 % of the starting sample weight remains on the sieve.
- (j) The digestion fluid is allowed to stand in the funnel for 30 minutes.
- (k) After 30 minutes, a 40 ml sample of digestion fluid is quickly run off into the measuring cylinder or centrifuge tube.
- (l) The digestion fluids and other liquid waste are kept in a tray until reading of the results is completed.
- (m) The 40 ml sample is allowed to stand for 10 minutes. 30 ml of supernatant is then carefully withdrawn by suction to remove the upper layers and leave a volume of not more than 10 ml.
- (n) The remaining 10 ml sample of sediment is poured into a larval counting basin or petri dish.
- (o) The cylinder or centrifuge tube is rinsed with not more than 10 ml of tap water, which has to be added to the sample in the larval counting basin or petri dish. Subsequently, the sample is examined by trichinoscope or stereo-microscope at a 15 to 20 times magnification. Visualisation using other techniques is allowed, provided examination of positive control samples has been shown to give an equal or better result than traditional visualisation methods. In all cases of suspect areas or parasite-like shapes, higher magnifications of 60 to 100 times must be used.
- (p) Digests are to be examined as soon as they are ready. Under no circumstances should examination be postponed until the following day.

Where the digests are not examined within 30 minutes of preparation, they must be clarified as follows. The final sample of about 40 ml is poured into a measuring

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

cylinder and allowed to stand for 10 minutes. 30 ml of the supernatant fluid is then removed, leaving a volume of 10 ml. This volume is made up to 40 ml with tap water. After a further settling period of 10 minutes, 30 ml of the supernatant fluid is withdrawn by suction, leaving a volume of no more than 10 ml for examination in a petri dish or larval counting basin. The measuring cylinder is washed with no more than 10 ml of tap water and these washings are added to the sample in the petri dish or the larval counting basin for examination.

If the sediment is found to be unclear on examination, the sample is poured into a measuring cylinder and made up to 40 ml with tap water and then the above procedure is followed. The procedure can be repeated 2 to 4 times until the fluid is clear enough for a reliable reading.

## II. Pools of less than 100 g

Where needed, up to 15 g can be added to a total pool of 100 g and examined together with these samples in accordance with 3(I). More than 15 g must be examined as a complete pool. For pools of up to 50 g, the digestion fluid and the ingredients may be reduced to 1 litre of water, 8 ml of hydrochloric acid and 5 g of pepsin.

## III. Positive or doubtful results

Where examination of a collective sample produces a positive or uncertain result, a further 20 g sample is taken from each pig in accordance with 2(a). The 20 g samples from five pigs are pooled and examined using the method described above. In this way samples from 20 groups of five pigs will be examined.

When *Trichinella* is detected in a pooled sample from five pigs, further 20 g samples are collected from the individual pigs in the group and each is examined separately using the method described above.

Parasite samples are to be kept in 90 % ethyl alcohol for conservation and identification at species level at the Community or national reference laboratory.

After parasite collection, positive fluids (digestive juice, supernatant fluid, washings, etc.) are to be decontaminated by heating to at least 60 °C.

## [<sup>F5</sup>IV. Cleaning and decontamination procedure after a positive or doubtful result

When the examination of a collective or individual sample produces a positive or doubtful latex agglutination result, all material in contact with meat (blender bowl, beaker, stirring rod, temperature sensor, conical filtration funnel, sieve and forceps) must be carefully decontaminated by soaking for few seconds in warm water (65 to 90 °C). Meat residues or inactivated larvae that could remain on their surface may be removed with a clean sponge and tap water. If required, a few drops of detergent can be added for degreasing equipment. It is then recommended to rinse each piece thoroughly to remove all traces of detergent.]

### Textual Amendments

- F5** Inserted by [Commission Regulation \(EU\) No 216/2014 of 7 March 2014 amending Regulation \(EC\) No 2075/2005 laying down specific rules on official controls for \*Trichinella\* in meat \(Text with EEA relevance\)](#).

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

## CHAPTER II

### EQUIVALENT METHODS

- A. Mechanically assisted pooled sample digestion method/sedimentation technique
1. Apparatus and reagents
    - (a) Knife or scissors for cutting specimens
    - (b) Trays marked off with 50 squares, each of which can hold samples of approximately 2 g of meat, or other tools giving equivalent guarantees as regards the traceability of the samples
    - (c) Meat mincer or electrical blender
    - (d) A stomacher lab-blender 3 500 thermo model
    - (e) Plastic bags suitable for the stomacher lab-blender
    - (f) Conical separation funnels, capacity 2 litres, preferably fitted with teflon safety plugs
    - (g) Stands, rings and clamps
    - (h) Sieves, mesh size 180 microns, external diameter 11 cm, with stainless steel or brash mesh
    - (i) Funnels, internal diameter not less than 12 cm, to support the sieves
    - (j) 100 ml glass measuring cylinders
    - (k) A thermometer accurate to 0,5 °C within the range 1 to 100 °C
    - (l) A vibrator, e.g. an electric shaver with the head removed
    - (m) A relay which will switch on and off at one-minute intervals
    - (n) A trichinoscope with a horizontal table or a stereo-microscope, with a sub-stage transmitted light source of adjustable intensity
    - (o) A larval counting basin and a number of 9 cm diameter petri dishes as in Chapter I(1) (l) and (m)
    - (p) 17,5 % hydrochloric acid
    - (q) [<sup>F4</sup>Pepsin, strength: 1: 10 000 NF (US National Formulary) corresponding to 1: 12 500 BP (British Pharmacopoeia) and to 2 000 FIP (Fédération internationale de pharmacie), or stabilized liquid pepsin with minimum 660 European Pharmacopoeia units/ml]
    - (r) A number of 10 litre bins to be used for decontamination of apparatus, e.g. with formol, and for digestive juice remaining where specimens test positive
    - (s) A balance accurate to 0,1 g.
  2. Collecting of specimens and quantity to be digested  
As stipulated in Chapter I(2).
  3. Procedure



---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

## I. Grinding

Grinding the meat samples in a meat mincer beforehand will improve the digestion quality. If an electrical blender is used, the blender must be operated three to four times for approximately one second each time.

## II. Digestion procedure

This procedure may involve complete pools (100 g of samples at a time) or pools of less than 100 g.

### (a) Complete pools (100 samples at a time)

- (i) The stomacher lab-blender 3 500 is fitted with a double plastic bag and the temperature control set at 40 to 41 °C.
- (ii) One and a half litres of water preheated to 40 to 41 °C is poured into the inner plastic bag.
- (iii) 25 ml of 17,5 % hydrochloric acid is added to the water in the stomacher.
- (iv) 100 samples weighing approximately 1 g each (at 25 to 30 °C) taken from each individual sample in accordance with 2 are added.
- (v) [F<sup>4</sup>Lastly, 6 g pepsin or 18 ml liquid pepsin is added. This order must be followed strictly to avoid decomposition of the pepsin.]
- (vi) The stomacher is then allowed to pound the content of the bag for 25 minutes.
- (vii) The plastic bag is removed from the stomacher and the digestion fluid is filtered through the sieve into a 3 litre beaker.
- (viii) The plastic bag is washed with approximately 100 ml of water, which is then used to rinse the sieve and lastly added to the filtrate in the beaker.
- (ix) Up to 15 individual samples can be added to a total pool of 100 samples and examined together with these samples.

### (b) Smaller pools (less than 100 samples)

- (i) The stomacher lab-blender 3 500 is fitted with a double plastic bag and the temperature control set at 40 to 41 °C.
- (ii) A digestion fluid is prepared by mixing about one and a half litres of water and 25 ml of 17,5 % hydrochloric acid. 6 g of pepsin is added and the whole mixed at a temperature of 40 to 41 °C. This order must be followed strictly to avoid decomposition of the pepsin.
- (iii) Of the digestion fluid, a volume corresponding to 15 ml per gram of sample is measured (e.g. for 30 samples the volume required is  $30 \times 15 \text{ ml} = 450 \text{ ml}$ ) and transferred to the inner of the two plastic bags, together with the meat samples weighing approximately 1 g (at 25 to 30 °C) taken from each individual sample in accordance with 2.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- (iv) Water at a temperature of approximately 41 °C is poured into the outer bag to make up a total volume in the two bags of one and a half litres. The stomacher is then allowed to pound the content of the bag for 25 minutes.
- (v) The plastic bag is removed from the stomacher and the digestion fluid is filtered through the sieve into a 3 litre beaker.
- (vi) The plastic bag is washed with approximately 100 ml of water (at 25 to 30 °C), which is then used to rinse the sieve and lastly added to the filtrate in the beaker.

### III. Recovery of larvae by sedimentation

- Ice (300 to 400 g of ice flakes, scaly ice or crushed ice) is added to the digestion fluid to bring its volume up to about 2 litres. The digestion fluid is then stirred until the ice has melted. In the case of smaller pools (see II(b)), the amount of ice must be reduced correspondingly.
- The chilled digestion fluid is transferred to a 2 litre separation funnel, equipped with a vibrator in an extra clamp.
- Sedimentation is allowed to proceed for 30 minutes, during which time the sedimentation funnel is vibrated intermittently, i.e. one minute vibration followed by a one-minute pause.
- After 30 minutes, a 60 ml sample of the sediment is quickly run off into a 100 ml measuring cylinder (the funnel is rinsed with detergent solution after use).
- The 60 ml sample is allowed to stand for at least 10 minutes, after which time the supernatant is withdrawn by suction to leave a volume of 15 ml, to be examined for presence of larvae.
- For suction, a disposable syringe, equipped with a plastic tube, can be used. The length of the tube must be such that 15 ml remains in the measuring cylinder when the flanges of the syringe rest on the cylinder's rim.
- The remaining 15 ml is poured into a larval counting basin or two petri dishes and examined using a trichinoscope or stereo-microscope.
- The measuring cylinder is washed with 5 to 10 ml of tap water and the washings are added to the sample.
- Digests are to be examined as soon as they are ready. Under no circumstances is examination to be postponed until the following day.

Where the digests are unclear or they are not examined within 30 minutes of their preparation, they must be clarified as follows:

- the final sample of 60 ml is poured into a measuring cylinder and allowed to stand for 10 minutes; 45 ml of supernatant fluid is then removed by suction and the remaining 15 ml is made up to 45 ml with tap water,
- after a further settling period of 10 minutes, 30 ml of supernatant fluid is removed by suction and the remaining 15 ml is poured into a petri dish or larval counting basin for examination,
- the measuring cylinder is washed with 10 ml of tap water and these washings are added to the sample in the petri dish or the larval counting basin for examination.

### IV. Positive or doubtful results

Where the result is positive or uncertain, the provisions laid down in Chapter I(3)(III) shall apply.

- B. Mechanically assisted pooled sample digestion method/'on filter isolation' technique

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

## 1. Apparatus and reagents

As stipulated in Chapter II(A)(1).

Additional equipment:

- (a) 1 litre Gelman funnel, complete with filter holder (diameter 45 mm);
- (b) filter discs, consisting of a circular stainless steel mesh with an aperture of 35 microns (disc diameter: 45 mm), two rubber rings 1 mm thick (external diameter: 45 mm; internal diameter: 38 mm), the circular mesh being placed between the two rubber rings and bonded to them using a two-component glue suitable for the two materials;
- (c) an Erlenmeyer flask, capacity 3 litres, fitted with a side tube for suction;
- (d) a filter pump;
- (e) plastic bags, capacity at least 80 ml;
- (f) equipment for sealing the plastic bags;
- (g) rennilase, strength 1: 150 000 soxhlet units per gram.

## 2. Collecting of specimens

As stipulated in Chapter I(2).

### 3. Procedure

#### I. Grinding

Grinding the meat samples in a meat mincer beforehand will improve the digestion quality. If an electrical blender is used, the blender must be operated three to four times for approximately one second each time.

#### II. Digestion procedure

This procedure may involve complete pools (100 g of samples at a time) or pools of less than 100 g.

- (a) Complete pools (100 samples at a time)

See Chapter II(A)(3)(II)(a).

- (b) Smaller pools (less than 100 samples)

See Chapter II(A)(3)(II)(b).

#### III. Recovery of larvae by filtration

- (a) Ice (300 to 400 g of ice flakes, scaly ice or crushed ice) is added to the digestion fluid to bring its volume up to about 2 litres. In the case of smaller pools, the amount of ice must be reduced correspondingly.
- (b) The digestion fluid is stirred until the ice has melted. The chilled digestion fluid is then left for at least three minutes to let the larvae coil.
- (c) The Gelman funnel, fitted with a filter holder and filter disc, is mounted on an Erlenmeyer flask connected to a filter pump.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- (d) The digestion fluid is poured into the Gelman funnel and filtered. Towards the end of filtration, the digestion fluid can be helped to pass through the filter by applying suction with the filter pump. Suction must cease before the filter becomes dry, i.e. when 2 to 5 ml of fluid is left in the funnel.
- (e) Once all the digestion fluid has been filtered, the filter disc is removed and placed in an 80 ml capacity plastic bag, together with 15 to 20 ml of rennilase solution. The rennilase solution is made by adding 2 g of rennilase to 100 ml of tap water.
- (f) The plastic bag is sealed twice and placed between the inner and outer bags in the stomacher.
- (g) The stomacher is allowed to pound for three minutes, e.g. while it is working on a complete or incomplete pool.
- (h) After three minutes, the plastic bag, complete with filter disc and rennilase solution, is removed from the stomacher and opened with scissors. The liquid contents are poured into a larval counting basin or petri dish. The bag is washed out with 5 to 10 ml of water, which is then added to the larval counting basin for examination by trichinoscope or to the petri dish for examination by stereo-microscope.
- (i) Digests must be examined as soon as they are ready. Under no circumstances is examination to be postponed until the following day.

*Note:* Filter discs must never be used when not completely clean. Unclean discs must never be allowed to dry out. Filter discs can be cleaned by leaving them in rennilase solution overnight. Before use, they must be washed in fresh rennilase solution using the stomacher.

#### IV. Positive or doubtful results

Where the result is positive or uncertain, the provisions laid down in Chapter I(3)(III) shall apply.

- C. Automatic digestion method for pooled samples of up to 35 g
  - 1. Apparatus and reagents
    - (a) Knife or scissors for cutting specimens
    - (b) Trays marked off with 50 squares, each of which can hold samples of approximately 2 g of meat, or other tools giving equivalent guarantees as regards the traceability of the samples
    - (c) A Trichomatic 35<sup>®</sup> blender with filtration insert
    - (d) Hydrochloric acid 8,5 ± 0,5 % weight
    - (e) Transparent polycarbonate membrane filters with a diameter of 50 mm and a pore size of 14 microns
    - (f) [F<sup>6</sup>Pepsin, strength 1: 10 000 NF (US National Formulary) corresponding to 1:12 500 BP (British Pharmacopoeia) and to 2 000 FIP (International Pharmaceutical Federation), or stabilised liquid pepsin with minimum 660 European Pharmacopoeia units/ml]
    - (g) A balance accurate to 0,1 g
    - (h) Tweezers with a flat tip

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

- (i) A number of microscope slides with a side-length of at least 5 cm or a number of petri dishes at least 6 cm in diameter, marked on their undersides into 10 × 10 mm square areas using a pointed instrument
- (j) A (stereo-)microscope with transmitted light (magnification 15 to 60 times) or a trichinoscope with a horizontal table
- (k) A bin for collection of waste liquids
- (l) A number of 10 litre bins to be used for decontamination of apparatus, e.g. with formol, and for digestive juice remaining where specimens test positive
- (m) a thermometer accurate to 0,5 °C within the range 1 to 100 °C.

#### Textual Amendments

**F6** Substituted by [Commission Implementing Regulation \(EU\) No 1109/2011 of 3 November 2011 amending Annex I to Regulation \(EC\) No 2075/2005 as regards the equivalent methods for Trichinella testing \(Text with EEA relevance\).](#)

## 2. Collecting of specimens

As stipulated in Chapter I(2).

## 3. Procedure

### I. Digestion procedure

- (a) Place the blender with the filtration insert, connect the waste tube and place the tube so it drains into the waste bin.
- (b) When the blender is switched on, heating will start.
- (c) Before this is done, the bottom valve located below the reaction chamber must be opened and closed.
- (d) Up to 35 samples weighing approximately 1 g each (at 25 to 30 °C) taken from each individual sample in accordance with point 2 are then added. Ensure that larger pieces of tendons are removed as they may clot the membrane filter.
- (e) Pour water up to the edge of a liquid chamber connected to the blender (approximately 400 ml).
- (f) Pour about 30 ml hydrochloric acid ((8,5 %) to the edge of the smaller, connected liquid chamber.
- (g) Place a membrane filter under the coarse filter in the filter holder in the filter insert.
- (h) [<sup>F4</sup>Lastly, add 7 g of pepsin or 21 ml liquid pepsin. This order must be followed strictly to avoid decomposition of the pepsin.]
- (i) Close the lids of the reaction and liquid chambers.
- (j) Select the period of digestion. A short digestion period (5 minutes) must be set for pigs at the normal slaughter age and a longer time (8 minutes) for other samples.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- (k) When the start button on the blender is turned on, the process of dispensing and digestion starts automatically, followed by filtration. After 10 to 13 minutes the process is completed and stops automatically.
- (l) Open the lid of the reaction chamber after checking that the chamber is empty. If there is foam or any digestion liquid remaining in the chamber, repeat the procedure in accordance with V.

#### II. Recovery of larvae

- (a) Remove the filter holder and transfer the membrane filter to a slide or Petri dish.
- (b) Examine the membrane filter using a (stereo-) microscope or a trichinoscope.

#### III. Cleaning equipment

- (a) Where the result is positive, fill the blender reaction chamber with boiling water until it is two-thirds full. Ordinary tap water is poured into the connecting liquid chamber until it covers the lower sensor. Automatic cleaning then takes place. Decontaminate the filter-holder and any other equipment, e.g. using formol.
- (b) After work is completed for the day, fill the blender liquid chamber with water and put it through a standard cycle.

#### IV. Use of membrane filters

Each polycarbonate membrane filter may be used no more than five times. The filter is to be turned between each use. In addition, the filter must be checked after each use for any damage which would make it unsuitable for further use.

#### V. Method to be applied when digestion is incomplete and filtration cannot be carried out

Once the blender has been put through an automatic cycle in accordance with C(3)(I), open the lid of the reaction chamber and check whether there is foam or any liquid remaining in the chamber. If this is the case, proceed as follows:

- (a) close the bottom valve below the reaction chamber;
- (b) remove the filter holder and transfer the membrane filter to a slide or Petri dish;
- (c) put a new membrane filter in the filter holder and attach the filter holder;
- (d) fill the blender liquid chamber with water until the lower sensor is covered;
- (e) carry out the automatic cleaning cycle;
- (f) after the cleaning cycle has ended, open the lid of the reaction chamber and check whether any liquid remains;
- (g) if the chamber is empty, remove the filter holder and transfer the membrane filter to a slide or Petri dish with tweezers;
- (h) examine the two membrane filters in accordance with C(3)(II). If the filters cannot be examined, repeat the entire digestion process with a longer digestion time in accordance with C(3)(I).

#### VI. Positive or doubtful results

Where the result is positive or uncertain, the provisions laid down in Chapter I(3)(III) shall apply.

[<sup>F7</sup>D. **Magnetic stirrer method for pooled sample digestion/‘on filter isolation’ and larva detection by a latex agglutination test**

***This method is only considered equivalent for the testing of meat of domestic swine.***

1. *Apparatus and reagents*
  - (a) Knife or scissors and tweezers for cutting specimens.
  - (b) Trays marked off into 50 squares, each of which can hold samples of approximately 2 g of meat, or other tools giving equivalent guarantees as regards the traceability of the samples.
  - (c) A blender with a sharp chopping blade. Where the samples are larger than 3 g, a meat mincer with openings of 2-4 mm or scissors must be used. In the case of frozen meat or tongue (after removal of the superficial layer, which cannot be digested), a meat mincer is necessary and the sample size will need to be increased considerably.
  - (d) Magnetic stirrers with thermostatically controlled heating plate and Teflon-coated stirring rods approximately 5 cm long.
  - (e) Glass beakers, capacity 3 litres.
  - (f) Sieves, mesh size 180 microns, external diameter 11 cm, with stainless steel mesh.
  - (g) Steel filtration apparatus for 20 µm mesh filters with a steel funnel.
  - (h) Vacuum pump.
  - (i) Metal or plastic tanks, capacity 10-15 litres, to collect the digestive juice.
  - (j) A 3D gyratory rocker.
  - (k) Aluminium foil.
  - (l) 25 % hydrochloric acid.
  - (m) Pepsin, strength: 1:10 000 NF (US National Formulary) corresponding to 1:12 500 BP (British Pharmacopoeia) and to 2 000 FIP (International Pharmaceutical Federation), or stabilised liquid pepsin with minimum 660 European Pharmacopoeia units/ml.
  - (n) Tap water heated to 46-48 °C.
  - (o) A balance accurate to 0,1 g.
  - (p) Pipettes of different sizes (1, 10 and 25 ml), micropipettes according to the latex agglutination manufacturer's instructions and pipette holders.
  - (q) 20 microns nylon mesh filters of a diameter that fits with the filtration system.
  - (r) Plastic or steel forceps of 10-15 cm.
  - (s) Conical vials of 15 ml.
  - (t) A pestle with a Teflon or steel conical tip to fit in the conical vials.
  - (u) A thermometer accurate to 0,5 °C within the range 1-100 °C.
  - (v) Latex agglutination cards of the Trichin-L antigen test kit validated under the code No EURLP\_D\_001/2011.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- (w) Buffer solution with preservative (sample diluent) of the Trichin-L antigen test kit validated under the code No EURLP\_D\_001/2011.
- (x) Buffer supplemented with preservative (negative control) of the Trichin-L antigen test kit validated under the code No EURLP\_D\_001/2011.
- (y) Buffer supplemented with *Trichinella spiralis* antigens and preservative (positive control) of the Trichin-L antigen test kit validated under the code No EURLP\_D\_001/2011.
- (z) Buffer with polystyrene particles coated with antibodies supplemented with preservative (latex beads) of the Trichin-L antigen test kit validated under the code No EURLP\_D\_001/2011.
- (aa) Disposable sticks.

## 2. *Collecting of specimens*

As stipulated in Chapter I(2).

### [<sup>F</sup>13. *Procedure*

- I. For complete pools (100 g of samples at a time)
  - (a)  $16 \pm 0,5$  ml of 25 % hydrochloric acid (0,2 % final) is added to a 3 litre beaker containing 2,0 litres  $\pm$  200 ml of tap water, preheated to 46 to 48 °C; a stirring rod is placed in the beaker, the beaker is placed on the preheated plate and the stirring is started.
  - (b)  $10 \pm 1$  g of powder pepsin (or  $30 \pm 3$  ml of liquid pepsin) is added.
  - (c) 100-115 g of samples collected in accordance with point 2 are chopped in the blender, with  $150 \pm 15$  ml of preheated digestion buffer.
  - (d) The chopped meat is transferred to the 3 litre beaker containing the water, pepsin and hydrochloric acid.
  - (e) The mincing insert of the blender is immersed repeatedly in the digestion fluid in the beaker and the blender bowl is rinsed with a small quantity of digestion fluid to remove any meat still adhering.
  - (f) The beaker is covered with aluminium foil.
  - (g) The magnetic stirrer must be adjusted so that it maintains a constant temperature of 44 to 46 °C throughout the operation. During stirring, the digestion fluid must rotate at a sufficiently high speed to create a deep whirl without splashing.
  - (h) The digestion fluid is stirred until the meat particles disappear (approximately 30 minutes). The stirrer is then switched off and the digestion fluid is poured through the sieve into the sedimentation funnel. Longer digestion times may be necessary (not exceeding 60 minutes) in the processing of certain types of meat (tongue, game meat, etc.).
  - (i) The digestion process is considered satisfactory if not more than 5 % of the starting sample weight remains on the sieve.
  - (j) The 20 microns nylon mesh filter is placed on the filtration support. The conical filtration steel funnel is fixed to the support with the block system and the steel sieve



---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- of 180 microns mesh size is placed on the funnel. The vacuum pump is connected with the filtration support and with the metal or plastic tank, to collect the digestive fluid.
- (k) Stirring is stopped and the digestion fluid is poured into the filtration funnel through the sieve. The beaker is rinsed with approximately 250 ml of warm water. The rinsing liquid is poured into the filtration ramp after the digested fluid has been successfully filtrated.
- (l) The filtration membrane is taken with the forceps, holding it by an edge. The filtration membrane is folded (minimal) in four and put in the 15 ml conical tube. The choice of conical tube must be adapted to the pestle.
- (m) The filtration membrane is pushed at the bottom of the 15 ml conical tube with the help of the pestle and strongly pressed by doing approximately 20 successive back and forth movements with the pestle which should be positioned inside the filtration membrane folding according to the manufacturer's instructions.
- (n)  $0,5 \pm 0,01$  ml of sample diluents is added into the 15 ml conical tube by pipette and the filtration membrane is homogenised with the pestle by doing successive low amplitude back and forth movements for approximately 30 seconds, avoiding abrupt movements to limit liquid splashes according to the manufacturer's instructions.
- (o) Each sample, the negative control, and the positive control, are dispensed into different fields of the agglutination card by pipettes, according to the manufacturer's instructions.
- (p) The latex beads are added into each field of the agglutination card by a pipette, according to the manufacturer's instructions, without making them come into contact with the sample/s and controls. In each field, the latex beads are then gently mixed with a disposable stick until the homogeneous liquid covers the entire field.
- (q) The agglutination card is put on the 3D rocker and is rocked for  $10 \pm 1$  minutes according to the manufacturer's instructions.
- (r) After the time established by the manufacturer's instructions, the rocking is stopped and the agglutination card is put on a plane surface and the reaction results are read immediately, according to the manufacturer's instructions. In the case of a positive sample, the beads aggregates must appear. In the case of a negative sample, the suspension remains homogeneous without beads aggregates.
- II. Pools of less than 100 g as set out in Chapter I(3)(II)

For pools of less than 100 g, the procedure set out in Chapter I(3)(II) must be followed.

III. Positive or doubtful results

Where examination of a collective sample produces a positive or uncertain latex agglutination result, a further 20 g sample is taken from each swine in accordance with Chapter I(2)(a). The 20 g samples from five swine are pooled and examined using the method described in Section I. In this way samples from 20 groups of five swine must be examined.

When a positive latex agglutination is obtained from a group of five swine, further 20 g samples are collected from the individuals in the group and each is examined separately using the method described in Section I.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

When a positive or uncertain latex agglutination result is obtained, at least 20 g of swine muscle must be sent to the national reference laboratory for confirmation using one of the methods described in Chapter I.

Parasite samples must be kept in 90 % ethyl alcohol for conservation and identification at species level at the EU or national reference laboratory.

After parasite collection, positive fluids must be decontaminated by heating to at least 60 °C.

IV. Cleaning and decontamination procedure after a positive or doubtful result.

When the examination of a collective or individual sample produces a positive or doubtful latex agglutination result, all material in contact with meat (blender bowl, beaker, stirring rod, temperature sensor, conical filtration funnel, sieve and forceps) must be carefully decontaminated by soaking for few seconds in warm water (65 to 90 °C). Meat residues or inactivated larvae that could remain on their surface may be removed with a clean sponge and tap water. If required, a few drops of detergent can be added for degreasing equipment. It is then recommended to rinse each piece thoroughly to remove all traces of detergent.]]

#### Textual Amendments

- F7** Inserted by [Commission Implementing Regulation \(EU\) No 1109/2011 of 3 November 2011 amending Annex I to Regulation \(EC\) No 2075/2005 as regards the equivalent methods for Trichinella testing \(Text with EEA relevance\)](#).

## CHAPTER III

### TRICHINOSCOPIC EXAMINATION

1. Apparatus
  - (a) An incandescent-lamp trichinoscope with 30 to 40 times and 80 to 100 times magnification or a stereomicroscope with a substage transmitted light source of adjustable intensity
  - (b) A compressorium being a pressure glass consisting of two glass plates (one of which is divided into equal fields)
  - (c) Small curved scissors
  - (d) Small forceps
  - (e) A knife for cutting specimens
  - (f) Small numbered containers for storing the specimens separately
  - (g) A dropping pipette
  - (h) A glass of acetic acid and a glass of potassium hydroxide solution for brightening any calcifications and softening dried meat.
2. Collecting of specimens

In the case of whole carcasses, several hazelnut-size samples are taken from each animal:

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- (a) in domestic swine, such samples are taken from both diaphragm pillars at the transition of the sinewy part;
- (b) in wild boar samples are taken from both diaphragm pillars at the transition of the sinewy part and in addition from the jaw, the muscles of the lower leg, the intercostal muscles and the tongue muscles, giving a total of six samples from each individual animal;
- (c) if certain muscles are not available for sampling, a total of four samples are taken from the muscles that are available;
- (d) in pieces of meat, four hazelnut-size samples of striated muscle tissue containing if possible no fat, taken from different points, are taken from each piece, where possible close to bones or tendons.

### 3. Procedure

- (a) In general a compressorium is filled with  $1,0 \pm 0,1$  g of meat, normally corresponding with 28 oat-kernel-size pieces. If necessary, two compressoria need to be filled to examine 56 oat-kernel-size pieces.
- (b) If both diaphragm pillars are present in a domestic swine, the *Trichinella* inspector cuts 28 oat-kernel-size pieces from each of the above specimens taken from a whole carcase, making 56 pieces in all.
- (c) If only one diaphragm pillar is present, 56 pieces are cut in different places, if possible from the transition to the sinewy part.
- (d) The samples collected from the other four muscles of wild boar are each cut into seven oat-kernel-size pieces, giving a total of 28 additional pieces.
- (e) The *Trichinella* inspector then compresses the 56 (or 84) pieces between the glass plates so that normal print can be clearly read through the slide preparation.
- (f) If the flesh of the specimens to be examined is dry and old, the preparations must be softened for 10 to 20 minutes before pressing with a mixture of one part of potassium hydroxide solution to about two parts of water.
- (g) From each of the samples taken from pieces of meat, the *Trichinella* inspector cuts 14 oat-kernel-size pieces, making 56 pieces in all.
- (h) The microscopic examination must be carried out by scanning each preparation slowly and carefully at a magnification of 30 to 40 times.
- (i) If the trichinoscopic examination reveals suspect areas, they must be examined at the trichinoscope's most powerful magnification (80 to 100 times).
- (j) Where the result is uncertain, the examination is repeated on other specimens and slide preparations until the information required is obtained. The trichinoscopic examination must be carried out for at least six minutes.
- (k) The minimum time fixed for the examination does not include the time necessary for taking samples and making the preparations.
- (l) As a general rule, the trichinoscopic examiner must not inspect more than 840 pieces a day, corresponding with examinations of 15 domestic swine or 10 wild boar.

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

## ANNEX II

### Freezing treatments

#### A. Freezing method 1

- (a) Meat brought in already frozen is to be kept in this condition.
- (b) The technical equipment and energy supply of the refrigeration room must be such as to ensure that the required temperature is reached very rapidly and maintained in all parts of the room and of the meat.
- (c) Insulated packaging must be removed before freezing, except in the case of meat that is already at the required temperature throughout when it is brought into the refrigeration room or meat so packaged that the packaging will not prevent it from reaching the required temperature within the specified time.
- (d) Consignments in the refrigeration room must be kept separately and under lock and key.
- (e) The date and time when each consignment is brought into the refrigeration room must be recorded.
- (f) The temperature in the refrigeration room must be at least  $-25^{\circ}\text{C}$ . It must be measured using calibrated thermo-electric instruments and recorded continuously. It may not be measured directly in the cold air flow. The instruments must be kept under lock and key. The temperature charts must include the relevant data from the meat inspection register on import and the date and time of commencement and completion of freezing, and must be retained for one year after compilation.
- (g) Meat of a diameter or thickness of up to 25 cm must be frozen for at least 240 consecutive hours, and meat of a diameter or thickness of between 25 and 50 cm must be frozen for at least 480 consecutive hours. This freezing process must not be applied to meat that is thicker or of a larger diameter. The freezing time is calculated from the point when the temperature in the freezing room reaches that specified in (f).

#### B. Freezing method 2

The general provisions of (a) to (e) of method 1 are complied with, and the following time-temperature combinations applied:

- (a) meat of a diameter or thickness of up to 15 cm must be frozen for one of the following time-temperature combinations:
  - 20 days at  $-15^{\circ}\text{C}$ ,
  - 10 days at  $-23^{\circ}\text{C}$ ,
  - 6 days at  $-29^{\circ}\text{C}$ ;
- (b) meat of a diameter or thickness of between 15 cm and 50 cm must be frozen for one of the following time-temperature combinations:
  - 30 days at  $-15^{\circ}\text{C}$ ,
  - 20 days at  $-25^{\circ}\text{C}$ ,
  - 12 days at  $-29^{\circ}\text{C}$ .

The temperature in the refrigeration room must be no higher than the level of the selected inactivation temperature. It must be measured using calibrated thermoelectric instruments and recorded continuously. It must not be measured directly in the cold air flow. The instruments must be kept under lock and key. The temperature charts

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

must include the relevant data from the meat inspection register on importation and the date and time of commencement and completion of freezing, and must be retained for one year after compilation.

Where freezing tunnels are used and the above procedures are not followed strictly, the food business operator must be able to prove to the competent authority that the alternative method is effective in killing *Trichinella* parasites in pigmeat.

### C. Freezing method 3

Treatment consists of commercial freeze-drying or freezing of meat for specified time-temperature combinations with temperature monitored at the centre of each cut.

- (a) The general provisions of (a) to (e) of Method 1 are to be complied with for the following time-temperature combinations:
- 106 hours at  $-18^{\circ}\text{C}$ ,
  - 82 hours at  $-21^{\circ}\text{C}$ ,
  - 63 hours at  $-23,5^{\circ}\text{C}$ ,
  - 48 hours at  $-26^{\circ}\text{C}$ ,
  - 35 hours at  $-29^{\circ}\text{C}$ ,
  - 22 hours at  $-32^{\circ}\text{C}$ ,
  - 8 hours at  $-35^{\circ}\text{C}$ ,
  - 1/2 hour at  $-37^{\circ}\text{C}$ .
- (b) The temperature is to be measured using calibrated thermoelectric instruments and recorded continuously. The thermometer probe is inserted in the centre of a cut of meat no smaller in size than the thickest piece of meat to be frozen. This cut must be placed at the least favourable position in the refrigeration room, not close to the cooling equipment or directly in the cold air flow. The instruments must be kept under lock and key. The temperature charts must include the data numbers from the meat inspection register on import and the date and time of commencement and completion of freezing, and must be retained for one year after compilation.

## ANNEX III

### Examination of animals other than swine

Horse meat, wild game meat and other meat that could contain *Trichinella* parasites must be examined in accordance with one of the digestion methods specified in Chapter I or II of Annex I, with the following changes:

- (a) specimens weighing at least 10 g are taken from the lingual or jaw muscle of horses and from the foreleg, tongue or diaphragm of wild boar;
- (b) in the case of horse, where those muscles are lacking, a larger-sized specimen is to be taken from a pillar of the diaphragm at the transition to the sinewy part. The muscle must be clean of connective tissue and fat;
- (c) at least 5 g of sample is digested following the reference method of detection in Chapter I of Annex I or an equivalent method in Chapter II. For each digest, the total weight of muscle examined must not exceed 100 g in the case of the method in

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- Chapter I and methods A and B in Chapter II and 35 g in the case of method C in Chapter II;
- (d) where the result is positive, a further 50 g specimen is taken for a subsequent independent examination;
- (e) without prejudice to the rules on conservation of animal species, all meat of game animals other than wild boar, such as bears, carnivorous mammals (including marine mammals) and reptiles, are to be tested by sampling 10 g of muscle at the predilection sites or larger amounts if those sites are not available. Predilection sites are:
- (i) in bear: diaphragm, masseter muscle and tongue;
  - (ii) in walrus: tongue;
  - (iii) in crocodiles: masseter, pterygoid and intercostal muscles;
  - (iv) in birds: muscles of the head (e.g. masseter and neck muscles).
- (f) The digestion time must suffice to ensure adequate digestion of the tissue of these animals but must not exceed 60 minutes.

## [<sup>F1</sup>ANNEX IV

### CHAPTER I

#### **OFFICIAL RECOGNITION OF A HOLDINGS OR A COMPARTMENT AS APPLYING CONTROLLED HOUSING CONDITIONS**

- A. The following requirements must be met by food business operators to obtain official recognition of holdings:
- (a) the operator must have taken all practical precautions with regard to building construction and maintenance in order to prevent rodents, any other kind of mammals and carnivorous birds from having access to buildings where animals are kept;
  - (b) the operator must apply a pest-control programme, in particular for rodents, effectively to prevent infestation of pigs. The operator must keep records of the programme to the satisfaction of the competent authority;
  - (c) the operator must ensure that all feed has been obtained from a facility that produces feed in accordance with the principles described in Regulation (EC) No 183/2005 of the European Parliament and of the Council<sup>(7)</sup>;
  - (d) the operator must store feed intended for *Trichinella* susceptible species in closed silos or other containers that are impenetrable to rodents. All other feed supplies must be heat-treated or produced and stored to the satisfaction of the competent authority;
  - (e) the operator must ensure that dead animals are collected, identified and transported without undue delay in accordance with Articles 21 and 22 of Regulation (EC) No 1069/2009 of the European Parliament and of the Council<sup>(8)</sup> and with Annex VIII to Commission Regulation (EU) No 142/2011<sup>(9)</sup>;

---

*Status: Point in time view as at 01/06/2014.*

*Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)*

---

- (f) if a rubbish dump is located in the neighbourhood of the holding, the operator must inform the competent authority. Subsequently, the competent authority must assess the risks involved and decide whether the holding is to be recognised as applying controlled housing conditions;
  - (g) the operator must ensure that piglets coming onto the holding from outside and pigs purchased are born and bred under controlled housing conditions;
  - (h) the operator must ensure that pigs are identified so each animal can be traced back to the holding;
  - (i) the operator may introduce new animals onto the holding only if they come from holdings also officially recognised as applying controlled housing conditions;
  - (j) none of the animals has access to outdoor facilities unless the food business operator can show by a risk analysis to the satisfaction of the competent authority that the time period, facilities and circumstances of outdoor access do not pose a danger for introduction of *Trichinella* in the holding.
- B. Food business operators of holdings officially recognised as applying controlled housing conditions shall inform the competent authority where any of the requirements laid down in point A is no longer fulfilled or where any other change has occurred that might affect the status of the holding.
- C. The competent authorities in Member States may only recognise a holding or a category of holdings provided that they have verified that the requirements laid down in point A are met.

## CHAPTER II

### REPORTING ON *TRICHINELLA* SITUATION

- (a) The number of cases (imported and autochthonous) of *Trichinella* in humans, including epidemiological data shall be reported in accordance with Commission Decision 2000/96/EC<sup>(10)</sup>.
- (b) The number of tests and the results of testing for *Trichinella* in domestic swine, wild boar, horses, game and any other susceptible animals shall be submitted in accordance with Annex IV to Directive 2003/99/EC. Data on domestic swine shall, at least, provide specific information related to:
  - (i) tests on animals raised under controlled housing conditions;
  - (ii) tests on breeding sows, boars and fattening pigs.]

---

**Status:** Point in time view as at 01/06/2014.

**Changes to legislation:** There are currently no known outstanding effects for the  
Commission Regulation (EC) No 2075/2005 (repealed). (See end of Document for details)

---

- (1) OJ L 139, 30.4.2004, p. 206, corrected by OJ L 226, 25.6.2004, p. 83.
- (2) OJ L 139, 30.4.2004, p. 55, corrected by OJ L 226, 25.6.2004, p. 22.
- (3) OJ L 165, 30.4.2004, p. 1, corrected by OJ L 191, 28.5.2004, p. 1.
- (4) OJ L 26, 31.1.1977, p. 67.
- (5) OJ L 157, 30.4.2004, p. 33, corrected by OJ L 195, 2.6.2004, p. 12.
- (6) OJ L 325, 12.12.2003, p. 31.
- (7) [<sup>F1</sup>OJ L 35, 8.2.2005, p. 1.
- (8) OJ L 300, 14.11.2009, p. 1.
- (9) OJ L 54, 26.2.2011, p. 1.
- (10) OJ L 28, 3.2.2000, p. 50.]

---

#### **Textual Amendments**

- F1** Substituted by Commission Regulation (EU) No 216/2014 of 7 March 2014 amending Regulation (EC) No 2075/2005 laying down specific rules on official controls for *Trichinella* in meat (Text with EEA relevance).



**Status:**

Point in time view as at 01/06/2014.

**Changes to legislation:**

There are currently no known outstanding effects for the Commission Regulation (EC) No 2075/2005 (repealed).