Commission Regulation (EU) No 277/2010 of 31 March 2010 concerning the authorisation of 6-phytase as a feed additive for poultry for fattening and breeding other than turkeys for fattening, for poultry for laying and for pigs other than sows (holder of authorisation Roal Oy) (Text with EEA relevance)

COMMISSION REGULATION (EU) No 277/2010

of 31 March 2010

concerning the authorisation of 6-phytase as a feed additive for poultry for fattening and breeding other than turkeys for fattening, for poultry for laying and for pigs other than sows (holder of authorisation Roal Oy)

(Text with EEA relevance)

THE EUROPEAN COMMISSION,

Having regard to the Treaty on the Functioning of the European Union,

Having regard to Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition⁽¹⁾, and in particular Article 9(2) thereof,

Whereas:

- (1) Regulation (EC) No 1831/2003 provides for the authorisation of additives for use in animal nutrition and for the grounds and procedures for granting such authorisation.
- (2) In accordance with Article 7 of Regulation (EC) No 1831/2003, an application was submitted for the authorisation of the preparation set out in the Annex to this Regulation. That application was accompanied by the particulars and documents required under Article 7(3) of Regulation (EC) No 1831/2003.
- (3) The application concerns the authorisation of the enzyme 6-phytase (EC 3.1.3.26) produced by *Trichoderma reesei* (CBS 122001) as a feed additive for poultry for fattening and breeding other than turkeys for fattening, for poultry for laying and for pigs other than sows, to be classified in the additive category 'zootechnical additives'.
- (4) The European Food Safety Authority (the Authority) concluded in its opinion of 11 November 2009⁽²⁾ that 6-phytase (EC 3.1.3.26) produced by *Trichoderma reesei* (CBS 122001) does not have an adverse effect on animal health, human health or the environment and that the use of that preparation can improve the performance of the animals. The Authority does not consider that there is a need for specific requirements of post market monitoring. It also verified the report on the method of analysis of the feed additive in feed submitted by the Community Reference Laboratory set up by Regulation (EC) No 1831/2003.
- (5) The assessment of that preparation shows that the conditions for authorisation, as provided for in Article 5 of Regulation (EC) No 1831/2003, are satisfied. Accordingly,

the use of that preparation should be authorised, as specified in the Annex to this Regulation.

(6) The measures provided for in this Regulation are in accordance with the opinion of the Standing Committee on the Food Chain and Animal Health,

HAS ADOPTED THIS REGULATION:

Article 1

The preparation specified in the Annex, belonging to the additive category 'zootechnical additives' and to the functional group 'digestibility enhancers', is authorised as an additive in animal nutrition subject to the conditions laid down in that Annex.

Article 2

This Regulation shall enter into force on the 20th day following its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 31 March 2010.

For the Commission The President José Manuel BARROSO

ANNEX

Identifica Niom e number of the of the holder additive of author		e Composi fipaç ic chemicalor formula, catego descripti of i, analyticalnima method	age ory		content f /kg plete stuff re		End period of authorisation
Catagory of zoo		dditives Eurotics	nal group: d		ty onhana	ore	
Category of zoo 4a12 Roal Oy	6- phytase EC 3.1.3.26	and breedir other than turkeys for fattenin Poultry for laying Pigs other than sows	ve — sition: ng Preparat of g 6- phytase (EC s 3.1.3.26) produced ng by <i>Trichode</i>	250 PPU ion d 125 PPU 250 PPU		1. 2.	21.4.2020 In the directions for use of the additive and premixture, indicate the storage temperature, storage life, and stability to pelleting. Maximum recommended dose per kilogram of complete feed for all authorised species: 1

a 1 PPU is the amount of enzyme which liberates 1 μ mol of inorganic phosphate from sodium phytate per minute at pH = 5.0 and 37C °.

b Details of the analytical methods are available at the following address of the Community Reference Laboratory: www.irmm.jrc.be/crl-feed-additives

Status: Point in time view as at 31/12/2020. Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EU) No 277/2010. (See end of Document for details)

active substance: 6- phytase (EC 3.1.3.26) produced by Trichoderma reesei (CBS 122001) Analytical method.* Colorimetric method grotection, quantifying the activity of 6- phytase by3.For use in feed containing more than 0.23 % phytin- bound phosphorus.4.For safety: boreathing protection, quantifying the activity of 6- phytase by measuring released inorganic phosphate from sodium phytate by reduction of a phytase4.For safety: breathing protection, glasses and gloves shall bc used during handling.			
6- use phytase feed (EC containing produced more by 0,23 % pressei 0,23 % phytin- bound ptotoderma phytin- reesei 0,23 % phytin- bound ptotoderma phytin- by the 122001) Analytical Analytical 4. method ⁺ Colorimetric method gloves of 6- phytase and gloves shall be used during measuring released inorganic phosphate from sodium phytate by analysing the colour colour formed by reduction of a phosphomolybdate a		2	Ean
in feed (EC 3.1.3.26) produced by <i>Trichoderma</i> <i>reesei</i> (CBS 122001) Analytical (CBS 122001) Analytical (CBS 122001) Analytical (Colorimetric method. ⁴ Colorimetric method activity of 6- phytase by measuring released inorganic phosphate from sodium physhate by measuring released inorganic phosphate from sodium physhate by measuring released inorganic phosphate from sodium physhate by measuring released inorganic phosphonolybdate		3.	
pinytate (ECfeed containing more than 0,23 % phytin- bound phosphorus. <i>Trichoderma</i> reesei (CBS 122001) Analytical analytical quantifying the activity of 6- phytase by measuring released inorganic phosphore from sodium phytate by reduction of 6- phytase by measuring released inorganic phosphore from sodium phytate by reduction of 6-4.For safety: breathing protection, glasses and during handling.4.	6-		
(EC 3.1.3.26) produced by Trichoderma reesei (CBS 	phytase		
3.1.3.26) containing produced by <i>Trichoderma</i> 0,23 % reesei (CBS 122001) Analytical Analytical 4. method.* Colorimetric quantifying the activity of 6- phytase by glasses shall be used during measuring released inorganic phosphare phosphare bo from sodium phytate by analysing the colour formed by reduction of a phosphate from sodium phytate by reduction of a phosphorulybdate by			
produced by <i>Trichoderma</i> <i>reesei</i> (CBS 122001) Analytical (CBS) 122001) Analytical (Colorimetric method. ⁴ Colorimetric method quantifying the activity of 6- phytase by measuring released inorganic phosphate from sodium phytate by y analysing the colour formed by reduction of a phosphomolybdate			
by Trichoderma reesei (CBS 122001) Analytical Analytical Analytical (CBS 122001) Analytical (CBS 122001) Analytical (CBS 122001) Analytical (CBS (Colorimetric method quantifying the activity of 6- phytase by measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			
Trichoderma reesei (CBS 122001)0,23 % phytin- bound phosphorus.Analytical method.*4.Colorimetric method quantifying the activity of 6- phytase by measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phytase4.			
reesei phytin-bound (CBS 122001) Analytical 4. method:* For Colorimetric protection, glasses and activity of 6- phytase by shall be used during measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of			0,23 %
Image: Construction of the colour formed by reduction of a phosphore.Description of the colour formed by reduction of a phosphore.Image: Colour formed by reduction of the colour formed by reduction of a phosphore.Image: Colour formed by reduction of a phosphore.			phytin-
122001) Analytical 4. For safety: Colorimetric method.* Safety: breathing Colorimetric and glasses and quantifying the and glasses the activity of shall be 0f by gloves shall be used during measuring released inorganic phosphate handling. released inorganic phosphate inorganic handling. phosphate from sodium phytate by handling. phosphate from sodium phytate by handling. by analysing the colour formed by indianality phosphomolybdate by reduction indianality indianality indianality			
Analytical Analytical Analytical Analytical Analytical Analytical Analytical Analytical Analytical Analytical 4. For safety: breathing protection, glasses and gloves shall be used during handling. 4. For safety: breathing protection, glasses and gloves shall be used during handling.			phosphorus.
method: Colorimetric method quantifying the activity of 6- phytase by measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			
Colorimetricbreathingmethodprotection,quantifyingglassestheandactivityglovesofshall6-bephytaseusedbyduringmeasuringreleasedinorganicphosphatefromsodiumphytatebybyanalysingthecolourformedbyaphosphomolybdate	Analytical	4.	
method protection, quantifying glasses the and activity gloves of shall 6- be phytase used by measuring released inorganic phosphate from from sodium phytae by analysing the colour formed by reduction inorganic phosphate from sodium phytae by analysing the colour formed by analysing the colour formed by a phosphomolybdate			
quantifying glasses activity gloves of shall 6- be phytase used by during measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate l			breathing
quantifying the activity of 6- phytase by measuring released inorganic phosphate from sodium phytate by analysing the colour formed by analysing the colour formed by analysing the colour formed by analysing the colour formed by analysing the colour formed by analysing the colour formed by ceduction of a phosphomolybdateglasses and glasses and gloves shall be used during handling.			
the and activity of 6- be phytase used by during measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate late			
activity of 6- phytase by measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			
of 6- shall phytase used by measuring released inorganic phosphate from from sodium phytate by by analysing the colour formed by value of a phosphomolybdate	activity		
6- phytase be by measuring released inorganic phosphate from sodium phytate by by analysing the colour formed by reduction of a phosphomolybdate in in	of		
phytase by measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate	6-		
by measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate	phytase		
measuring released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			
released inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			
inorganic phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			nununng.
phosphate from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate	inorganic		
from sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			
sodium phytate by analysing the colour formed by reduction of a phosphomolybdate			
phytate by analysing the colour formed by reduction of a phosphomolybdate			
by analysing the colour formed by reduction of a phosphomolybdate			
analysing the colour formed by reduction of a phosphomolybdate			
the colour formed by reduction of a phosphomolybdate			
colour formed by reduction of a phosphomolybdate			
formed by reduction of a phosphomolybdate			
by reduction of a phosphomolybdate			
reduction of a phosphomolybdate			
of a phosphomolybdate			
a phosphomolybdate			
phosphomolybdate			
complex.			
	complex.		

a 1 PPU is the amount of enzyme which liberates 1 μ mol of inorganic phosphate from sodium phytate per minute at pH = 5.0 and 37C °.

b Details of the analytical methods are available at the following address of the Community Reference Laboratory: www.irmm.jrc.be/crl-feed-additives

Status: Point in time view as at 31/12/2020. Changes to legislation: There are currently no known outstanding effects for the Commission Regulation (EU) No 277/2010. (See end of Document for details)

- (1) OJ L 268, 18.10.2003, p. 29.
- (2) *The EFSA Journal* 2009; 7(11):1380.

Status:

Point in time view as at 31/12/2020.

Changes to legislation:

There are currently no known outstanding effects for the Commission Regulation (EU) No 277/2010.