Commission Implementing Regulation (EU) No 1119/2012 of 29 November 2012 concerning the authorisation of preparations of Pediococcus acidilactici CNCM MA 18/5M DSM 11673, Pediococcus pentosaceus DSM 23376, NCIMB 12455 and NCIMB 30168, Lactobacillus plantarum DSM 3676 and DSM 3677 and Lactobacillus buchneri DSM 13573 as feed additives for all animal species (Text with EEA relevance)

COMMISSION IMPLEMENTING REGULATION (EU) No 1119/2012

of 29 November 2012

concerning the authorisation of preparations of *Pediococcus acidilactici* CNCM MA 18/5M DSM 11673, *Pediococcus pentosaceus* DSM 23376, NCIMB 12455 and NCIMB 30168, *Lactobacillus plantarum* DSM 3676 and DSM 3677 and *Lactobacillus buchneri* DSM 13573 as feed additives for all animal species

(Text with EEA relevance)

THE EUROPEAN COMMISSION,

Having regard to the Treaty on the Functioning of the European Union,

Having regard to Regulation (EC) No 1831/2003 of the European Parliament and of the Council of 22 September 2003 on additives for use in animal nutrition⁽¹⁾, and in particular Article 9(2) thereof,

Whereas:

- (1) Regulation (EC) No 1831/2003 provides for the authorisation of additives for use in animal nutrition and for the grounds and procedures for granting such authorisation. Article 10(7) of Regulation (EC) No 1831/2003 in conjunction with Article 10(1) to (4) thereof sets out specific provisions for the evaluation of products used in the Union as silage additives at the date that Regulation became applicable.
- In accordance with Article 10(1) of Regulation (EC) No 1831/2003, preparations of *Pediococcus acidilactici* CNCM MA 18/5M DSM 11673, *Pediococcus pentosaceus* DSM 23376, *Pediococcus pentosaceus* NCIMB 12455, *Pediococcus pentosaceus* NCIMB 30168, *Lactobacillus plantarum* DSM 3676, *Lactobacillus plantarum* DSM 3677 and *Lactobacillus buchneri* DSM 13573 were entered in the Community Register of Feed Additives as existing products belonging to the functional group of silage additives, for all animal species.
- (3) In accordance with Article 10(2) of Regulation (EC) No 1831/2003 in conjunction with Article 7 thereof, applications were submitted for the authorisation of those preparations as feed additives for all animal species, requesting those additives to be classified in the category 'technological additives' and in the functional group 'silage additives'. Those applications were accompanied by the particulars and documents required under Article 7(3) of Regulation (EC) No 1831/2003.

- The European Food Safety Authority ('the Authority') concluded in its opinions of (4)23 May 2012⁽²⁾ and 14 June 2012⁽³⁾ that, under the proposed conditions of use, the preparations concerned do not have an adverse effect on animal health, human health or the environment. The Authority also concluded that the preparations of *Pediococcus* acidilactici CNCM MA 18/5M DSM 11673, Pediococcus pentosaceus DSM 23376, Pediococcus pentosaceus NCIMB 12455, Pediococcus pentosaceus NCIMB 30168 concerned have the potential to improve the production of silage from all forages by reducing the pH and increasing the preservation of dry matter and/or protein. It also concluded that the preparations of Lactobacillus plantarum DSM 3676 and Lactobacillus plantarum DSM 3677 concerned have the potential to improve the production of silage from easy and moderately difficult to ensile material by increasing the lactic acid content and the preservation of dry matter, by reducing the pH and moderately the loss of protein. It also concluded that the preparation of *Lactobacillus* buchneri DSM 13573 concerned has the potential to increase acetic acid concentration for a wide range of forages. The Authority does not consider that there is a need for specific requirements of post-market monitoring. It also verified the report on the method of analysis of the feed additives in feed submitted by the Community Reference Laboratory set up by Regulation (EC) No 1831/2003.
- (5) The assessment of the preparations concerned shows that the conditions for authorisation, as provided for in Article 5 of Regulation (EC) No 1831/2003, are satisfied. Accordingly, the use of those preparations should be authorised as specified in the Annex to this Regulation.
- (6) Since safety considerations do not require the immediate application of the modifications to the conditions of authorisation, it is appropriate to allow a transitional period for interested parties to prepare themselves to meet the new requirements resulting from the authorisation.
- (7) The measures provided for in this Regulation are in accordance with the opinion of the Standing Committee on the Food Chain and Animal Health,

HAS ADOPTED THIS REGULATION:

Article 1

Authorisation

The preparations specified in the Annex belonging to the additive category 'technological additives' and to the functional group 'silage additives', are authorised as additives in animal nutrition, subject to the conditions laid down in that Annex.

Article 2

Transitional measures

The preparations specified in the Annex and feed containing them, which are produced and labelled before 20 June 2013 in accordance with the rules applicable before 20

Status: This is the original version (as it was originally adopted).

December 2012 may continue to be placed on the market and used until the existing stocks are exhausted.

Article 3

Entry into force

This Regulation shall enter into force on the twentieth day following that of its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 29 November 2012.

For the Commission

The President

José Manuel BARROSO

Status: This is the original version (as it was originally adopted).

ANNEX

Identifi	ca Niom e	Additiv	e Compo	si Sipa çies	Maxim	umMinim	ııMaxim	u 10 ther	End
numbei	of the		chemica		age	content	content	provisio	nsf
of the	holder		formula	a, categor		CFU/kg	of	•	period
additive	e of		descrip	ti of i,		fresh m			of
	authori	sation		aanimal					authorisation
			method						
Categor	y of techr	ological	additives.	Function	al group:	silage ado	ditives		
1k2104	_	Pedioco	ccus	Add ditive	_				20 December
		acidilaci	ici	animadsi	tion			1.	December
		CNCM		Prepiers at					194992
		MA		of					directions
		18/5M		Pedioco	ccus				for
		DSM		acidilaci					use
		11673		CNCM					of
		11075		MA					the
				18/5M					additive
				DSM					and
				11673					premixture,
				containii	າຕ				indicate
				a	15				the
				minimur	_				storage
				of	11				temperature
				$\frac{01}{3}$ ×					and
									storage
				10 ⁹					life.
				CFU/					
				g				2.	Minimum
				additive					dose
				Characte	erisation				of
				of					the
				the					additive
				active					when
				substanc	e				used
				Pedioco	ccus				without
				acidilaci					combination
				CNCM					with
				MA					other
				18/5M					micro-
				DSM					organisms
				11673					as
				Analytic	al				silage
				method ^a					additives:
				Enumera					3 ×
				in	111011				
				the					10 ⁷
				feed					CFU/
									Kg
				additive:					

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

		spread plate method (EN 15786) Identific pulsed field gel electropl (PFGE)			3.	fresh material. For safety: it is recommended to use breathing protection and gloves during handling.
1k2105	Pediococ pentosac DSM 23376	Additive animadsic species of Pediococopentosac DSM 23376 containing a minimum of 1 × 10 ¹¹ CFU/g additive Character of the active substance Pediococopentosac DSM 23376 Analytic method Enumerating the feed	tion tion ccus ecus ng m erisation eeccus eeus		 2. 	December 2022 directions for use of the additive and premixture, indicate the storage temperature and storage life. Minimum dose of the additive when used without combination with other microorganisms as

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

Document Generated: 2023-12-24

Status: This is the original version (as it was originally adopted).

		additive: spread plate method (EN 15786) Identification: pulsed field gel electrophoresis (PFGE)	silage additives: 1 × 10 ⁸ CFU/ Kg fresh material. 3. For safety: it is recomment to use breathing protection and gloves during handling.	nded
1k2106 —	Pediococcus pentosaceus NCIMB 12455	Additive — animalsition sprepasation of Pediococcus pentosaceus NCIMB 12455 containing a minimum of 3 × 109 CFU/ g additive Characterisation of the active substance Pediococcus pentosaceus NCIMB 12455	1. 20 December 2022 directions for use of the additive and premixtur indicate the storage temperatu and storage life. 2. Minimum dose of the additive when used without	e, re

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

		Analytical method ^a Enumeration in the feed additive: spread plate method (EN 15786) Identification: pulsed field gel electrophoresis (PFGE)	3.	combination with other micro- organisms as silage additives: 3×10^7 CFU/ Kg fresh material. For safety: it is recommended to use breathing protection and gloves during handling.
1k2107 —	Pediococcus pentosaceus NCIMB 30168	Additive animalsition Preparation of Pediococcus pentosaceus NCIMB 30168 containing a minimum of 5 × 10 ¹⁰ CFU/ g additive Characterisation of the	2.	20 December 22 directions for use of the additive and premixture, indicate the storage temperature and storage life. Minimum dose

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

		active substance Pediococcus pentosaceus NCIMB 30168 Analytical methoda Enumeration in the feed additive: spread plate method (EN 15786) Identification: pulsed field gel electrophoresis (PFGE)	3.	of the additive when used without combination with other microorganisms as silage additives: 1 × 10 ⁸ CFU/ Kg fresh material. For safety: it is recommended to use breathing protection and gloves during handling.
1k20731 —	Lactobacillus plantarum DSM 3676	Additive — animal sition species at on of Lactobacillus plantarum DSM 3676 containing a minimum of 6 × 10 ¹¹ le at the following address of the Reference animal sition and si		December 20 Decemb

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

	i I		ODI!				1
			CFU/				and
			g				storage
			additive				life.
			Characte	erisation		2	Minimo
			of			2.	Minimum
			the				dose
			active				of
			substanc	e			the
			Lactoba				additive
			plantaru				when
			DSM				used
			3676				without
			Analytic	al			combination
			method ^a	ai			with
			Enumera	4:			other
				HIOH			micro-
			in				organisms
			the				as
			feed				silage
			additive:				additives:
			spread				1 ×
			plate				10^{8}
			method				CFU/
			(EN				
			15787)				Kg
			Identific	ation:			fresh
			pulsed				material.
			field			3.	The
			gel			٥.	additive
			electropl	noresis			shall
			(PFGE)				be
			,				used
							in
							easy
							and
							moderately
							difficult
							to
							ensile
							material ^b .
						4	Г
						4.	For
							safety:
							it
							is
							recommended
							to
							use
							breathing

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

Document Generated: 2023-12-24

Status: This is the original version (as it was originally adopted).

						protection and gloves during handling.
1k20732	Lactobac plantaru DSM 3677	Additive animadsi spepiarati of Lactobal plantaru DSM 3677 containin a minimur of 4 × 10 ¹¹ CFU/g additive Characte of the active substance Lactobal plantaru DSM 3677 Analytic method Enumera in the feed additive spread plate method (EN 15787) Identific pulsed field gel	tion ion cillus m ng n erisation ce cillus al		 2. 3. 	20 December 222 directions for use of the additive and premixture, indicate the storage temperature and storage life. Minimum dose of the additive when used without combination with other microorganisms as silage additives: 1 × 108 CFU/ Kg fresh material. The additive and premixture, indicate the additive when a storage life.

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

		electrophoresis (PFGE)	4.	shall be used in easy and moderately difficult to ensile material ^b . For safety: it is recommended to use breathing protection and gloves during handling.
1k20733 —	Lactobacillus buchneri DSM 13573	Additive — animadsition Speciasation of Lactobacillus buchneri DSM 13573 containing a minimum of 2 × 10 ¹¹ CFU/ g additive Characterisation of the active substance	2.	December 2022 directions for use of the additive and premixture, indicate the storage temperature and storage life. Minimum dose of the additive

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

	Lactobacillus buchneri DSM 13573 Analytical methoda Enumeration in the feed additive: spread plate method (EN 15787) Identification: pulsed field gel electrophoresis (PFGE)	3.	when used without combination with other micro-organisms as silage additives: 1 × 108 CFU/ Kg fresh material. For safety: it is recommended to use breathing protection and gloves during handling.
--	--	----	---

a Details of the analytical methods are available at the following address of the Reference Laboratory: http://irmm.jrc.ec.europa.eu/EURLs/EURL_feed_additives/Pages/index.aspx

b Easy to ensile forage: > 3 % soluble carbohydrates in fresh material. Moderately difficult to ensile forages: 1,5-3,0 % soluble carbohydrate in fresh material. Commission Regulation (EC) No 429/2008 (OJ L 133, 22.5.2008, p. 1).

- (1) OJ L 268, 18.10.2003, p. 29.
- (2) EFSA Journal 2012; 10(6):2733.
- (3) EFSA Journal 2012; 10(7):2780.