Commission Implementing Regulation (EU) 2017/2470 of 20 December 2017 establishing the Union list of novel foods in accordance with Regulation (EU) 2015/2283 of the European Parliament and of the Council on novel foods (Text with EEA relevance)

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)



^{F2}... LIST OF NOVEL FOODS

Textu	al Amendments
F1	Substituted by Commission Implementing Regulation (EU) 2018/1023 of 23 July 2018 correcting Implementing Regulation (EU) 2017/2470 establishing the Union list of novel foods (Text with EEA relevance).
F2	Word in Annex heading omitted (31.12.2020) by virtue of The Novel Food (Amendment) (EU Exit) Regulations 2019 (S.I. 2019/702), regs. 1, 64(a) ; 2020 c. 1, Sch. 5 para. 1(1)
Conte	nt of the list
1.	The ^{F3} list shall consist of Tables 1 and 2.
	 Textual Amendments F3 Word in Annex para. 1 omitted (31.12.2020) by virtue of The Novel Food (Amendment) (EU Exit) Regulations 2019 (S.I. 2019/702), regs. 1, 64(b); 2020 c. 1, Sch. 5 para. 1(1)
2.	Table 1 includes the authorised novel foods and contains the following information: U.K.
Colum Colum	
Colum Colum	n 3 : Additional specific labelling requirements
3.	Table 2 includes the specifications on novel foods and contains the following information: U.K.
Colum Colum	

TABLE 1: AUTHORISED NOVEL FOODS

Authorised novel food	Conditions u the novel foo used		Additional specific labelling requirements	Other requirements	[^{F8} Data Protection]
N - Acetyl-D- neuraminic acid	Specified food category Infant and follow-on formulae as defined by Regulation (EU) No 609/2013 ^a	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' <i>N</i> -acetyl-D-		

Processed cereal-based foods and baby foods for infants and young children as defined by Regulation (EU) No 609/2013 Foods for special medical purposes for infants and young children as defined by Regulation (EU) No 609/2013	0,05 g/kg for solid foods	neuraminic acid' Food supplements containing <i>N</i> -acetyl-D- neuraminic acid shall bear a statement that the food supplement should not be given to infants, young children and children under 10 years of age where they consume breast milk or other foods with added <i>N</i> -acetyl-D- neuraminic acid within the same twenty four hour period.	
Total diet replacement foods for weight control as defined by Regulation (EU) No 609/2013	0,2 g/L (drinks) 1,7 g/kg (bars)		
Foods bearing statements on the absence or reduced presence of gluten in accordance with the	1,25 g/kg		

requirements of Commission Implementing Regulation (EU) No 828/2014 ^b		
Unflavoured pasteurised and sterilised (including UHT) milk-based products	0,05 g/L	1
Unflavoured fermented milk-based products, heat treated after fermentation, flavoured fermented milk products including heat-treated products	0,05 g/L (beverages) 0,4 g/kg (solids)	1
Dairy analogues, including beverage whiteners	0,05 g/L (beverages) 0,25 g/kg (solids)	
Cereal bars	0,5 g/kg	
Table top sweeteners	8,3 g/kg	
Fruit and vegetable- based drinks	0,05 g/L	
Flavoured drinks	0,05 g/L	
Speciality coffee, tea, herbal and fruit infusions, chicory; tea, herbal and fruit infusions and chicory	0,2 g/kg	

	extracts; tea, plant, fruit and cereal preparations for infusions Food Supplements as defined in Directive 2002/46/EC ^c	300 mg/day for general population older than 10 years 55 mg/day for infants 130 mg/day for young children 250 mg/day for children between 3 to 10 years of age		
<i>Adansonia digitata</i> (Baobab) dried fruit pulp	Not specified		The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Baobab fruit pulp '	
<i>Ajuga</i> <i>reptans</i> extract from cell cultures	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels		
L-Alanyl-L- Glutamine	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels		

	Foods for special medical purposes as defined in Regulation (EU) No 609/2013 excluding foods for infants and young children			
	Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen			
Algal oil from the	Specified food	Maximum levels of	The designation	
microalgae	category	DHA	of the novel	
<i>Ulkenia</i> sp.	Bakery products (breads, rolls and sweet biscuits)	200 mg/100 g	food on the labelling of the foodstuffs containing it shall be 'Oil from the	
	Cereal bars	500 mg/100 g	micro-algae	
	Non- alcoholic beverages (including milk based beverages)	60 mg/100 ml	Ulkenia sp. '	
[^{F9} Allanblacki seed oil	a Specified food category	Maximum levels	The designation of the novel	
	Yellow fat spreads and cream based spreads	30 g/100 g	food on the labelling of the foodstuffs containing it shall be '	
	Mixtures of vegetable oils (*) and milk (falling under	30 g/100 g	<i>Allanblackia</i> seed oil'	

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27. June 2024. There are changes that may be brought into force at a future date

known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	and o oils a in Pa Anne Regu	pt olive oils live pomace s defined rt VIII of x VII of lation (EU) 308/2013.]		
<i>Aloe macroclada</i> Baker leaf	Specified food category	Maximum levels		
extract	Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of the similar gel derived <i>from Aloe</i> <i>vera</i> (L.) Burm.		
Antarctic Krill oil from <i>Euphausia</i> superba	Specified food category	Maximum levels of combined DHA and EPA	The designation of the novel food on the labelling of	
	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g	the foodstuffs containing it shall be 'Lipid extract	
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g	from the crustacean Antarctic Krill (<i>Euphausia</i> <i>superba</i>)'	
	Non- alcoholic beverages Milk-based drinks Dairy analogue drinks	80 mg/100 ml		

Spreadable fat and dressings	600 mg/100 g
Cooking fats	360 mg/100 ml
Breakfast cereals	500 mg/100 g
Bakery products (breads, rolls and sweet biscuits)	200 mg/100 g
Nutrition bars/cereal bars	500 mg/100 g
Food Supplements as defined in Directive 2002/46/EC	3 000 mg/ day for the general population 450 mg/day for pregnant and lactating women
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal
Processed cereal-based food and baby food	200 mg/100 ml

	intended for infants and young children covered by Regulation (EU) No 609/2013			
	Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen			
	Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014			
Antarctic Krill oil rich in phospholipids from	Specified food category	Maximum levels of combined DHA and EPA	The designation of the novel food on the labelling of	
Euphausia superba	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g	the foodstuffs containing it shall be 'Lipid extract	
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g	from the crustacean Antarctic Krill (<i>Euphausia</i> <i>superba</i>)'	
	Non- alcoholic beverages	80 mg/100 ml		

Milk-based drinks Dairy analogue drinks	
Spreadable fat and dressings	600 mg/100 g
Cooking fats	360 mg/100 ml
Breakfast cereals	500 mg/100 g
Bakery products (breads, rolls and sweet biscuits)	200 mg/100 g
Nutrition bars/cereal bars	500 mg/100 g
Food Supplements as defined in Directive 2002/46/EC	3 000 mg/ day for the general population 450 mg/day for pregnant and lactating women
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements	250 mg/meal

	for weight control Processed cereal-based food and baby food	200 mg/100 ml	
	intended for infants and young children covered by Regulation (EU) No 609/2013		
	Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen		
	Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014		
Arachidonic acid-rich oil from	Specified food category	Maximum levels	The designation of the novel
the fungus Mortierella alpina	Infant formula and follow-on formula as defined in Regulation (EU) No 609/2013	In accordance with Regulation (EU) No 609/2013	food on the labelling of the foodstuffs containing it shall be 'Oil from <i>Mortierella</i> <i>alpina</i> ' or ' <i>Mortierella</i> <i>alpina</i> oil'

Argan oil from	Foods for special medical purposes for premature infants as defined in Regulation (EU) No 609/2013 Specified food	In accordance with Regulation (EU) No 609/2013 Maximum levels	The designation	
Argania	category		of the novel	
spinosa	As seasonings	Not specified	food on the labelling of	
	Food Supplements as defined in Directive 2002/46/EC	In line with normal food use of vegetable oils	the foodstuffs containing it shall be ' Argan oil ' and if used as seasoning ' Vegetable oil only for seasoning ' shall be mentioned on the label	
Astaxanthin- rich oleoresin	Specified food	Maximum levels	The designation	
	Category		ot the novel	
from	<i>category</i> Food	40-80 mg/day	of the novel food on the	
from <i>Haematococcu</i>	Food Supplements	of oleoresin,	food on the labelling of	
from Haematococcu pluvialis	Food Supplements as defined	of oleoresin, resulting	food on the labelling of the foodstuffs	
from <i>Haematococcu</i>	Food Supplements	of oleoresin,	food on the labelling of	
from Haematococcu pluvialis	Food Supplements as defined in Directive	of oleoresin, resulting in ≤ 8 mg astaxanthin	food on the labelling of the foodstuffs containing it shall be '	
from Haematococcu pluvialis algae Basil seeds (Ocimum	Food Supplements as defined in Directive 2002/46/EC Specified food	of oleoresin, resulting in ≤ 8 mg astaxanthin per day <i>Maximum</i>	food on the labelling of the foodstuffs containing it shall be '	
from Haematococcu pluvialis algae Basil seeds (Ocimum	Food Supplements as defined in Directive 2002/46/EC Specified food category Fruit juice and fruit/ vegetable blend	of oleoresin, resulting in ≤ 8 mg astaxanthin per day Maximum levels 3 g/200 ml for addition of whole basil seeds (Ocimum	food on the labelling of the foodstuffs containing it shall be '	Authorised on 22 August 2019. This

isotonic and energy drinks intended for sportsmen Protein and cereal bars intended for sportsmen	500 mg/100 g	the foodstuffs containing it shall be 'betaine'. The labelling of foods containing betaine shall bear	proprietary scientific evidence and scientific data protected in accordance with Article 26 of Regulation
Meal replacements intended for sportsmen	20 mg/100 g	a statement that the foods should not be used if food	(EU) 2015/2283. Applicant: DuPont
Total diet replacement for weight control as defined under Regulation (EU) No 609/2013 Foods for Special Medical Purposes as defined under Regulation (EU) No 609/2013 for adults	500 mg/100 g (bar) 136 mg/100 g (soup) 188 mg/100 g (porridge) 60 mg/100 g (beverages) 400 mg/day]	supplements containing betaine are consumed the same day.	Nutrition Biosciences ApS, Langebrogade 1 Copenhagen K, DK-1411, Denmark. During the period of data protection, the novel food betaine is authorised for placing on the market within the Union only by DuPont Nutrition Biosciences ApS unless a subsequent applicant obtainsauthorisation for the novel food without reference to the proprietary scientific evidence or scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283

					or with the agreement of DuPont Nutrition Biosciences ApS, End date of the data protection: 22 August 2024.
Fermented black bean extract	Specified food category	Maximum levels	The designation of the novel		
	Food Supplements as defined in Directive 2002/46/EC	4,5 g/day	food on the labelling of the foodstuffs containing it shall be ' Fermented black bean (Soya) extract ' ' or ' Fermented Soya extract '		
Bovine lactoferrin	Specified food	Maximum levels	The designation		
	<i>category</i> Infant formula and follow-on formula as defined in Regulation (EU) No 609/2013 (ready to drink)	100 mg/100 ml	of the novel food on the labelling of the foodstuffs containing it shall be ' Lactoferrin from cows ' milk'		
	Foods on dairy basis intended for young children (ready to eat/ drink)	200 mg/100 g			
	Processed cereal food (solid)	670 mg/100 g			
	Foods for special medical	Depending on the needs of			

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27. June 2024. There are changes that may be brought into force at a future date

known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	purposes as defined in Regulation (EU) No 609/2013	the individual up to 3 g/day			
	Beverages based on milk	200 mg/100 g			
	Powdered drink mixes based on milk (ready to drink)	330 mg/100 g			
	Beverages based on fermented milk (including yoghurt drinks)	50 mg/100 g			
drinks Product based of	alcoholic	120 mg/100 g			
	Products based on yoghurt	80 mg/100 g			
	Products based on cheese	2 000 mg/100 g			
	Ice cream	130 mg/100 g			
	Cakes and pastries	1 000 mg/100 g			
	Candies	750 mg/100 g			
	Chewing gum	3 000 mg/100 g			
[^{F10} Bovine milk basic whey protein	Specified food category	Maximum levels	The designation of the novel food on the		Authorised on 20 November 2018 Thia
isolate	Infant formulae as defined in Regulation (EU) No 609/2013 Follow-on formulae as defined in	30 mg/100 g (powder) 3,9 mg/100 mL (reconstituted) 30 mg/100 g (powder)	lood on the labelling of the foodstuffs containing it shall be 'Milk whey protein isolate'. Food supplements containing		2018 . This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article

Regulation	4,2 mg/100	bovine milk		26 of
(EU) No	mL	basic whey		Regulation
609/2013	(reconstituted)	protein isolate		(EU)
	· · · · · · · · · · · · · · · · · · ·	*		
Total diet	300 mg/day	shall bear the		2015/2283.
replacement	30 mg/100	following		Applicant:
foods for	g (powder	statement:		Armor
weight	formula for	'This food		Protéines
control as	infants during	supplement		S.A.S., 19
defined by	the first	should not		bis, rue de la
Regulation	months of	be consumed		Libération
(EU) No	life until the	by infants/		35460 Saint-
609/2013	introduction	children/		Brice-en-
Foods for	of appropriate	adolescents		Coglès,
special	complementary			France.
medical	feeding)	of one/three/		During the
purposes as	3,9 mg/100	eighteen (*)		period of data
defined in	mL	years'		protection the
Regulation	(reconstituted	(*)		novel food
	formula for			bovine milk
(EU) No		Depending on		
609/2013	infants during	the age group		basic whey
Food	the first	the food		protein isolate
Supplements	months of	supplement is		is authorised
as defined	life until the	intended for.		for placing
in Directive	introduction			on the market
2002/46/EC	of appropriate			within the
	complementary	7		Union only
	feeding)			by Armor
	30 mg/100			Protéines
	g (powder			S.A.S. unless
	formula for			a subsequent
	infants when			applicant
	appropriate			obtains
	complementary	7		authorisation
	feeding is	1		for the novel
	introduced)			food without
	4,2 mg/100			reference
	4,2 mg/100 mL			
				to the
	(reconstituted			proprietary
	formula for			scientific
	infants when			evidence or
	appropriate			scientific data
	complementary			protected in
	feeding is			accordance
	introduced)			with Article
	58 mg/day			26 of
	for young			Regulation
	children			(EU)
	380 mg/			2015/2283
	day for			or with the
	children and			agreement
	adolescents			
	from 3 to 18			
	years of age			
	years of age	I	l	

Status: Point in time view as at 31/12/2020.	
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes	
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.	
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)	

		610 mg/day for adults 25 mg/day for infants 58 mg/day for young children 250 mg/ day for children and adolescents from 3 to 18 years of age 610 mg/day for adults]			of Armor Protéines S.A.S. End date of the data protection: 20 November 2023 .
Buglossoides arvensis seed oil	Specified food category	Maximum levels of stearidonic acid (STA)	The designation of the novel food on the		
	Dairy	250 mg/100 g	labelling of		
	products and analogues	75 mg/100 g for drinks	the foodstuffs containing it shall be		
	Cheese and cheese products	r and fat and nulsions ling ds (not oking ing			
	Butter and other fat and oil emulsions including spreads (not for cooking or frying purposes)				
	Breakfast cereals 625 mg/100 g	625 mg/100 g			
	Food supplements as defined in Directive 2002/46/EC, excluding food supplements for infants and young children	500 mg/day			
	Foods for special medical	In accordance with the particular			

	purposes as defined in Regulation (EU) No 609/2013, excluding foods for special medical purposes intended for infants and young children	nutritional requirements of the persons for whom the products are intended		
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal		
<i>Calanus</i> <i>finmarchicus</i> oil	Specified food category	Maximum levels	The designation of the novel	
	Food supplements as defined	2,3 g/day	food on the labelling of	
	in Directive 2002/46/EC		the foodstuffs containing it shall be 'oil from <i>Calanus</i> <i>finmarchicus</i> (crustacean)'	
Chewing gum base	2002/46/EC Specified food	Maximum levels	containing it shall be 'oil from <i>Calanus</i> <i>finmarchicus</i> (crustacean)' The designation	
gum base	2002/46/EC Specified		containing it shall be 'oil from <i>Calanus</i> <i>finmarchicus</i> (crustacean)' The	

			homopolymer, maleated, esters with polyethylene glycol mono- Me ether) ' or ' Gum base (including CAS No: 1246080-53-4)	
Chewing gum base (Methyl	Specified food category	Maximum levels	The designation of the novel	
vinyl ether- maleic anhydride copolymer)	Chewing gum	2 %	food on the labelling of the foodstuffs containing it shall be ' Gum base (including methyl vinyl ether-maleic anhydride copolymer) ' or ' Gum base (including CAS No 9011-16-9) '	
Chia oil from <i>Salvia</i> <i>hispanica</i>	Specified food category	Maximum levels	The designation of the novel	
F	Fats and oils	10 %	food on the labelling of the foodstuffs containing it shall be 'Chia oil (<i>Salvia</i> <i>hispanica</i>)'	
	Pure chia oil	2 g/day		
	Food Supplements as defined in Directive 2002/46/EC	2 g/day		
[^{F11} Chia seeds (<i>Salvia</i>	Specified food category	Maximum levels	The designation of the novel	
hispanica)	Bread products	5 % (whole or ground chia seeds)	food on the labelling of the foodstuffs	
	Baked products	10 % whole chia seeds	containing it shall be 'Chia seeds	
	Breakfast cereals	10 % whole chia seeds	(Salvia hispanica)'	

Sterilised	5 % whole	
ready to eat meals based on cereal grains, pseudocereals grains and/or pulses	chia seeds	
Fruit, nut and seed mixes		
Pre-packaged Chia seed as such		
Confectionery (including chocolate and chocolate products), excluding chewing gums		
Dairy products (including yoghurt) and analogues		
Edible ices		
Fruit and vegetables products (including fruit spreads, compotes with/without cereals, fruit- preparations to underlay or to be mixed with dairy products, fruit desserts, mixed fruits with coconut milk for a twin pot)		
Non- alcoholic beverages (including		

	fruit juice and fruit/ vegetable blend beverages) Puddings that do not require heat treatment at or above 120 °C in their manufacture, processing or preparation]			
Chitin- glucan from <i>Aspergillus</i> <i>niger</i>	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels 5 g/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chitin- glucan from <i>Aspergillus</i> <i>niger</i> '	
Chitin- glucan complex from <i>Fomes</i> <i>fomentarius</i>	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels 5 g/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chitin- glucan from <i>Fomes</i> <i>fomentarius</i> '	
Chitosan extract from fungi (<i>Agaricus</i> <i>bisporus</i> ; <i>Aspergillus</i> <i>niger</i>)	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chitosan extract from <i>Agaricus</i> <i>bisporus</i> ' or 'Chitosan	

			extract from Aspergillus niger '		
Chondroitin sulphate	Specified food category	Maximum levels	The designation of the novel		
	Food supplements as defined in Directive 2002/46/ EC for adult population, excluding pregnant and lactating women	1 200 mg/day			
Chromium Picolinate	food categorylevels of totaldesignation	The designation of the novel food on the			
	Foods covered by Regulation (EU) No 609/2013	250 μg/day	labelling of the foodstuffs containing it shall be ' Chromium Picolinate '		
	Foods fortified in accordance with Regulation (EC) No 1925/2006 ^d				
<i>Cistus</i> <i>incanus</i> L. Pandalis	Specified food category	Maximum levels	The designation of the novel		
herb	Herbal infusions	Intended daily intake: 3 g herbs/day (2 cups/day)	ed food on the ntake: labelling of bs/day the foodstuffs		
Citicoline	Specified food category	Maximum levels	1. The design of the	nation	

	Food Supplements as defined in Directive 2002/46/EC Foods for special medical purposes as defined in Regulation (EU) No 609/2013	500 mg/day 250 mg per serving and a maximum daily consumption level of 1 000 mg	novel food on the labelling of the foodstuffs containing it shall be 'Citicoline' 2. The labelling of foods containing citicoline shall bear a statement that the product is not intended to be consumed by children
Clostridium butyricum	Specified food category	Maximum levels	The designation of the novel
	Food Supplements as defined in Directive 2002/46/EC	1,35 × 10 ⁸ CFU/day	food on the labelling of the foodstuffs containing it shall be ' <i>Clostridium</i> <i>butyricum</i> MIYAIRI 588 (CBM 588)' or ' <i>Clostridium</i> <i>butyricum</i> (CBM 588)'

[^{F8} D-ribose	Specified food category	Maximum levels	The designation of the novel		Authorised on 16 April 2019. This		
	Cereal bars	0,20 g/100 g	food on the		inclusion		
	Fine bakery wares	0,31 g/100 g	labelling of the foodstuffs containing it		is based on proprietary scientific		
	Chocolate confectionery (excluding chocolate bars)	0,17 g/100 g	shall be 'D- ribose'. The labelling of foods containing D-ribose		evidence and scientific data protected in accordance with Article 26 of		
	Milk- based drinks (excluding malts and shakes)	0,08 g/100 g	D-ribose shall bear a statement that the foods should not be used if food supplements containing D-ribose are consumed the same day.		Regulation (EU) 2015/2283. Applicant: Bioenergy Life Science,		
	Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen, isotonic and energy drinks	0,80 g/100 g			Inc., 13840 Johnson St. NE, Minneapolis, Minnesota, 55304, USA. During the period of data protection, the novel food D-ribose is authorised for placing on the market within the Union only by Bioenergy Life Science, Inc. unless a subsequent		
	Bars intended to meet the expenditure of intense muscular effort especially for sportsmen	3,3 g/100 g					
	Meal replacement for weight control (as drinks)'	0,13 g/100 g					applicant obtains authorisation for the novel food without reference
	Meal replacement for weight control (as bars)	3,30 g/100 g			to the proprietary scientific evidence or scientific data		
	Confectionery	0,20 g/100 g			protected in accordance with Article		

	Tea and infusions (in powder form to be reconstituted)	0,23 g/100 g]		26 of Regulation (EU) 2015/2283 or with the agreement of Bioenergy Life Science, Inc. End date of the data protection: 16 April 2024 (5 years).
Extract of defatted cocoa	Specified food category	Maximum levels	Consumers shall be instructed not	
powder	Nutrition bars	1 g/day and	to consume	
	Milk based beverages	300 mg polyphenols corresponding	more than 600 mg polyphenols	
	Any other foods (including food supplements as defined in Directive 2002/46/ EC) which have become established vehicles for functional ingredients and which are typically positioned for consumption by health conscious adults	to not more than 550 mg of extract of defatted cocoa powder in one portion of food (or food supplement)	corresponding to 1,1 g of extract of defatted cocoa powder per day	
Low fat cocoa extract		Maximum levels	Consumers shall be	
	<i>category</i>	730 mg nor	instructed not to consume	
	Foods including food supplements as defined	730 mg per serving and around 1,2 g/ day	more than 600 mg of cocoa	

Coriander seed oil from Coriandrum sativum	in Directive 2002/46/EC Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels 600 mg/day	flavanols per day The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Coriander seed oil '	
[^{F12} Cranberry extract powder	Specified food category Food Supplements as defined in Directive 2002/46/EC for the adult population	Maximum levels 350 mg/day]	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' cranberry extract powder '	Authorised on 20 November 2018. This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283. Applicant: Ocean Spray Cranberries Inc. One Ocean Spray Drive Lakeville- Middleboro, MA, 02349, USA. During the period of data protection the novel food, cranberry extract powder, is authorised for placing on the market within the Union only by Ocean Spray Cranberries

	Snecified	Maximum	The	Inc. unless a subsequent applicant obtains authorisation for the novel food without reference to the proprietary scientific evidence or scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283 or with the agreement of Ocean Spray Cranberries Inc. End date of the data protection: 20 November 2023.
<i>Crataegus pinnatifida</i> dried fruit	Specified food category	Naximum levels	The designation of the novel	
	Herbal infusions	In line with normal	l labelling of the foodstuffs containing	
	Jams and jellies in accordance with Directive 2001/113/EC e	food use of Crataegus laevigata		
	Compotes			
α- cyclodextrin	Not specified		The designation of the novel food on the labelling of the foodstuffs containing	

γ- cyclodextrin	Not specified		it shall be ' Alpha- cyclodextrin ' or ' α - cyclodextrin ' The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Gamma- Cyclodextrin ' or ' γ - Cyclodextrin '	
[^{F13} Decorticate grains of <i>Digitaria</i> <i>exilis</i> (Kippist) Stapf (Traditional food from a third country)	dNot specified		The designation of the novel food on the labelling of the foodstuffs containing it shall be 'decorticated fonio (<i>Digitaria</i> <i>exilis</i>) grains']	
Dextran preparation produced by <i>Leuconostoc</i> mesenteroides	Specified food category Bakery products	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Dextran '	
Diacylglycero oil of plant origin	Specified food categoryCooking oilsFat spreadsSalad dressingsMayonnaise	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Diacylglycerol oil of plant origin (at	

	Meal replacement for weight control (as drinks) Bakery products Yoghurt type products		least 80 diacylgly			
Dihydrocapsi (DHC)	atSpecified food category	Maximum levels	1.	The desig	nation	
	Cereal bars	9 mg/100 g		the		
Biscuits, cookies and crackers Rice based snacks Carbonated drinks, dilutable drinks, fruit juice based beverages	cookies and	9 mg/100 g		novel food on the labell of the foods conta it shall be 'Dihy	ing tuffs	
		12 mg/100 g				
	drinks, dilutable drinks, fruit juice based	1,5 mg/100 ml				
	Vegetable drinks	2 mg/100 ml	2.	Food suppl conta	ements	
	Coffee based drinks, tea based drinks	1,5 mg/100 ml		synth dihyd will		
	Flavoured water — still	1 mg/100 ml		be labell	nded	
	Precooked oatmeal cereal	2,5 mg/100 g		as 'not intene for		
	Other cereals	4,5 mg/100 g		childı up	ren	
	Ice cream, dairy desserts	4 mg/100 g		to 4.5	,	
	Pudding mixes (ready to eat)	2 mg/100 g		years		
	Products based on yoghurt	2 mg/100 g				

	Chocolate confectionery Hard candy Sugar-free	7,5 mg/100 g 27 mg/100 g 115 mg/100 g		
	gum Whitener/	40 mg/100 g		
	creamer Sweeteners	200 mg/100 g		
	Soup (ready to eat)	1,1 mg/100 g		
	Salad dressing	16 mg/100 g		
	Vegetable protein	5 mg/100 g		
	Ready to eat meals	3 mg/meal		
	Meal replacements for weight control	3 mg/meal		
	Meal replacement for weight control (as drinks)	1 mg/100 ml		
	Food Supplements as defined in Directive 2002/46/EC	3 mg/single intake 9 mg/day		
	Non- alcoholic powdered drink mixes	14,5 mg/kg equivalent to 1,5 mg/100 ml		
[^{F14} Dried aerial parts of <i>Hoodia</i>	Specified food category	Maximum levels	The designation of the novel	Authorised on 3 September 2018. This
parviflora	Food Supplements as defined in Directive 2002/46/ EC for adult population	9,4 mg/day]	food on the labelling of the foodstuffs containing it shall be 'dried aerial parts of <i>Hoodia</i> parviflora '	inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

26 of Regulation (EU) 2015/2283. Applicant: Desert Labs, Ltd Kibbutz Yotvata. 88820 Israel. During the period of data protection the novel food dried aerial parts of Hoodia parviflora is authorised for placing on the market within the Union only by Desert Labs, Ltd unless a subsequent applicant obtains authorisation for the novel food without reference to the proprietary scientific evidence or scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283 or with the agreement of Desert Labs, Ltd. End date of the data protection: 3 September 2023.

Dried extract of <i>Lippia</i> <i>citriodora</i> from cell cultures	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels In line with normal use in food supplements of a similar extract from the leaves of <i>Lippia</i> <i>citriodora</i>	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'dried extract of <i>Lippia</i> <i>citriodora</i> from cell cultures HTN [®] Vb'	
<i>Echinacea</i> <i>angustifolia</i> extract from cell cultures	Specified food category	Maximum levels	-	
	Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of a similar extract from the root of <i>Echinacea</i> <i>angustifolia</i>		
[^{F15} Echinacea purpurea extract from	Specified food category	Maximum levels	The designation of the novel	
extract from cell cultures	Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of a similar extract from florets within the flower head of <i>Echinacea</i> <i>purpurea</i>]	food on the labelling of the foodstuffs containing it shall be 'dried extract of <i>Echinacea</i> <i>purpurea</i> from cell cultures EchiPure- PCTM'	
<i>Echium plantagineum</i> oil	Specified food category	Maximum levels of stearidonic acid (STA)	The designation of the novel food on the	
	Milk-based products and drinkable yoghurt products	250 mg/100 g; 75 mg/100 g for drinks	labelling of the foodstuffs containing it shall be ' Refined echium oil '	

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	delivered in a single dose Cheese preparations Spreadable fat and dressings Breakfast cereals Food supplements as defined in Directive 2002/46/EC	750 mg/100 g 750 mg/100 g 625 mg/100 g 500 mg/day		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal		
[^{F16} Ecklonia cava phlorotannins	Specified food category Food supplements as defined in Directive 2002/46/EC intended for the general population, excluding children	Maximum levels 163 mg/ day for adolescents from 12 to 14 years of age 230 mg/ day for adolescents above 14 years of age	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' <i>Ecklonia</i> <i>cava</i> Phlorotannins' Food supplements containing	

under the age of 12 years	263 mg/day for adults]	<i>Ecklonia</i> phlorotan	inins
		shall bear	
		following statement	
			This
			food
			supplement
			should
			not
			be
			consumed
			by
			children/
			adolescents
			under
			the
			age
			of
			twelve/
			fourteen/
			eighteen (*)
			years.
			This
			food
			supplement
			should
			not
			be
			consumed
			by
			persons
			with
			thyroid disease
			or
			by
			persons
			who
			are
			aware
			of
			or
			have
			been
			identified
			as
			being
			at
			risk
			of
			developing

			 shoul not be consult of the consult of	se. lement ld umed lements lining e umed. mding	
[^{F17} Egg membrane	Specified food	Maximum levels	The designation		Authorised on 25 November
hydrolysate	<i>category</i> Food Supplements as defined in Directive 2002/46/ EC intended for the general adult population	450 mg/day]	of the novel food on the labelling of the foodstuffs containing it shall be ' egg membrane hydrolysate '.		2018. This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283. Applicant: Biova, LLC., 5800 Merle Hay Rd, Suite 14 PO Box 394 Johnston 50131,

				Iowa USA. During the period of data protection the novel food egg membrane hydrolysate is authorised for placing on the market within the Union only by Biova, LLC. unless a subsequent applicant obtains authorisation for the novel food without reference to the proprietary scientific evidence or scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283 or with the agreement of Biova, LLC. End date of the data protection: 25 November 2023
Epigallocatech gallate as a purified extract from green tea leaves (<i>Camellia</i> <i>sinensis</i>)	Food food category Foods including food supplements as defined in Directive 2002/46/EC	Maximum levels	The labelling shall bear a statement that consumers should not consume more than 300 mg of extract per day	

[^{F18} L- ergothioneine	Specified food category Alcohol-free beverages Milk-based drinks ' Fresh ' milk	<i>Maximum</i> <i>levels</i> 0,025 g/kg 0,025 g/kg 0,040 g/kg	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' L- ergothioneine	
	products(*) Cereal bars Chocolate	0,2 g/kg 0,25 g/kg	-	
	confectionery Food supplements as defined in Directive 2002/46/EC	30 mg/day for general population (excluding pregnant and lactating women) 20 mg/day for children older than 3 years		
	milk ergot not re or in	n used in products L- hioneine may eplace in whole part, any milk ituent]		
[^{F16} Extract of three herbal		Maximum levels	The designation	
roots (<i>Cynanchum</i> <i>wilfordii</i> Hemsley, <i>Phlomis</i> <i>umbrosa</i> Turcz. and <i>Angelica</i> <i>gigas</i> Nakai)	Food supplements as defined in Directive 2002/46/ EC for adult population	175 mg/day]	of the novel food on the labelling of the foodstuffs containing it shall be 'extract of three herbal roots (<i>Cynanchum</i> <i>wilfordii</i> Hemsley, <i>Phlomis</i> <i>umbrosa</i> Turcz. and <i>Angelica</i> <i>gigas</i> Nakai)'.	

			The labelling of food supplements containing the extract of mixture of the three herbal roots shall bear a statement in close proximity to the list of ingredients indicating that it should not be consumed by individuals with known celery allergy.		
Ferric Sodium EDTA	Specified food category Food supplements as defined in Directive 2002/46/EC	Maximum levels (expressed as anhydrous EDTA) 18 mg/day for children 75 mg/day for adults	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Ferric Sodium EDTA '		
	Foods covered by Regulation (EU) No 609/2013	12 mg/100 g			
	Foods fortified in accordance with Regulation (EC) No 1925/2006				
Ferrous ammonium phosphate	Specified food category	Maximum levels	The designation of the novel	1	
	Food supplements as defined	To be used in compliance with	food on the labelling of the foodstuffs containing		

	in Directive 2002/46/EC Foods covered by Regulation (EU) No 609/2013 Foods fortified in accordance with Regulation (EC) No 1925/2006	Directive 2002/46/EC, Regulation (EU) No 609/2013 and/or Regulation (EC) No 1925/2006	it shall be ' Ferrous ammonium phosphate '		
Fish peptides from <i>Sardinops</i>	Specified food category	Maximum levels fish peptide product	The designation of the novel food on the		
sagax	Foods based on yoghurt, yoghurt drinks, fermented milk products, and powdered milk	0,48 g/100 g (ready to eat/ drink)	labelling of the foodstuffs containing it shall be 'Fish (<i>Sardinops</i> <i>sagax</i>) peptides'		
	Flavoured water, and vegetable- based drinks	0,3 g/100 g (ready to drink)			
	Breakfast cereals	2 g/100 g	-		
	Soups, stews and soup powders	0,3 g/100 g (ready to eat)	-		
Flavonoids from Glycyrrhiza glabra	Specified food category	Maximum levels of flavonoids from Glycyrrhiza glabra	of the	Beverages natitaining flavonoids shall be presented to the final	
	Beverages based on milk	120 mg/day	on the	consumer as single	
	Beverages based on yoghurt		fabell of the foods conta		

Beverages based on fruit or vegetables			it sh be 'F
Food Supplements as defined in Directive 2002/46/EC	120 mg/day	2.	fro Gl gl L. Tl
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	120 mg/day		lal of th fo w th pr w ad
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	120 mg/day		as no fo in she a stath (a
			(b

1 vonoids n cyrrhiza bra lling ls re duct eđ el d edient 1 ement the product should not be consumed by pregnant and breast feeding women, children and young adolescents and people taking prescription drugs should only consume the

Status: Point in time view as at 31/12/2020.

			3.	(c) The amou of flavor in the final food shall be indica on the labell of the food conta it.	noids nted ing
[^{F19} Fruit pulp, pulp juice, concentrated pulp juice from <i>Theobroma</i> <i>cacao</i> L. (Traditional food from a third country)	Not specified	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp', 'cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp juice' or 'cocoa (<i>Theobroma</i>			

	Succi Cad	<i>cacao</i> L.) concentrated pulp juice' depending on the form used.]			
Fucoidan extract from	Specified food	Maximum levels	The designation		
the seaweed	category		of the novel		
Fucus vesiculosus	Foods including food supplements as defined in Directive 2002/46/ EC for the general population	250 mg/day	food on the labelling of the foodstuffs containing it shall be 'Fucoidan extract from seaweed <i>Fucus</i> <i>vesiculosus</i> '.		
Fucoidan extract from the seaweed	Specified food category	Maximum levels	The designation of the novel		
Undaria pinnatifida	Foods including food supplements as defined in Directive 2002/46/ EC for the general population	250 mg/day	food on the labelling of the foodstuffs containing it shall be ' Fucoidan extract from seaweed Undaria pinnatifida '		
2'- Fucosyllactoso	Specified food category	Maximum levels	1. The desig of	nation	
	Unflavoured pasteurised and sterilised (including UHT) milk-based products	1,2 g/l	the novel food on the labell of	ing	
	Unflavoured fermented milk-based products	1,2 g/l beverages 19,2 g/kg products other than beverages	the foods conta it shall be '2'- fucos		

Flavoured fermented milk-based products including heat-treated products Dairy analogues, including beverage whiteners	1,2 g/l beverages 19,2 g/kg products other than beverages 1,2 g/l beverages 12 g/kg for products other than	2.	The labelling of food supplements containing 2'- fucosyllactose shall bear a statement that
	beverages 400 g/kg for whitener		the supplements should not
Cereal bars	12 g/kg		be used
Table-top sweeteners	200 g/kg		if other
Infant formula as defined in Regulation (EU) No 609/2013	1,2 g/l alone or in combination with up to 0,6 g/l of lacto- N - neotetraose at a ratio of 2:1 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	3.	foods with added 2'- fucosyllactose are consumed the same day. The labelling of food supplements containing 2'- fucosyllactose
Follow-on formula as defined in Regulation (EU) No 609/2013	1,2 g/l alone or in combination with up to 0,6 g/l of lacto- N - neotetraose at a ratio of 2:1 in the final product ready for use, marketed as such or reconstituted		intended for young children shall bear a statement that the supplements should not

Processed cereal-based food and baby food for infants and young children as defined in Regulation (EU) No 609/2013	as instructed by the manufacturer 12 g/kg for products other than beverages 1,2 g/l for liquid food ready for use, marketed as such or reconstituted as instructed by the manufacturer	be used if breast milk or other foods with added 2'- fucosyllactose are consumed the same day.
Milk-based drinks and similar products intended for young children	1,2 g/l for milk-based drinks and similar products added alone or in combination with up to 0,6 g/l lacto- N -neotetraose, at a ratio of 2:1 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended	
Total diet replacement for weight control as	4,8 g/l for drinks 40 g/kg for bars	

Status: Point in time view as at 31/12/2020.

defined in Regulation (EU) No 609/2013		
Bread and pasta products bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014	60 g/kg	
Flavoured drinks	1,2 g/l	
Coffee, tea (excluding black tea), herbal and fruit infusions, chicory; tea, herbal and fruit infusions and chicory extracts; tea, plant, fruit and cereal preparations for infusions, as well as mixes and instant mixes of these products	9,6 g/l — the maximum level refers to the products ready to use	
Food supplements as defined in Directive	3,0 g/day for general population	
2002/46/EC, excluding food	1,2 g/day for young children	

	supplements for infants				
[^{F20} 2'- Fucosyllactos Difucosyllact	Specified Se ^f lood Oscategory	Maximum levels	The designation of the novel		Authorised on 19.12.2019. This inclusion
mixture (' 2'-FL/DFL ') (microbial source)	Unflavoured pasteurised and unflavoured sterilised (including UHT) milk products	2,0 g/L	food on the labelling of the foodstuffs containing it shall be '2'- Fucosyllactose/ Difucosyllactos mixture'. The labelling		is based on proprietary scientific evidence and scientific data protected in accordance with Article 26 of
	Unflavoured fermented milk-based products	2,0 g/L (beverages) 20 g/kg (products other than beverages)	of food supplements containing the 2'- Fucosyllactose Difucosyllactos mixture		Regulation (EU) 2015/2283. Applicant: Glycom A/ S, Kogle Allé 4, DK-2970
	Flavoured fermented milk-based products including heat-treated products	2,0 g/L (beverages) 20 g/kg (products other than beverages)	shall bear a statement that they should not be used if breast milk or other foods containing		Hørsholm, Denmark. During the period of data protection, the novel food 2'-
	Beverages (flavoured drinks)	2,0 g/L	added 2'- Fucosyllactose and/or		Fucosyllactose/ Difucosyllactose mixture is
	Cereal bars	20 g/kg	Difucosyllactos	se	authorised for placing
	Infant formula as defined under Regulation (EU) No 609/2013	1,6 g/L in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	the same day.		on the market within the Union only by Glycom A/S, unless a subsequent applicant obtains authorisation for the novel
	Follow-on formula as defined under Regulation (EU) No 609/2013	1,2 g/L in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer			for the novel food without reference to the proprietary scientific evidence or scientific data protected in accordance

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes

known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Processed cereal-based food and baby food for infants and young children as defined under Regulation (EU) No 609/2013	1,2 g/L (beverages) in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer 10 g/kg for products other than beverages		with 26 of Regu (EU) 2015 or wi agree Glyc End o of the prote 19.12
	Total diet replacement foods for weight control as defined under Regulation (EU) No 609/2013	4,0 g/L (beverages) 40 g/kg (products other than beverages)		
	Food for special medical purposes as defined under Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Food Supplements as defined in Directive 2002/46/EC intended for the general population excluding infants	4,0 g/day]		
Galacto- oligosaccharic	Specified food category	Maximum levels (expressed as ratio kg galacto- oligosacchari	de/	

Status: Point in time view as at 31/12/2020.

I	ka fu -l
	kg final food)
Food Supplements as defined in Directive 2002/46/EC	0,333
Milk	0,02
Milk drinks	0,03
Meal replacement for weight control (as drinks)	0,02
Dairy analogue drinks	0,02
Yoghurt	0,033
Dairy based deserts	0,043
Frozen dairy deserts	0,043
Fruit drinks and energy drinks	0,021
Infant meal replacement drinks	0,012
Baby juice	0,025
Baby yogurt drink	0,024
Baby desert	0,027
Baby snack	0,143
Baby cereals	0,027
Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen	0,013
Juice	0,021

	Fruit pie fillings	0,059
	Fruit preparations	0,125
	Bars	0,125
	Cereals	0,125
	Infant formula and follow-on formula as defined in Regulation (EU) No 609/2013	0,008
Glucosamine HCl	Specified food category	Maximum levels
	Food Supplements as defined in Directive 2002/46/EC	In line with normal food use of glucosamine from shell
	Foods covered by Regulation (EU) No 609/2013	fish
	Meal replacement for weight control	
	Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen	
	Foods bearing statements on the absence or reduced presence of gluten in	

	accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014					
Glucosamine sulphate KCl	Specified food category	Maximum levels				
	Food Supplements as defined in Directive 2002/46/EC	In line with normal food use of glucosamine from shell fish				
Glucosamine sulphate NaCl	Specified food category	Maximum levels				
	Food Supplements as defined in Directive 2002/46/EC	In line with normal food use of glucosamine from shell fish				
		11511				
Guar Gum	Specified food category	Maximum levels	1.	The design	nation	
Guar Gum		Maximum	1.	desig of the novel food on the labell of the foods	ing tuffs	
Guar Gum	<i>food</i> <i>category</i> Fresh dairy products such as yogurts, fermented milks, fresh cheeses and other dairy-based	Maximum levels	2.	design of the novel food on the labell of the	ing tuffs ining	

based compotes		risks of
Cereals accompanied by a dairy product, in packaging containing two compartments	10 g/100 g in the cereals None in the accompanying dairy product 1 g/100 g in the product when ready to eat	digestive discomfort linked to the exposure of children aged under 8 to guar gum must be visible on the label of any foodstuffs containing it. For example, 'Excessive consumption of these products may cause digestive discomfort, especially for children under 8 years of age'. 3. In the case of
		products with

			conta dairy and cerea produ respe the instru for use must clearl speci the need to mix the cerea and the dairy produ consu in order to take into accou the poten risk of gastro intest	I nets ctively, netions y fy I netbefore imption, int tial	
Heat- treated milk products	Specified food category	Maximum levels			
fermented with Bacteroides xylanisolvens	Fermented milk products (in liquid, semi-liquid and spray- dried powder forms)				
Hydroxytyros	oSpecified food category	Maximum levels	The designation of the novel		

	Fish and vegetable oils, (except	0,215 g/kg	food on the labelling of the food
	olive oils and olive pomace oils as defined in Part VIII of Annex VII of Regulation (EU) No 1308/2013 ^f), placed as		products containing it shall be 'hydroxytyrosol'. The labelling of the food products containing hydroxytyrosol shall bear the
	such on the market		following statements: (a) This
	Spreadable fats as defined in Part VII of Annex VII of Regulation (EU) No 1308/2013, placed as such on the market	0,175 g/kg	food product should not be consumed by children under the age of three years, pregnant women, and lactating women; (b) This food product should not be used for cooking, baking or frying'
Ice Structuring Protein type	Specified food category	Maximum levels	The designation of the novel
III HPLČ 12	Edible ices	0,01 %	food on the labelling of

Aqueous extracts of dried leaves of <i>Ilex guayusa</i>	Specified food category Herbal infusions Food Supplements as defined in Directive 2002/46/EC	Maximum levels In line with normal use in herbal infusions and food supplements of a similar aqueous extract of dried leaves of <i>Ilex</i> paraguariensis	the foodstuffs containing it shall be ' Ice Structuring Protein ' The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Extracts of dried leaves of <i>Ilex</i> <i>guayusa</i> '		
[^{F21} Infusion from coffee leaves of <i>Coffea</i> <i>arabica</i> L. and/ or <i>Coffea</i> <i>canephora</i> Pierre ex A. Froehner (Traditional food from a third country)	Specified food category Herbal infusions]	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Infusion from coffee leaves of <i>Coffea</i> <i>arabica</i> and/ or <i>Coffea</i> <i>canephora</i> '.		
Isomalto- oligosaccharic	Specified efood category Energy- Reduced Soft Drinks Energy Drinks Foods intended to meet the	Maximum levels 6,5 % 5,0 % 6,5 %	of the novel food on the labell of the foods	tuffs	
	expenditure of intense muscular efforts,		conta it shall	ining	

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes

known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	especially for sportsmen (including isotonic drinks) Fruit Juices Processed Vegetables and Vegetable Juices Other Soft Drinks Cereals Bars Cookies, Biscuits Breakfast Cereal Bars Hard Candies Soft Candies/ Chocolate Bars	5 % 5 % 5 % 10 % 20 % 25 % 97 % 25 %	2.	be 'Isoma Foods contai the novel ingred must be labelle as 'a source of glucos	ning lient ed	ride'.
	Meal replacement for weight control (as bars or milk based)	20 %				
Isomaltulose	Not specified		1.	The design of the novel food on the labelli of the foodst contai it shall be 'Isoma The design of	ng uffs ning altulose'.	

	Specified	Marian	the novel food on the labelling shall be accompanied by indication that the 'Isomaltulose is a source of glucose and fructose'.	
[^{F22} Lactitol	Specified food	Maximum levels	The designation	
	<i>category</i> Food Supplements as defined in Directive 2002/46/EC (capsules, tablets or powder) intended for the adult population	20 g/day]	of the novel food on the labelling of the food supplements containing it shall be ' Lactitol '	
Lacto- N - neotetraose	Specified food category	Maximum levels	1. The designation of	
	Unflavoured pasteurised and sterilised (including UHT) milk-based products0,6 g/lflUnflavoured fermented0,6 g/l for beverages cff	the novel food on the labelling of the foodstuffs containing		
	milk-based products	9,6 g/kg for products other than beverages	it shall be 'lacto-	

Flavoured fermented milk-based products including heat-treated products Dairy analogues, including	0,6 g/l for beverages 9,6 g/kg for products other than beverages 0,6 g/l for beverages 6 g/kg for	N - neotetraose'. 2. The labelling of food supplements containing lacto- N - neotetraose
beverage whiteners	products other than beverages 200 g/kg for whitener	shall bear a statement that the
Cereal bars	6 g/kg	supplements
Table-top sweeteners	100 g/kg	should not
Infant formula as defined in Regulation (EU) No 609/2013	0,6 g/l in combination with up to 1,2 g/l of 2'- fucosyllactose at a ratio of 1:2 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	be used if other foods with added lacto- N - neotetraose are consumed the same day. 3. The
Follow-on formula as defined in Regulation (EU) No 609/2013	0,6 g/l in combination with up to 1,2 g/l of 2'- fucosyllactose at a ratio of 1:2 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	13. The labelling of food supplements containing lacto- N - neotetraose intended for young children shall bear a statement that

Processed cereal-based food and baby food for infants and young children as defined in Regulation (EU) No 609/2013	6 g/kg for products other than beverages 0,6 g/l for liquid food ready for use, marketed as such or reconstituted as instructed by the manufacturer	the supplements should not be used if breast milk or other foods with added
Milk-based drinks and similar products intended for young children	0,6 g/l for milk-based drinks and similar products added alone or in combination with 2'- fucosyllactose, at a ratio of 1:2 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	added lacto- N - neotetraose are consumed the same day.
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended	
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	2,4 g/l for drinks 20 g/kg for bars	

Status: Point in time view as at 31/12/2020.

Bread and pasta products bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014	30 g/kg
Flavoured drinks	0,6 g/l
Coffee, tea (excluding black tea), herbal and fruit infusions, chicory; tea, herbal and fruit infusions and chicory extracts; tea, plant, fruit and cereal preparations for infusions, as well as mixes and instant mixes of these products	4,8 g/l — the maximum level refers to the products ready to use
Food supplements as defined in Directive 2002/46/EC, excluding food supplements for infants	1,5 g/day for general population 0,6 g/day for young children

гF23т	Specified	Manimaria	The	Authorized
[^{F23} Lacto- N-tetraose	Specified food category	Maximum levels	The designation of the novel	Authorised on 23.4.2020.
(microbial source) pa an ur sta (in U) fe m pr in fe m pr in he pr Ba (ff	Unflavoured pasteurised and unflavoured sterilised (including UHT) milk products	1,0 g/l	food on the labelling of the foodstuffs containing it shall be 'lacto- <i>N</i> - tetraose'. The labelling	This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article
	Unflavoured fermented milk-based products	1,0 g/l (beverages) 10 g/kg (products other than beverages)	of food supplements containing lacto- <i>N</i> - tetraose shall bear a statement that	26 of Regulation (EU) 2015/2283. Applicant: Glycom A/
	Flavoured fermented milk-based products including heat-treated products	1,0 g/l (beverages) 10 g/kg (products other than beverages)	they should not be used if breast milk or other foods containing added lacto- N- tetraose	S, Kogle Allé 4, DK-2970 Hørsholm, Denmark. During the period of data protection, the novel
	Beverages (flavoured drinks)	1,0 g/l	are consumed the same day.	food lacto- N- tetraose is authorised
	Cereal bars	10 g/kg		for placing on the market
	Infant formula as defined under Regulation (EU) No 609/2013	0,8 g/l in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	LU b A a a o o a a f c f c f c f c f c f c f c f c f c	within the Union only by Glycom A/S, unless a subsequent applicant obtains authorisation for the novel food without
	Follow-on formula as defined under Regulation (EU) No 609/2013	0,6 g/l in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer		to the proprietary scientific evidence or scientific data protected in accordance with Article
	Processed cereal- based food,	0,6 g/l (beverages) in the final		26 of Regulation (EU) 2015/2283

baby food for infants and young children as defined under Regulation (EU) No 609/2013	product ready for use, marketed as such or reconstituted as instructed by the manufacturer 5 g/kg for products other than beverages
Milk based drinks and similar products intended for young children	0,6 g/l (beverages) in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer 5 g/kg for products other than beverages
Total diet replacement foods for weight control as defined under Regulation (EU) No 609/2013	2,0 g/l (beverages) 20 g/kg (products other than beverages)
Food for special medical purposes as defined under Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended
Food Supplements as defined in Directive 2002/46/EC,	2,0 g/day for young children, children,

	excluding infants	adolescents, and adults]		
<i>I^{F24}Lonicera</i> <i>caerulea</i> L. berries (haskap) (Traditional food from a third country)	Not specified		The designation of the novel food on the labelling of the foodstuffs containing it shall be 'haskap (<i>Lonicera</i> <i>caerulea</i>) berries']	
Lucerne leaf extract from <i>Medicago</i>	Specified food category	Maximum levels	The designation of the novel	
sativa	Food supplements as defined in Directive 2002/46/EC	10 g/day	food on the labelling of the foodstuffs containing it shall be 'Lucerne (<i>Medicago</i> <i>sativa</i>) protein' or 'Alfalfa (<i>Medicago</i> <i>sativa</i>) protein'.	
Lycopene	Specified food category	Maximum levels	The designation of the novel	
	Fruit/ vegetable juice-based drinks (including concentrates)	2,5 mg/100 g	food on the labelling of the foodstuffs containing it shall be ' Lycopene '	
	Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen	2,5 mg/100 g		
	Total diet replacement	8 mg/meal		

Status: Point in time view as at 31/12/2020.

	for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control			
	Breakfast cereals	5 mg/100 g		
	Fats and dressings	10 mg/100 g		
	Soups other than tomato soups	1 mg/100 g		
	Bread (including crispy breads)	3 mg/100 g		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Food supplements as defined in Directive 2002/46/EC	15 mg/day		
Lycopene from <i>Blakeslea</i>	Specified food category	Maximum levels	The designation of the novel	
trispora	Fruit/ vegetable juice-based drinks (including concentrates)	2,5 mg/100 g	food on the labelling of the foodstuffs containing it shall be ' Lycopene '	
	Drinks intended to meet the expenditure of intense	2,5 mg/100 g		

	muscular effort especially for sportsmen Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight	8 mg/meal		
	control Breakfast cereals	5 mg/100 g		
	Fats and dressings	10 mg/100 g	-	
	Soups other than tomato soups	1 mg/100 g		
	Bread (including crispy breads)	3 mg/100 g		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Food supplements as defined in Directive 2002/46/EC	15 mg/day		
Lycopene from tomatoes	Specified food category	Maximum levels	The designation of the novel	
	Fruit/ vegetable juice-based drinks	2,5 mg/100 g	food on the labelling of the foodstuffs containing	

(including concentrates)		it shall be ' Lycopene '	
Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen	2,5 mg/100 g		
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	8 mg/meal		
Breakfast cereals	5 mg/100 g		
Fats and dressings	10 mg/100 g		
Soups other than tomato soups	1 mg/100 g		
Bread (including crispy breads)	3 mg/100 g		
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
Food supplements as defined in Directive 2002/46/EC	15 mg/day		

Lycopene oleoresin from tomatoes	Specified food category Fruit/ vegetable juice-based drinks (including concentrates) Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen	Maximum levels of lycopene 2,5 mg/100 g 2,5 mg/100 g	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Lycopene oleoresin from tomatoes '	
	Total diet replacement for weight control covered by Regulation (EU) No 609/2013 and meal replacements for weight control	8 mg/meal		
	Breakfast cereals Fats and	5 mg/100 g 10 mg/100 g		
	dressings Soups other than tomato soups Bread (including	1 mg/100 g 3 mg/100 g		
	crispy breads) Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the		

		products are intended		
[^{F16} Hen egg white	Specified food category	Maximum levels	The designation	
lysozyme hydrolysate	Food supplements as defined in Directive 2002/46/ EC intended for adult population	1000 mg/ day]	of the novel food on the labelling of food supplements containing it shall be ' Hen egg white lysozyme hydrolysate '	
Magnesium citrate malate	Specified food category	Maximum Ievels	The designation of the novel	
	Food Supplements as defined in Directive 2002/46/EC		food on the labelling of the foodstuffs containing it shall be ' Magnesium citrate malate	
Magnolia Bark Extract	Specified food category	Maximum levels	The designation of the novel	
	Mints (confectionary products)	0,2 % for breath freshening	food on the labelling of the foodstuffs	
	Chewing gum	purposes. Based on a 0,2 % maximum incorporation level and a maximum gum/mint size of 1,5 g each, each gum or mint serving will contain no more than 3 mg of magnolia	containing it shall be ' Magnolia Bark Extract '	
		bark extract. <i>Maximum</i>		

unsaponifiable matter Methylcellulos	Supplements as defined in Directive 2002/46/EC Chewing gum	2 g/day 2 % Maximum levels 2 %	food on the labelling of the foodstuffs containing it shall be ' Maize-germ oil extract ' The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Methylcellulos	Methylcellulos is not to be used in foods specially prepared for young children e	e
[^{F25} 1- Methylnicotin chloride	Specified affitte category Food Supplements as defined in Directive 2002/46/EC for the adult population excluding pregnant and lactating women	Maximum levels 58 mg/day]	The designation of the novel food on the labelling of the foodstuffs containing it shall be '1- Methylnicotina chloride'. Food supplements containing 1- Methylnicotina shall bear the following statement: This food supplement should be consumed by		Authorised on 2 September 2018. This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283. Applicant: Pharmena SA, Wolczanska 178, 90 530

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes

known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

			adults only excluding pregnant and lactating women	Lodz, Poland. During the period of data protection thenovel food 1- methylnicotinamide chloride is authorised for placing on the market within the Union only by Pharmena S.A. unless a subsequent applicant obtains authorisation for the novel food without reference to the proprietary scientific evidence or scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283 or with the agreement of Pharmena S.A. End date of the data protection: 2 September 2023
(6S)-5- methyltetrahy acid, glucosamine salt	Specified d f%ff ic category	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' (6S)-5-	

	Food Supplements as defined in Directive 2002/46/EC as a source of folate		methyltetrahyd acid, glucosamine salt ' or ' 5MTHF- glucosamine '	rofolic	
Monomethyls (Organic Silicon)	il SRacif ied food category	Maximum levels of silicon	The designation of the novel		
	Food Supplements as defined in Directive 2002/46/ EC for adult population (in liquid form)	10,40 mg/day	food on the labelling of the food supplements containing it shall be ' Organic silicon (monomethylsi	lanetriol)	
Mycelial extract from	Specified food	Maximum levels	The designation		
Shiitake	category		of the novel		
mushroom (<i>Lentinula</i>	Bread products	2 ml/100 g	food on the labelling of		
edodes)			the feedstuffs		
	Soft drinks	0,5 ml/100 ml	the foodstuffs containing		
	Soft drinks Ready prepared meals	0,5 ml/100 ml 2,5 ml per meal	containing it shall be 'extract from the		
	Ready prepared	2,5 ml per	containing it shall be 'extract from the mushroom <i>Lentinula</i> <i>edodes</i> '		
	Ready prepared meals Foods based	2,5 ml per meal	containing it shall be 'extract from the mushroom <i>Lentinula</i>		

in Directive 2002/46/EC population, excluding pregnant and lactating women 230 mg/day for pregnant and lactating women]	scientific evidence and scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283. Applicant: ChromaDex Inc., 10900 Wilshire Boulevard Suite 600, Los Angeles, CA 90024 USA. During the period of data protection, the novel food is authorised for placing on the market within the Union only by ChromaDex Inc. unless a subsequent applicant obtains
--	---

				ChromaDex Inc. End date of the data protection: 20 February 2025.
Noni fruit juice (<i>Morinda</i> <i>citrifolia</i>)	Specified food category Pasteurised fruit and fruit nectar based drinks	Maximum levels 30 ml with one serving (up to 100 % noni juice) or 20 ml twice a day, not more than 40 ml	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Noni juice' or 'Juice of <i>Morinda</i>	
Noni fruit juice powder (<i>Morinda</i> <i>citrifolia</i>)	Food supplements as defined in Directive 2002/46/EC	6,6 g/day (equivalent to 30 ml of noni juice)	<i>citrifolia</i> ' The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Noni juice powder' or 'Juice powder of <i>Morinda</i> <i>citrifolia</i> '	
Noni fruit puree and concentrate (<i>Morinda</i> <i>citrifolia</i>)	Specified food category Candy/ confectionery Cereal bars Powdered nutritional drink mixes (dry weight) Carbonated beverages Ice cream & sorbet	Maximum levels Fruit puree 45 g/100 g 53 g/100 g 53 g/100 g 11 g/100 g 31 g/100 g	The designation of the novel food on the labelling of the foodstuffs containing it shall be: For fruit puree: ' <i>Morinda</i> <i>citrifolia</i> fruit puree' or 'Noni fruit puree' For fruit concentrate: ' <i>Morinda</i> <i>citrifolia</i>	

Yoghurt	12 g/100 g	fruit concentrate'
Biscuits	53 g/100 g	or 'Noni fruit
Buns, cakes and pastries	53 g/100 g	concentrate'
Breakfast cereals (wholegrain)	88 g/100 g	
Jams and jellies in accordance with Directive 2001/113/EC	133 g/100 g Based on pre- processing quantity to produce final 100 g product	
Sweet spreads, fillings and icings	31 g/100 g	
Savoury sauces, pickles, gravies and condiments	88 g/100 g	
Food Supplements as defined in Directive 2002/46/EC	26 g/day	
	Fruit concentrate	
Candy/ Confectionery	10 g/100 g	
Cereal bars	12 g/100 g	
Powdered nutritional drink mixes (dry weight)	12 g/100 g	
Carbonated beverages	3 g/100 g	
Ice cream & sorbet	7 g/100 g	
Yoghurt	3 g/100 g	
Biscuits	12 g/100 g	
Buns, cakes and pastries	12 g/100 g	

	Due al-f	20 - /100]		I		
	Breakfast cereals (wholegrain)	20 g/100 g					
	Jams and jellies in accordance with Directive 2001/113/EC	30 g/100 g					
	Sweet spreads, fillings and icings	7 g/100 g					
	Savoury sauces, pickles, gravies and condiments	20 g/100 g	-				
	Food Supplements as defined in Directive 2002/46/EC	6 g/day					
Noni leaves (<i>Morinda</i> citrifolia)	Specified food category	Maximum levels	1.	The designation of			
citrifolia)	For the preparation of infusions	A cup of infusion to be consumed shall not be prepared with more than 1 g of dried and roasted leaves of <i>Morinda</i> <i>citrifolia</i>			the novel food on the labelling of the foodstuffs containing it shall be 'Noni leaves' or 'leaves of <i>Morinda</i> <i>citrifolia</i> '.		
			2.	Instructions shall be			

			given to the consu that a cup of infust shoul not be prepa with more than 1 g of dried and roaste leave of Morin	imer ion d red s nda	
Noni fruit powder (Specified food	Maximum levels	The designation		
Morinda citrifolia)	<i>category</i> Food Supplements as defined in Directive 2002/46/EC	2,4 g per/day	of the novel food on the labelling of the foodstuffs containing it shall be ' Morinda citrifolia fruit powder ' or ' Noni fruit powder '		
Odontella	Specified food	Maximum levels	The		
<i>aurita</i> microalgae	category		designation of the novel		
_	Flavoured pasta	1,5 %	food on the labelling of		
	Fish soups	1 %	the foodstuffs containing		
	Marine terrines	0,5 %	it shall be ' Odontella aurita		
	Broth preparations	1 %	microalgae'		
	Crackers	1,5 %]		

	Frozen breaded fish	1,5 %			
Oil enriched	Specified	Maximum	In accordance		
with	food	levels of	with Annex		
phytosterols/	category	phytosterols/	III.5 to		
phytostanols		phytostanols	Regulation		
	Spreadable	1. The	(EU) No		
	fats as	produ	dts69/2011		
	defined in	conta	ining		
	Annex VII,	the			
	Part VII and	novel			
	Appendix	food	diant		
	II, points B and C of	ingre shall	ulent		
	Regulation	be			
	(EU) No	prese	nted		
	1308/2013,	in	liteu		
	and excluding	such			
	cooking and	а			
	frying fats	mann	er		
	and spreads	that			
	based on	they			
	butter or	can			
	other animal	be			
	fat	easily divid			
	Milk based	into	cu		
	products,	portic	ns		
	such as	that	,115		
	products	conta	in		
	based	either			
	on semi- skimmed	а			
	and skimmed	maxi	num		
	milk	of 3			
	products,	g (in			
	possibly with	case of			
	the addition	one			
	of fruits and/	portic	n		
	or cereals,	per	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,		
	products	day)			
	based on fermented	or a			
	milk such as	maxi	num		
	yoghurt and	of 1			
	cheese based	g (in			
	products (fat	case			
	content ≤ 12	of three			
	g per 100	three portio	ne		
	g), where	portic	115		
	possibly the	day)			
	milk fat has	of			
	I		I	I	I

	been reduced and the fat or protein has been partly or fully replaced by vegetable fat or protein Soya drinks Salad dressings, mayonnaise and spicy sauces	 2. The amound of phytophytophytophytophytophytophytophyto	sterols/ stanols. nt sterols/ stanols iner ages ed ings, nnaise s	
Oil extracted	Specified food	Maximum levels	The designation	
from squids	category	of DHA and EPA combined	of the novel food on the labelling of	
	Dairy products except milk-based beverages	200 mg/100 g or for cheese products 600 mg/100 g	the foodstuffs containing it shall be ' Squid oil '.	
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g		
	Spreadable fat and dressings	600 mg/100 g		
	Breakfast cereals	500 mg/100 g		

	Bakery products (breads and bread rolls)	200 mg/100 g			
	Cereal bars	500 mg/100 g			
	Non- alcoholic beverages (including milk-based beverages)	60 mg/100 ml			
	Food Supplements as defined in Directive 2002/46/EC	3 000 mg/day for general population 450 mg/day for pregnant and lactating women			
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products intended			
	Total diet replacement for weight control defined in Regulation (EU) No 609/2013 and meal replacements for weight control	200 mg/meal			
[^{F5} Partially defatted chia	Specified food category	Maximum levels	The designation		
seed (<i>Salvia</i> <i>hispanica</i>) powders	Powder with his content		of the novel food on the labelling of		
P	Unflavoured fermented milk products, including	0,7 %	the foodstuffs containing it shall be 'Partially defatted chia		

natural unflavoured buttermilk (excluding sterilised buttermilk) non-heat- treated after fermentation		seed (<i>Salvia</i> <i>hispanica</i>) powder'	
Unflavoured fermented milk products, heat- treated after fermentation	0,7 %		
Flavoured fermented milk products including heat-treated products	0,7 %		
Confectionery	10 %		
Fruit juices as defined by Directive 2001/112/ EC ^h and vegetable juices	2,5 %		
Fruit nectars as defined by Directive 2001/112/EC and vegetable nectars and similar products	2,5 %		
Flavoured drinks	3 %		
Food supplements as defined in Directive 2002/46/EC excluding food supplements for infants	7,5 g/day		

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	and young children			
	Powder with his content	igh fibre		
	Confectionery	4 %		
	Fruit juices as defined by Directive 2001/112/EC and vegetable juices	2,5 %		
	Fruit nectars as defined by Directive 2001/112/EC and vegetable nectars and similar products	4 %		
	Flavoured drinks	4 %		
	Food supplements as defined in Directive 2002/46/EC excluding food supplements for infants and young children	12 g/day]		
Pasteurised fruit-based preparations	Specified food category	Maximum levels	The wording ' pasteurised by high-	
produced using high- pressure treatment	Types of fruit: apple, apricot, banana, blackberry, blueberry, cherry, coconut, fig, grape, grapefruit, mandarin, mango, melon, peach, pear,		pressure treatment ' shall be displayed next to the name of the fruit preparations as such and in any product in which it is used	

_

[^{F27} Phenylcaps	pineapple, prune, raspberry, rhubarb, strawberry	Maximum	The		Authorised on
	food category	levels	designation		19 December
	Foods for special medical purposes as defined under Regulation (EU) No 609/2013 excluding foods for infants, young children and children under the age of 11 years Food supplements as defined in Directive 2002/46/EC intended for the general population, excluding children under the age of 11 years	2,5 mg/day 2,5 mg/day]	of the novel food on the labelling of the foodstuffs containing it shall be ' phenylcapsaici '.	n	2019. This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283. Applicant: aXichem AB, Södergatan 26, SE 211 34, Malmö Sweden. During the period of data protection, the novel food phenylcapsaicin is authorised for placing on the market within the Union only by aXichem AB, unless a subsequent applicant obtains authorisation for the novel food without reference to the proprietary scientific evidence or scientific data
					to the properties of the total properties of the

					protected in accordance with Article 26 of Regulation (EU) 2015/2283 or with the agreement of aXichem AB.
Phosphated maize starch	Specified food category	Maximum levels	The designation of the novel		
	Baked bakery products	15 %	food on the labelling of		
	Pasta		the foodstuffs containing		
	Breakfast cereals		it shall be ' Phosphated maize starch '		
	Cereal bars		maize staten		
Phosphatidyls from fish phospholipids	food	Maximum levels of phosphatidyls	The designation esim Re novel		
	Beverages based on yoghurt	50 mg/100 ml	food on the labelling of the foodstuffs		
	Powders based on milk powders	3 500 mg/100 g (equivalent to 40 mg/100 ml ready to drink)	containing it shall be ' Fish phosphatidylse	rine	
	Foods based on yoghurt	80 mg/100 g			
	Cereal bars	350 mg/100 g			
	Chocolate based confectionary	200 mg/100 g			
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In compliance with Regulation (EU) No 609/2013			
	Food supplements as defined	300 mg/day			

	in Directive 2002/46/EC				
Phosphatidyls from soya phospholipids	food	Maximum levels of phosphatidyls	The designation		
	Beverages based on yoghurt	50 mg/100 ml	food on the labelling of the foodstuffs containing it shall be ' Soya phosphatidylse		
	Powders based on milk powder	3,5 g/100 g (equivalent to 40 mg/100 ml ready to drink)		rine	
	Foods based on yoghurt	6 6			
	Cereal bars	350 mg/100 g			
	Chocolate 200 mg/100 g based confectionary				
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In compliance with Regulation (EU) No 609/2013			
Phospholipid product	Specified food category	Maximum levels of phosphatidyls	The designation	The product is not	
containing equal amounts of phosphatidyls	Breakfast cereals	80 mg/100 g	food on the labelling of the foodstuffs	intended to be marketed to pregnant	
and	Cereal bars	350 mg/100 g	containing	or breast- feeding	
phosphatidic acid	Foods based on yogurt	80 mg/100 g	shall be ' Soy phosphatidylse and	women rine	
	Soy-based yogurt-like products	80 mg/100 g	phosphatidic acid '		
	Yogurt based- drinks	50 mg/100 g			
	Soy-based yogurt-like drinks	50 mg/100 g			
	Powders based on milk powder	3,5 g/100 g (equivalent to 40 mg/100			

	Food Supplements as defined in Directive 2002/46/EC Foods for special medical purposes as defined in Regulation (EU) No 609/2013	ml ready-to drink) 800 mg/day In compliance with Regulation (EU) No 609/2013		
Phospholipide	_S Specified food	Maximum levels		
from egg yolk	jooa category	ieveis		
•	Not specified			
Phytoglycoger	Specified food category	Maximum levels	The designation of the novel	
	Processed foods	25 %	food on the labelling of the foodstuffs containing it shall be ' Phytoglycogen '	
Phytosterols/ phytostanols	Specified food category	Maximum levels	In accordance with Annex III.5 of	
	Rice drinks	1 They	Regulation	
	Rye bread with flour containing $\geq 50 \%$ rye (wholemeal rye flour, whole or cracked rye kernels and rye flakes) and $\leq 30 \%$ wheat; and with $\leq 4 \%$ added sugar but no fat added.	1. They shall be prese in such a mann that they can be easily divide into portic that	nted er ed	

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.

Changes that have b	een made a	ppear in th	e content and	are referer	iced with a	nnotations	. (See ei	nd of Do	cument for	details)

[
Salad	contain
dressings,	either
mayonnaise	a .
and spicy	maximum
sauces.	of 3
0 1 1	g (in
Soya drink	case
Milk type	of 1
products,	portion/
such as semi-	day)
skimmed and	or a
skimmed	maximum
milk type	of 1
products,	g (in
possibly with	case
the addition	of 3
of fruits and/	portions/
	day)
or cereals,	of
where	added
possibly the	phytosterols/
milk fat has	phytostanols.
been reduced,	The
or where milk	amount of
fat and/or	phytosterols/
protein has	phytostanols
been partly or	added to a
fully replaced	container of
by vegetable	
fat and/or	beverages
protein.	shall not
Draduata	exceed 3 g.
Products	Salad
based on	dressings,
fermented	mayonnaise
milk such	and spicy
as yoghurt	sauces shall
and cheese	be packed
type products	as single
(fat content	portions
< 12 % per	
100 g), where	
possibly the	
milk fat has	
been reduced,	
or where milk	
fat and/or	
protein has	
been partly or	
fully replaced	
by vegetable	
fat and/or	
protein	
protein	

Status: Point in time view as at 31/12/2020.
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Spreadable fats as defined in Annex VII, Part VII and Appendix II, points B and C of Regulation (EU) No 1308/2013, and excluding cooking and frying fats and spreads based on butter or other animal fat. Food Supplements as defined in Directive	3 g/day		
Plum kernel oil	2002/46/EC Specified food category	Maximum levels		
	For frying and as seasoning	In line with normal food use of vegetable oils		
Potato proteins (coagulated) and hydrolysates thereof	Not specified	1	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Potato protein '	
Prolyl oligopeptidaso (enzyme preparation)	Specified food category Food Supplements as defined in Directive 2002/46/ EC for	Maximum levels 120 PPU/ day (2,7 g of enzyme preparation/ day) (2 × 10 ⁶ PPI/day)	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' Prolyl	

[^{F28} Protein	general adult population Specified	PPU – Prolyl Peptidase Units or Proline Protease Units PPI – Protease Picomole International Maximum	oligopeptidase	
extract from pig kidneys	<i>food</i> <i>category</i> Food Supplements as defined in Directive 2002/46/EC Food for special medical purposes as	<i>levels</i> 3 capsules or 3 tablets/day; equalising 12,6 mg pig kidney extract a day Diamine oxidase (DAO) content: 0,9		
[^{F29} Pyrroloqui	defined in Regulation (EU) No 609/2013] nonecified food	mg/day (3 capsules or 3 tablets with a content of DAO of 0,3 mg/capsule or 0,3 mg/tablet) Maximum levels	The designation	Authorised on 2 September
quinone disodium salt	<i>category</i> Food Supplements as defined in Directive 2002/46/ EC intended for the adult population, excluding pregnant and lactating women	20 mg/day]	of the novel food on the labelling of the foodstuffs containing it shall be 'Pyrroloquinol quinone disodium salt'. Food supplements containing Pyrroloquinolin quinone disodium salt shall bear the following statement:	2018. This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283. Applicant: Mitsubishi Gas Chemical Company, Inc., Mitsubishi Building 5-2

This foo supplem should b consume adults or excludir pregnan and lacts women	ent 2-chome, Chiyoda- ku, Tokyo nly 100-8324, Japan. During t the period
--	---

Rapeseed oil high in unsaponifiab	Specified food le category	Maximum levels	The designation of the novel	
matter	Food Supplements as defined in Directive 2002/46/EC	1,5 g per portion recommended for daily consumption	food on the labelling of the foodstuffs containing it shall be ' Rapeseed oil extract '	
Rapeseed Protein	As a vegetable protein source in foods except in infant formula and follow-on formula		 The design of the novel food on the labell of the foods contain it shall be 'Rape protei Any foods contain 'rapes protein shall bear a statem that this ingrea a statem that this ingrea a llerg reacting to mustain and 	ing tuffs ining eseed n'. tuff ining seed n' nent dient dient ic on mers ic

			produ thered When releva this staten shall appea in close proxit to the list of ingred	of. e ant, nent r	
[^{F30} Refined shrimp peptide concentrate	Specified food category Food Supplements as defined in Directive 2002/46/EC for the adult population	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be ' refined shrimp peptide concentrate '.		Authorised on 20 November 2018. This inclusion is based on proprietary scientific evidence and scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283. Applicant: Marealis AS., Stortorget 1, Kystens Hus, 2nd floor, N-9008 Tromsø Postal address: P.O. Box 1065, 9261 Tromsø, Norway. During the period of data protection the novel food refined shrimp peptideconcentrate is authorised

						for placing on the market within the Union only by Marealis AS unless a subsequent applicant obtains authorisation for the novel food without reference to the proprietary scientific evidence or scientific data protected in accordance with Article 26 of Regulation (EU) 2015/2283 or with the agreement of Marealis AS. End date of the data protection: 20 November 2023.
Trans- resveratrol	Specified food category	Maximum levels	1.	The desig of	nation	
	Food Supplements as defined in Directive 2002/46/ EC for adult population (capsule or tablet form)	150 mg/day		the novel food on the labell of the food	ements ining	

			2.	- resveratrol'. The labelling of food supplements containing trans- resveratrol shall bear a statement that people using medicines should only consume the product under medical supervision.
Trans- resveratrol (microbial source)	Specified food category Food supplements as defined in Directive 2002/46/EC	Maximum levels	1.	The designation of the novel food on the labelling of the food supplements containing it shall be ' <i>Trans</i> - resveratrol'. The labelling of food supplements containing of the labelling of the labelling of food supplements containing

			trans- resveratrol shall bear a statement that people using medicines should only consume the product under medical supervision.	
Rooster comb extract	Specified food category Milk-based	<i>Maximum</i> <i>levels</i> 40 mg/100 g	The designation of the novel food on the	
	drinks	or mg/100 ml	labelling of	
	Milk based fermented drinks	80 mg/100 g or mg/100 ml	the foodstuffs containing it shall be ' Rooster comb	
	Yoghurt-type products	65 mg/100 g or mg/100 ml	extract ' or ' Cockerel	
	Fromage frais	110 mg/100 g or mg/100 ml	comb extract	
Sacha inchi oil from <i>Plukenetia</i>	Specified food category	Maximum levels	The designation of the novel	
volubilis	As for linseed oil	normal food use of linseed oil	food on the labelling of the foodstuffs containing it shall be ' Sacha inchi oil (Plukenetia volubilis) '	
Salatrims	Specified food category	Maximum Ievels	1. The designation of	
	Bakery products and confectionary		the novel food on	

		that the produ are not intend for use by child	ded
d levels egory of DH and H comb	S de HA of EPA foo bined : lat mg/day the co it s 'D EF	esignation `the novel od on the belling of e foodstuffs ontaining shall be DHA and PA-rich	
	l levels gory of Di and l comb l 3 000 blements efined irective	Itevels Itevels def of DHA of DHA of fo and EPA fo combined : la diments state control offined it control irrective fill control or adult oil control	cified I goryMaximum levels of DHA and EPA combined :The designation of the novel food on the labelling of the foodstuffs containing it shall be 'DHA and EPA-rich oil from the

excluding pregnant and lactating women		Schizochytrium sp.'	ł
Food Supplements as defined in Directive 2002/46/EC for pregnant and lactating women	450 mg/day		
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal		
Milk-based drinks and similar products intended for young children	200 mg/100 g		
Processed cereal based food and baby food for infants and young children as defined in Regulation			

(EU) No 609/2013		
Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen		
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014		
Bakery Products (Breads, Rolls and Sweet Biscuits)	200 mg/100 g	
Breakfast Cereals	500 mg/100 g	
Cooking Fats	360 mg/100 g	
Dairy Analogues except drinks	600 mg/100 g for cheese; 200 mg/100 g for soy and imitation milk products (excluding drinks)	
Dairy Products except milk- based drinks	600 mg/100 g for cheese; 200 mg/100 g for milk products (including	

	Non- alcoholic Beverages (including dairy analogue and milk-based drinks)	milk, fromage frais and yoghurt products; excluding drinks) 80 mg/100 g			
	Cereal/ Nutrition Bars	500 mg/100 g			
	Spreadable Fats and Dressings	600 mg/100 g			
[^{F31} Schizochyth sp. (ATCC PTA-9695)	ritmecified food category	Maximum levels of DHA	The designation of the novel		
oil	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g	food on the labelling of the foodstuffs containing it shall be		
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g	'Oil from the microalgae <i>Schizochytrium</i> sp.'	t	
	Spreadable fats and dressings	600 mg/100 g			
	Breakfast cereals	500 mg/100 g			
	Food Supplements as defined in Directive	250 mg DHA/day for general population			
	2002/46/EC	450 mg DHA/day for pregnant and lactating women			

Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal
Milk-based drinks and similar products intended for young children	200 mg/100 g
Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen	
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014	
Foods for special medical purposes as defined in Regulation	In accordance with the particular nutritional requirements

	(EU) No 609/2013	of the persons for whom the products are intended		
	Bakery products (breads, rolls, and sweet biscuits)	200 mg/100 g		
	Cereal bars	500 mg/100 g		
	Cooking fats	360 mg/100 g	-	
	Non- alcoholic beverages (including dairy analogue and milk-based drinks)	80 mg/100 ml		
	Infant formula and follow-on formula as defined in Regulation (EU) No 609/2013	In accordance with Regulation (EU) No 609/2013	-	
	Processed cereal-based foods and baby foods for infants and young children as defined in Regulation (EU) No 609/2013	200 mg/100 g		
	Fruit/ vegetable puree	100 mg/100 g]	-	
<i>[^{F32}Schizochyth</i> sp. oil	ritimecified food category	Maximum levels of DHA	The designation of the novel	
	Dairy products	200 mg/100 g or for	food on the labelling of	

except milk- based drinks	cheese products 600 mg/100 g	the foodstuffs containing it shall be
Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g	'Oil from the microalgae <i>Schizochytrium</i> sp.'
Spreadable fat and dressings	600 mg/100 g	
Breakfast cereals	500 mg/100 g	
Food Supplements as defined in Directive	250 mg DHA/day for general population	
2002/46/EC	450 mg DHA/day for pregnant and lactating women	
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal	
Milk-based drinks and similar products intended for young children	200 mg/100 g	
Processed cereal-based foods and baby foods for infants and young		

children as defined in Regulation (EU) No 609/2013		
Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen		
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Implementing Regulation (EU) No 828/2014		
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended	
Bakery products (breads, rolls, and, sweet biscuits)	200 mg/100 g	
Cereal bars	500 mg/100 g	
Cooking fats	360 mg/100 g	

	Non- alcoholic beverages (including dairy analogue and milk-based drinks) Fruit/ vegetable puree	80 mg/100 ml 100 mg/100 g]			
<i>[^{F18}Schizochyta</i> sp. (T18) oil	food category Dairy products except milk- based drinks Dairy analogues	Maximum levels 200 mg/100 g or for cheese products 600 mg/100 g 200 mg/100 g or for	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Oil from the microalgae		
	except drinks Spreadable fats and dressings Breakfast	analogues to cheese products 600 mg/100 g 600 mg/100 g	Schizochytrium sp.'.		
	Food supplements as defined in Directive 2002/46/EC	500 mg/100 g 250 mg DHA/day for general population 450 mg			
		DHA/day for pregnant and lactating women			
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements	250 mg/meal			

for weight control	
Milk-based drinks and similar products intended for young children	200 mg/100 g
Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen	
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014	
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended
Bakery products (breads, rolls and, sweet biscuits)	200 mg/100 g

			٦	1	1
	Cereal bars	500 mg/100g			
	Cooking fats	360 mg/100 g			
	Non- alcoholic beverages (including dairy analogue and milk-based drinks)	80 mg/100 ml			
	Infant formula and follow-on formula as defined in Regulation (EU) No 609/2013	In accordance with Regulation (EU) No 609/2013	-		
	Processed cereal-based foods and baby foods for infants and young children as defined in Regulation (EU) No 609/2013	200 mg/100 g			
	Fruit/ vegetable puree	100 mg/100 g]			
[^{F33} Syrup from <i>Sorghum</i> <i>bicolor</i> (L.) Moench (Traditional food from a third country)	Not specified		The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Sorghum (<i>Sorghum</i> <i>bicolor</i>) syrup']		
Fermented soybean extract	Specified food category	Maximum levels	1. The desig of the	nation	

	Food Supplements as defined in Directive 2002/46/EC (capsules, tablets or powder form) intended for the adult population, excluding pregnant and lactating women	100 mg/day	it shall be 'Ferm soybe extra 2. The label of food suppl conta fermo soybe extra shall bear a staten that perso takin medi shoul only consu	ling stuffs ining mented ean ct'. ling ements ining ented ean ct ment ms g cation d ume act	
Spermidine- rich wheat germ extract (<i>Triticum</i>	Specified food category	Maximum levels	The designation of the novel food on the		
(Trucum aestivum)	Food Supplements as defined in Directive 2002/46/ EC intended for the adult population,	Equivalent of max. 6 mg/day spermidine	labelling of the food supplements containing it shall be ' spermidine-		

	excluding pregnant and lactating women		rich wheat germ extract '	
Sucromalt	Specified food category	Maximum levels	of	gnation
	Not specified		the nove food on the label of the food conta it shall be 'Suc2.The desig of the nove food on the label 	Illing stuffs aining romalt'. gnation Illing mpanied cation uct
			sourd of gluce and fruct	ose
Sugar cane fibre	Specified food category	Maximum levels		
	Bread	8 %		
	1	1	1	1

			_	
	Meat and muscle products	3 %		
	Seasonings and spices	3 %		
	Grated cheeses	2 %	_	
	Special diet foods	5 %	_	
	Sauces	2 %		
	Beverages	5 %		
[^{F34} Sugars obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp	Not specified		The designation of the novel food on the labelling of the foodstuffs containing it shall be 'sugars obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp', 'Glucose obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp' or 'Fructose obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp' or 'Fructose obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp', depending on the form used.]	
Sunflower oil extract	Specified food	Maximum levels	The designation	
	category	1 1 ~/d	of the novel	
	Food Supplements as defined in Directive 2002/46/EC	1,1 g/day	food on the labelling of the foodstuffs containing it shall be '	

			Sunflower oil extract '		
Dried <i>Tetraselmis</i> <i>chuii</i> microalgae	Specified food category	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Dried microalgae <i>Tetraselmis</i> <i>chuii</i> ' or 'Dried microalgae <i>T.</i> <i>chuii</i> ' Food supplements containing dried microalgae <i>Tetraselmis</i> <i>chuii</i> shall bear the following statement: 'Contains negligible amounts of iodine'		
	Sauces	20 % or 250mg/day			
	Special salts	1 %			
	Condiment	250 mg/day			
	Food Supplements as defined in Directive 2002/46/EC	250 mg/day			
<i>Therapon barcoo /</i> Scortum	Intended use identical to that of the salmon, namely the preparation of culinary fish products and dishes, including cooked, raw, smoked and baked fish products				
D-Tagatose	Specified food category	Maximum levels	1. The desig of	nation	
	Not specified		the novel food on the labell of the foods conta	ling	

			2. The label of any produce when the level of D-Tagar exceed 15 g per servities and all beven contained great than 1 % D-Tagar (as consist shall bear a statem 'exceed the contained of the conta	act e tose eds ng tages ining er tose umed) ment ssive umption ace ive	
[^{F18} Taxifolin- rich extract	Specified food category	Maximum levels	The designation of the novel		
	Yogurt plain/ Yogurt with fruits ^(*)	0,020 g/kg	food on the labelling of the foodstuffs		
	Kephir ^(*)	0,008 g/kg	containing it shall be '		
	Buttermilk (*)	0,005 g/kg	taxifolin-rich extract '		

	Milk powder (*) Cream (*) Sour cream (*) Cheese (*) Butter (*) Chocolate confectionery Non- alcoholic	0,052 g/kg 0,070 g/kg 0,050 g/kg 0,090 g/kg 0,164 g/kg 0,070 g/kg 0,020 g/L				
	beverages Food supplements as defined in Directive 2002/46/EC intended for the general population, excluding infants, young children, children and adolescents younger than 14 years	100 mg/day				
	produ rich e repla in par	n used in milk acts Taxifolin- extract may not ce in whole or rt, any milk ituent]	-			
Trehalose	Specified food category Not specified	Maximum levels	1.	The design of the novel food on the labelli of the foodst contain	ng uffs	

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

			2.	it shall be 'Trehalose' and shall be displayed on the labelling of the product as such or in the list of ingredients of foodstuffs containing it. The designation of the novel food on the labelling shall be accompanied by indication that the 'Trehalose is a source of glucose'.
[^{F18} UV- treated mushrooms (<i>Agaricus bisporus</i>)	Specified food category Mushrooms (Agaricus bisporus)	Maximum levels of vitamin D ₂ 20 μg of vitamin D ₂	1.	The designation on the label of

/100 g fresh		the
weight]		novel
		food
		as
		such
		or
		of
		the
		foodstuffs
		containing
		it
		shall
		be
		'UV-
		treated
		mushrooms
		(
		Agaricus
		bisporus
)'.
	2.	The
		designation
		on
		the
		label
		of
		the
		novel
		food
		as
		such
		or
		of
		the
		foodstuffs
		containing
		it
		shall
		be
		accompanied
		by
		indication
		that
		a
		'controlled
		light
		treatment
		was
		used
		to
		increase
		vitamin
		D
I		i I

			levels' or 'UV treatment was used to increase vitamin D_2 levels'.
[^{F18} UV- treated baker's yeast (<i>Saccharomyce</i> <i>cerevisiae</i>)	Specified food category Yeast- leavened breads and rolls Yeast- leavened fine bakery wares Food supplements as defined in Directive 2002/46/EC	Maximum levels of vitamin D ₂ 5 μg of vitamin D ₂ /100 g 5 μg of vitamin D ₂ /100 g	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Vitamin D yeast' or 'Vitamin D ₂ yeast'
	Pre-packed fresh or dry yeast for home baking	45 μg/100 g for fresh yeast 200 μg/100 g for dried yeast	1.The designation of the novel food on the labelling of the foodstuffs shall be 'Vitamin D yeast' or 'Vitamin D 2 yeast'.2.The labelling of of

	the
	novel
	food
	shall
	bear
	a
	statement
	that
	the
	foodstuff
	is
	only
	intended
	for
	baking
	and
	that
	it
	should
	not
	be
	eaten
	raw.
3.	The
	labelling
	of
	the
	novel
	food
	shall
	bear
	instructions
	for
	use
	for
	the
	final
	consumers
	so
	that
	a
	a maximum
	concentration
	of 5
	μg/100
	g of
	vitamin
	D 2
	in
	final
	home#
	baked
	products

UV-treated bread	Specified food category Yeast leavened bread and rolls (without toppings)	Maximum levels of vitamin D ₂ 3 μg vitamin D ₂ /100 g	is not exceeded.] The designation on the label of the novel food shall be accompanied by ' contains vitamin D produced by UV-treatment	
UV-treated milk	Specified food category Pasteurised whole milk as defined in Regulation (EU) No 1308/2013 to be consumed as such Pasteurised semi- skimmed milk as defined in Regulation (EU) No 1308/2013 to be consumed as such	Maximum levels of vitamin D ₃ 5-32 μg/kg for general population excluding infants 1-15 μg/kg for general population excluding infants	1. The designation on the label of the novel food shall be 'UV- treated'. 2. Where UV- treated milk contains an amount of vitamin D that is considered significant in accordance with Point 2 of Part A of Annex XIII to Regulation	

			(EU) No 1169/2011 of the European Parliament and of the Council, the designation for the labelling shall be accompanied by 'contains vitamin D produced by UV- treatment' or 'milk containing vitamin D resulting from UV- treatment'.	
[^{F7} Vitamin D ₂ mushroom powder	Specified food category	Maximum levels of vitamin D 2 ^k	The designation of the novel food on the	Authorised on 27 August 2020. This inclusion
	Breakfast cereals	2,25 μg of vitamin D ₂ /100 g	labelling of the foodstuffs containing it shall be	is based on proprietary scientific evidence and
	Yeast- leavened bread and pastries	2,25 μg of vitamin D ₂ /100 g	'UV-treated mushroom powder containing	scientific data protected in accordance with Article
	Grain products and pastas	2,25 μg of vitamin D ₂ /100 g	vitamin D' or 'UV-treated mushroom powder	26 of Regulation (EU) 2015/2283.

Fruit juice and fruit/ vegetable blend beverages Milk and dairy products (excluding fluid milks)	1,125 μg of vitamin D ₂ /100 mL 2,25 μg of vitamin D ₂ /100 g/1,125 μg of vitamin D ₂ /100 mL	containing vitamin D ₂ ' The labelling of food supplements containing vitamin D ₂ mushroom powder shall bear a statement	Applicant: Oakshire Naturals, LP., PO Box 388 Kennett Square, Pennsylvania 19348, United States. During the period of data		
Cheese (excluding cottage cheese, ricotta cheese, and hard-grating cheeses)	(beverages) 2,25 μg of vitamin D ₂ /100 g	that they should not be consumed by infants	should not be consumed by	should not be consumed by	protection, the novel food vitamin D_2 mushroom powder is authorised for placing on the market within the
Meal replacement bars and beverages	$\begin{array}{c} 2,25 \ \mu g \ of \\ vitamin \ D_{2} \\ /100 \ g/1,125 \\ \mu g \ of \ vitamin \\ D_{2} \ /100 \ mL \\ (beverages) \end{array}$			Union only by Oakshire Naturals, LP., unless a subsequent applicant	
Dairy analogues	$\begin{array}{c} 2,25 \ \mu g \ of \\ vitamin \ D \ _2 \\ /100 \ g/1,125 \\ \mu g \ of \ vitamin \\ D \ _2 \ /100 \ mL \\ (beverages) \end{array}$				obtains authorisation for the novel food without reference to the proprietary
Meat analogues	2,25 μg of vitamin D ₂ /100 g	-	scientific evidence or scientific data protected in		
Soups and broths	2,25 μg of vitamin D ₂ /100 g	-	accordance with Article 26 of		
Extruded vegetable snacks	2,25 μg of vitamin D ₂ /100 g	-	Regulation (EU) 2015/2283 or with the		
Foods for Special Medical Purposes as defined under Regulation (EU) No	15 μg/day		agreement of Oakshire Naturals, LP. End date of the data protection: 27 August 2025.		

Status: Point in time view as at 31/12/2020.
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	609/2013 excluding those intended for infants Food supplements as defined in Directive 2002/46/EC intended for the general population excluding infants	15 μg/day]			
Vitamin K ₂ (menaquinone	To be used in c with Directive EC, Regulatior 609/2013 and/o (EC) No 1925/	2002/46/ n (EU) No or Regulation	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Menaquinone or 'Vitamin K 2'		
Wheat bran extract	Specified food category	Maximum levels	The designation of the novel	The 'Wheat Bran Extract 'may not be	
	Beer and substitutes	0,4 g/100 g	food on the labelling of the foodstuffs containing it shall be ' Wheat bran extract '	introduced onto the market as a food supplement or food supplement ingredient. Nor may it be added to infant formula.	
	Ready to eat cereals	9 g/100 g			
	Dairy products	2,4 g/100 g			
	Fruit and vegetable juices	0,6 g/100 g			
	Soft drinks	0,6 g/100 g			
	Meat preparations	2 g/100 g			
[^{F35} Xylo- oligosaccharic	Specified efood category	Maximum levels ^j	The designation of the novel		
	White bread	14 g/kg	food on the		
	Wholemeal bread	14 g/kg	labelling of the foodstuffs containing		

	Breakfast cereals	14 g/kg	it shall be ' Xylo-	
	Biscuits	14 g/kg	oligosaccharides	
	Soy drink	3,5 g/kg		
	Yoghurt ⁱ	3,5 g/kg		
	Fruit spreads	30 g/kg		
	Chocolate confectionery	30 g/kg		
	Food supplements as defined in Directive 2002/46/ EC for the general adult population	2 g/day]		
[^{F36} Yarrowia lipolytica yeast biomass	Specified food category	Maximum levels	The designation of the novel	
	Food Supplements as defined in Directive 2002/46/EC, excluding food supplements for infants and young children	6 g/day for children from 10 years of age, adolescents and general adult population 3 g/day for children from 3 to 9 years of age]	food on the labelling of the foodstuffs containing it shall be ' Yarrowia lipolytica yeast heat- killed biomass'	
Yeast beta- glucans	Specified food category	cervisiae)	The designation of the novel food on the labelling of Ch e foodstuffs containing	
	Food supplements as defined in Directive 2002/46/EC, excluding food supplements for infants	1,275 g/day for children older than 12 years and general adult population 0,675 g/day for children younger than 12 years	it shall be 'Yeast (<i>Saccharomyces</i> <i>cerevisiae</i>) beta-glucans'	

and young children	
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	1,275 g/day
Food for special medical purposes as defined in Regulation (EU) No 609/2013, excluding food for special medical purposes intended for infants and young children	1,275 g/day
Beverages based on fruit and/or vegetable juices including concentrate and dehydrated juices	1,3 g/kg
Fruit- flavoured drinks	0,8 g/kg
Cocoa beverages preparation powder	38,3 g/kg (powder)
Other beverages	0,8 g/kg (ready to drink)
	7 g/kg (powder)

			1		
	Cereal bars	6 g/kg			
	Breakfast cereals	15,3 g/kg			
	Wholegrain and high fibre instant hot breakfast cereals	1,5 g/kg			
	Cookie-type biscuits	6,7 g/kg			
	Cracker-type biscuits	6,7 g/kg			
	Milk based beverages	3,8 g/kg			
	Fermented milk products	3,8 g/kg			
	Milk product analogues	3,8 g/kg			
	Dried milk/ milk powder	25,5 g/kg			
	Soups and soup mixes	0,9 g/kg (ready to eat)			
		1,8 g/kg (condensed)			
		6,3 g/kg (powder)			
	Chocolate and confectionery	4 g/kg			
	Protein bars and powders	19,1 g/kg			
	Jam, marmalade and other fruit spreads	11,3 g/kg			
[^{F37} Zeaxanthin	Specified	Maximum	The		
	food category	levels	designation of the novel		
	Food Supplements as defined in Directive 2002/46/EC	2 mg/day]	food on the labelling of the foodstuffs containing it shall be ' Zeaxanthin '.		

Zinc L- pidolate	Specified food category	Maximum levels	The designation of the novel		
	Foods covered by Regulation (EU) No 609/2013	3 g/day	food on the labelling of the foodstuffs containing it shall be ' Zinc L- pidolate '		
	Milk based drinks and similar products intended for young children				
	Meal replacement for weight control				
in to ex of m ef es	Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen	-			
	Food bearing statement on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014				
	Food Supplements as defined in Directive 2002/46/EC				

a Regulation (EU) No 609/2013 of the European Parliament and of the Council of 12 June 2013 on food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control and repealing

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

Council Directive 92/52/EEC, Commission Directives 96/8/EC, 1999/21/EC, 2006/125/EC and 2006/141/EC, Directive 2009/39/EC of the European Parliament and of the Council and Commission Regulations (EC) No 41/2009 and (EC) No 953/2009 (OJ L 181, 29.6.2013, p. 35).

- **b** Commission Implementing Regulation (EU) No 828/2014 of 30 July 2014 on the requirements for the provision of information to consumers on the absence or reduced presence of gluten in food (OJ L 228, 31.7.2014, p. 5).
- c Directive 2002/46/EC of the European Parliament and of the Council of 10 June 2002 on the approximation of the laws of the Member States relating to food supplements (OJ L 183, 12.7.2002, p. 51).
- **d** Regulation (EC) No 1925/2006 of the European Parliament and of the Council of 20 December 2006 on the addition of vitamins and minerals and of certain other substances to foods (OJ L 404, 30.12.2006, p. 26).
- e Council Directive 2001/113/EC of 20 December 2001 relating to fruit jams, jellies and marmalades and sweetened chestnut purée intended for human consumption (OJ L 10, 12.1.2002, p. 67).
- f Regulation (EU) No 1308/2013 of the European Parliament and of the Council of 17 December 2013 establishing a common organisation of the markets in agricultural products and repealing Council Regulation (EEC) No 922/72, (EEC) No 234/79, (EC) No 1037/2001 and (EC) 1234/2007 (OJ L 347, 20.12.2013, p. 671).
- g [^{F4}Maximum use levels in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer.]
- h [^{F5}Council Directive 2001/112/EC of 20 December 2001 relating to fruit juices and certain similar products intended for human consumption (OJ L 10, 12.1.2002, p. 58).]
- i [^{F6}When used in milk products xylo-oligosaccharides shall not replace, in whole or in part, any milk constituent.
- j Maximum levels calculated on the basis of the specifications of Powder form 1.]
- **k** [^{F7}The minimum specification for vitamin D content in vitamin D₂ mushroom powder of 1 000 μ g vitamin D₂/gram of mushroom powder is used.]

Textual Amendments

- **F4** Inserted by Commission Implementing Regulation (EU) 2019/1294 of 1 August 2019 authorising the placing on the market of betaine as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- F5 Inserted by Commission Implementing Regulation (EU) 2020/500 of 6 April 2020 authorising the placing on the market of partially defatted chia seed (Salvia hispanica) powders as novel foods under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- F6 Inserted by Commission Implementing Regulation (EU) 2020/916 of 1 July 2020 authorising the extension of use of xylo-oligosaccharides as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- F7 Inserted by Commission Implementing Regulation (EU) 2020/1163 of 6 August 2020 authorising the placing on the market of vitamin D2 mushroom powder as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F8** Inserted by Commission Implementing Regulation (EU) 2019/506 of 26 March 2019 authorising the placing on the market of D-ribose as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council, and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- F9 Substituted by Commission Implementing Regulation (EU) 2019/110 of 24 January 2019 authorising an extension of use of Allanblackia seed oil as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F10** Substituted by Commission Implementing Regulation (EU) 2019/1686 of 8 October 2019 authorising the extension of use of bovine milk basic whey protein isolate as a novel food under Regulation (EU)

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

- F11 Substituted by Commission Implementing Regulation (EU) 2020/24 of 13 January 2020 authorising an extension of use of chia seeds (Salvia hispanica) as a novel food and the change of the conditions of use and the specific labelling requirements of chia seeds (Salvia hispanica) under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- F12 Inserted by Commission Implementing Regulation (EU) 2018/1631 of 30 October 2018 authorising the placing on the market of cranberry extract powder as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F13** Inserted by Commission Implementing Regulation (EU) 2018/2016 of 18 December 2018 authorising the placing on the market of decorticated grains of Digitaria exilis as a traditional food from a third country under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- F14 Inserted by Commission Implementing Regulation (EU) 2018/1133 of 13 August 2018 authorising the placing on the market of dried aerial parts of Hoodia parviflora as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F15** Substituted by Commission Implementing Regulation (EU) 2019/1272 of 29 July 2019 correcting Implementing Regulation (EU) 2017/2470 establishing the Union list of novel foods and Implementing Decision (EU) 2017/2078 authorising an extension of use of yeast beta-glucans as a novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council (Text with EEA relevance).
- **F16** Inserted by Commission Implementing Regulation (EU) 2020/1559 of 26 October 2020 amending Implementing Regulation (EU) 2017/2470 establishing the Union list of novel foods (Text with EEA relevance).
- F17 Inserted by Commission Implementing Regulation (EU) 2018/1647 of 31 October 2018 authorising the placing on the market of egg membrane hydrolysate as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F18** Substituted by Commission Implementing Regulation (EU) 2020/1559 of 26 October 2020 amending Implementing Regulation (EU) 2017/2470 establishing the Union list of novel foods (Text with EEA relevance).
- **F19** Inserted by Commission Implementing Regulation (EU) 2020/206 of 14 February 2020 authorising the placing on the market of fruit pulp, pulp juice, concentrated pulp juice from Theobroma cacao L. as a traditional food from a third country under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F20** Inserted by Commission implementing Regulation (EU) 2019/1979 of 26 November 2019 authorising the placing on the market of 2'-Fucosyllactose/Difucosyllactose mixture as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F21** Inserted by Commission Implementing Regulation (EU) 2020/917 of 1 July 2020 authorising the placing on the market of infusion from coffee leaves of Coffea arabica L. and/or Coffea canephora Pierre ex A. Froehner as a traditional food from a third country under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F22** Substituted by Commission Implementing Regulation (EU) 2018/1293 of 26 September 2018 amending Implementing Regulation (EU) 2017/2470 as regards the conditions of use of the novel food lactitol (Text with EEA relevance).
- **F23** Inserted by Commission Implementing Regulation (EU) 2020/484 of 2 April 2020 authorising the placing on the market of lacto-N-tetraose as a novel food under Regulation (EU) 2015/2283 of the European

Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

- F24 Inserted by Commission Implementing Regulation (EU) 2018/1991 of 13 December 2018 authorising the placing on the market of berries of Lonicera caerulea L. as a traditional food from a third country under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F25** Inserted by Commission Implementing Regulation (EU) 2018/1123 of 10 August 2018 authorising the placing on the market of 1-methylnicotinamide chloride as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F26** Inserted by Commission Implementing Regulation (EU) 2020/16 of 10 January 2020 authorising the placing on the market of nicotinamide riboside chloride as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F27** Inserted by Commission Implementing Regulation (EU) 2019/1976 of 25 November 2019 authorising the placing on the market of Phenylcapsaicin as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F28** Substituted by Commission Implementing Regulation (EU) 2020/973 of 6 July 2020 authorising a change of the conditions of use of the novel food 'protein extract from pig kidneys' and amending Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F29** Inserted by Commission Implementing Regulation (EU) 2018/1122 of 10 August 2018 authorising the placing on the market of pyrroloquinoline quinone disodium salt as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F30** Inserted by Commission Implementing Regulation (EU) 2018/1633 of 30 October 2018 authorising the placing on the market of refined shrimp peptide concentrate as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- F31 Substituted by Commission Implementing Regulation (EU) 2019/387 of 11 March 2019 authorising an extension of use of Schizochytrium sp. (ATCC PTA-9695) oil as a novel food and the change of the designation and of the specific labelling requirement of Schizochytrium sp. (ATCC PTA-9695) oil under Regulation (EU) 2015/2283 of the European Parliament and of the Council, and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F32** Substituted by Commission Implementing Regulation (EU) 2019/109 of 24 January 2019 authorising an extension of use of Schizochytrium sp. oil as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council, and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F33** Inserted by Commission Implementing Regulation (EU) 2018/2017 of 18 December 2018 authorising the placing on the market of syrup from Sorghum bicolor (L.) Moench as a traditional food from a third country under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F34** Inserted by Commission Implementing Regulation (EU) 2020/1634 of 4 November 2020 authorising the placing on the market of sugars obtained from cocoa (Theobroma cacao L.) pulp as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F35** Substituted by Commission Implementing Regulation (EU) 2020/916 of 1 July 2020 authorising the extension of use of xylo-oligosaccharides as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F36** Inserted by Commission Implementing Regulation (EU) 2019/760 of 13 May 2019 authorising the placing on the market of Yarrowia lipolytica yeast biomass as a novel food under Regulation (EU)

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

F37 Substituted by Commission Implementing Regulation (EU) 2018/1132 of 13 August 2018 authorising the change of the designation and specific labelling requirement of the novel food synthetic zeaxanthin under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

Authorised Novel Food	Specifications
	Description
<i>N</i> -Acetyl-D- neuraminic acid	Description: N A satul D nouraminia said is a white to off white areatelling nourder
neuraminic aciu	<i>N</i> -Acetyl-D-neuraminic acid is a white to off-white crystalline powder Definition:
	Chemical name:
	IUPAC names:
	N -Acetyl-D-neuraminic acid (dihydrate)
	5-Acetamido-3,5-dideoxy-D-glycero-D-galacto-non-2-ulopyranosonic
	acid (dihydrate)
	Synonyms:
	Sialic acid (dihydrate)
	Chemical formula:
	$C_{11} H_{19} NO_9$ (acid)
	$C_{11}H_{23}NO_{11}$ (C $_{11}H_{19}NO_{9}$ * 2H $_{2}O$) (dihydrate)
	Molecular mass:
	309,3 Da (acid)
	345,3 (309,3 + 36,0) (dihydrate)
	CAS No.:
	131-48-6 (free acid)
	50795-27-2 (dihydrate)
	Specifications:
	Description: white to off-white crystalline powder
	pH (20 °C, 5 % solution): $1,7 - 2,5$
	N -Acetyl-D-neuraminic acid (dihydrate): $> 97,0\%$
	Water (dihydrate calculates to $10,4$ %): $\leq 12,5$ % (w/w)
	Ash, sulphated: $< 0.2 \%$ (w/w)
	Acetic acid (as free acid and/or sodium acetate): $< 0.5 \%$ (w/w)
	Heavy Metals:
	Iron: $< 20,0 \text{ mg/kg}$
	Lead: $< 0.1 \text{ mg/kg}$
	Residual proteins: < 0,01 % (w/w) Residual solvents:
	2-Propanol: < 0,1 % (w/w)
	Acetone: $< 0,1 \% (w/w)$
	Ethyl acetate: $< 0,1 \% (w/w)$
	Microbiological criteria:
	Salmonella: Absence in 25 g
	Aerobic mesophilic total count:< 500 CFU/g
	Enterobacteriaceae: Absence in 10 g
	Cronobacter (Enterobacter) sakazakii: Absence in 10 g
	Listeria monocytogenes: Absence in 25 g
	Bacillus cereus: < 50 CFU/g
	Yeasts: < 10 CFU/g

TABLE 2: SPECIFICATIONS

	Moulds: < 10 CFU/g Residual endotoxins: < 10 EU/mg CFU: Colony Forming Units; EU: Endotoxin Units.
<i>Adansonia digitata</i> (Baobab) dried fruit pulp	Description/Definition: The Baobab (<i>Adansonia digitata</i>) fruits are harvested from trees. The hard shells are cracked open and the pulp is separated from the seeds and the shell. This is milled, separated into coarse and fine lots (particle size 3 to 600 μ) and then packaged. Typical nutritional components: Moisture (loss on drying) (g/100 g): 4,5-13,7 Protein (g/100 g): 1,8-9,3 Fat (g/100 g): 0-1,6 Total carbohydrate (g/100 g): 76,3-89,5 Total sugars (as glucose): 15,2-36,5 Sodium (mg/100 g): 0,1-25,2 Analytical specifications: Foreign matter: Not more than 0,2 % Moisture (loss on drying) (g/100 g): 4,5-13,7 Ash (g/100 g): 3,8-6,6
<i>Ajuga reptans</i> extract from cell cultures	Description/Definition: Hydroalcoholic extract from <i>Ajuga reptans</i> L. tissue cultures which is substantially equivalent to extracts from flowering aerial parts of <i>Ajuga reptans</i> obtained by traditional cultures.
L-Alanyl-L- Glutamine	Description/Definition: L-Alanyl-L-Glutamine is produced by fermentation with a genetically modified strain of <i>Escherichia coli</i> . During the fermentation process, the ingredient is secreted into the growth medium from which it is subsequently separated and purified to a concentration of > 98 %. Appearance: White crystalline powder Purity: > 98 % Infrared spectroscopy: Conformity with ref. standard Appearance of solution: Colourless and clear Assay (dry basis): 98-102 % Related substances (each): $\leq 0,2$ % Residue on ignition: $\leq 0,1$ % Loss on drying: $\leq 0,5$ % Optical rotation: +9,0 - +11,0 ° pH (1 %; H ₂ O): 5,0-6,0 Ammonium (NH ₄): $\leq 0,020$ % Chloride (Cl): $\leq 0,020$ % Sulphate (SO ₄): $\leq 0,020$ % Microbiological criteria: <i>Escherichia coli</i> : Absence/g
Algal oil from the microalgae <i>Ulkenia</i> sp.	Description/Definition: Oil from the micro-algae <i>Ulkenia</i> sp. Acid value: $\leq 0,5$ mg KOH/g Peroxide value (PV): $\leq 5,0$ meq/kg oil Moisture and volatiles: $\leq 0,05$ % Unsaponifiables: $\leq 4,5$ %

DHA content: \geq 32 %

[^{F9} Allanblackia seed oil	Description/Definition: Allanblackia seed oil is obtained from the seeds of the allanblackia species: A. floribunda (synonymous with A. parviflora) and A. stuhlmannii . Composition of fatty acids (as a % of the total fatty acids): Lauric acid — Myristic acid — Palmitic acid (C12:0 – C14:0 – C16:0): sum of these acids < 4,0 % Stearic acid (C18:0): 45-58 % Oleic acid (C18:1): 40-51 % Poly unsaturated fatty acids (PUFA): < 2 % Characteristics: Free fatty acids: max 0,1 % of total fatty acids Trans fatty acids: max 1,0 % of total fatty acids Peroxide value: max 1,0 meq/kg Unsaponifiable matter: max 1,0 % (w/w) of the oil Saponification value: 185-198 mg KOH/g]
<i>Aloe macroclada</i> Baker leaf extract	Description/Definition: Powdered gel extract derived from the leaves of <i>Aloe macroclada</i> Baker which is substantially equivalent to the same gel derived from <i>Aloe vera</i> (L.) Burm.f. leaves. Ash: 25 % Dietary fibres: 28,6 % Fat: 2,7 % Moisture: 4,7 % Polysaccharides: 9,5 % Protein: 1,63 % Glucose: 8,9 %
[^{F39} Antarctic Krill oil from <i>Euphausia</i> <i>superba</i>	Description/Definition: To produce lipid extract from Antarctic Krill (<i>Euphausia superba</i>) deep- frozen crushed krill or dried krill meal is subjected to lipid extraction with an approved extraction solvent (under Directive 2009/32/EC). Proteins and krill material are removed from the lipid extract by filtration. The extraction solvents and residual water are removed by evaporation. Saponification value: $\leq 230 \text{ mg KOH/g}$ Peroxide value (PV): $\leq 3 \text{ meq O}_2/\text{kg oil}$ Oxidative stability: All food products containing Antarctic Krill oil from <i>Euphausia superba</i> should demonstrate oxidative stability by appropriate and recognised national/international test methodology (e.g. AOAC). Moisture and volatiles: $\leq 3 \%$ or 0,6 expressed as water activity at 25 °C Phospholipids: $\geq 35 \%$ to $< 60 \%$ Trans-fatty acids: $\leq 1 \%$ EPA (eicosapentaenoic acid): $\geq 9 \%$ DHA (docosahexaenoic acid): $\geq 5 \%$]
Antarctic Krill oil rich in phospholipids from <i>Euphausia</i> superba	Description/Definition: Oil rich in phospholipids is produced from Antarctic krill (<i>Euphausia superba</i>) by repeated solvent washings with an approved solvent (under Directive 2009/32/EC) to increase phospholipid content of the oil. Solvents are removed from the final product by evaporation. Saponification value: \leq 230 mg KOH/g

Status: Point in time view as at 31/12/2020.
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

Arachidonic acid-rich oil	Peroxide value (PV): $\leq 3 \mod O_2/kg$ oil Moisture and volatiles: $\leq 3 \%$ or 0,6 expressed as water activity at 25 °C Phospholipids: $\geq 60 \%$ Trans-fatty acids: $\leq 1 \%$ EPA (eicosapentaenoic acid): $\geq 9 \%$ DHA (docosahexaenoic acid): $\geq 5 \%$ Description/Definition: The clear yellow arachidonic acid-rich oil is obtained by fermentation
from the fungus Mortierella alpina	of the non-genetically modified strains IS-4, I49-N18, FJRK-MA01 and CBS 210.32 of the fungus <i>Mortierella alpina</i> using a suitable liquid. The oil is then extracted from the biomass and purified. Arachidonic acid: ≥ 40 % by weight of the total fatty acid content Free fatty acids: $\leq 0,45$ % of the total fatty acid content Trans fatty acids: $\leq 0,5$ % of the total fatty acid content Unsaponifiable matter: $\leq 1,5$ % Peroxide value (PV): ≤ 5 meq/kg Anisidin value: ≤ 20 Acid value: $\leq 1,0$ KOH/g Moisture: $\leq 0,5$ %
Argan oil from <i>Argania spinosa</i>	Description/Definition: Argan oil is the oil obtained by cold pressing of the almond like kernels of the fruits of <i>Argania spinosa</i> (L.) Skeels. Kernels may be roasted prior to pressing, but with no direct contact with a flame. Composition: Palmitic acid (C16:0): 12-15 % Stearic acid (C18:0): 5-7 % Oleic acid (C18:1): 43-50 % Linoleic acid (C18:2): 29-36 % Unsaponifiable matter: 0,3-2 % Total sterols: 100-500 mg/100 g Total tocopherols: 16-90 mg/100 g Oleic acidity: 0,2-1,5 % Peroxide value (PV): < 10 meq O ₂ /kg
Astaxanthin- rich oleoresin from <i>Haematococcus</i> <i>pluvialis</i> algae	Description/Definition: Astaxanthin is a carotenoid produced by <i>Haematococcus pluvialis</i> algae. Production methods for the growth of the algae are variable; using either closed systems exposed to sunlight or strictly controlled illuminated light; alternatively open ponds may be used. The algal cells are harvested and dried; the oleoresin is extracted using either super critical CO ₂ or a solvent (ethyl acetate). The Astaxanthin is diluted and standardized to 2,5 %, 5,0 %, 7,0 %, 10 %, 15 % or 20 % using olive oil, safflower oil, Sunflower oil or MCT (Medium Chain Triglycerides). Composition of the Oleoresin: Fat: 42,2- 99 % Protein: 0,3-4,4 % Carbohydrate: 0-52,8 % Fibre: < 1,0 % Ash: 0,0-4,2 % Specification of Carotenoids w/w% Total Astaxanthins: 2,9-11,1 % 9-cis-astaxanthin: 0,3-17,3 % 13-cis-astaxanthin: 0,2-7,0 % Astaxanthin monoesters: 79,8-91,5 %

<i>Status:</i> Point in time view as at 31/12/2020.
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Astaxanthin diesters: 0,16-19,0 % B-Carotene: 0,01-0,3 % Lutein: 0-1,8 % Canthaxanthin: 0-1,30 % Microbiological criteria: Total aerobic bacteria: < 3 000 CFU/g Yeast and Moulds: < 100 CFU/g Coliforms: < 10 CFU/g <i>E. coli</i> : Negative <i>Salmonella</i> : Negative <i>Staphylococcus</i> : Negative
Basil seeds (Ocimum basilicum)	Description/Definition: Basil (<i>Ocimum basilicum</i> L.) belongs to the family ' <i>Lamiaceae</i> ' within the order 'Lamiales'. Post-harvest the seeds are cleaned mechanically. Flowers, leaves and other parts of the plant are removed. Highest level of purity of Basil seeds has to be ensured by filtering (optical, mechanical). Production process of fruit juice and fruit/vegetable blend beverages containing Basil seeds (<i>Ocimum basilicum</i> L.) includes seed pre-hydration and pasteurisation steps. Microbiological controls and monitoring systems are in place. Dry Matter: 94,1 % Protein: 20,7 % Fat: 24,4 % Carbohydrate: 1,7 % Dietary Fibre: 40,5 % (Method: AOAC 958,29) Ash: 6,78 %
[^{F4} Betaine	Description/Definition: Betaine (N,N,N-trimethylglycine or carboxy-N,N,N- trimethylmethanaminium), in anhydrous (CH $_3$) $_3$ N $^+$ CH $_2$ COO $^-$ (CAS No: 107-43-7) and monohydrate (CH $_3$) 3N $^+$ CH $_2$ COO $^-$ (CAS No: 590-47-6) forms is obtained from processing of sugar beets (i.e. molasses, vinasses or betaine-glycerol). Characteristics/Composition Appearance: Free-flowing white crystals Betaine: \geq 99.0 % (w/w on dry weight basis) Moisture: \leq 2,0 % (anhydrous); \leq 15,0 % (monohydrate) Ash: \leq 0,1 % pH: 5,0-7,0 Residual protein: \leq 1,0 mg/g Heavy metals: Arsenic: < 0,1 mg/kg Mercury: < 0,005 mg/kg Cadmium: < 0,01 mg/kg Lead: < 0,05 mg/kg Microbiological criteria: Total viable count: \leq 100 CFU/g Coliforms: Negative/10 g Salmonella sp.: Negative/25 g Yeast: \leq 10 CFU/g CFU: Colony Forming Units.]

_

Document Generated: 2024-06-27

Status: Point in time view as at 31/12/2020.

Fermented black bean extract	Description/Definition: Fermented black bean extract (Touchi extract) is a fine light-brown protein-rich powder obtained by water extraction of small soybeans (<i>Glycine max (L.) Merr.</i>) fermented with <i>Aspergillus oryzae</i> . The extract contains an α -glucosidase inhibitor. Characteristics: Fat: $\leq 1,0 \%$ Protein: $\geq 55 \%$ Water: $\leq 7,0 \%$ Ash: $\leq 10 \%$ Carbohydrate: $\geq 20 \%$ α -glucosidase inhibitory activity: IC50 min 0,025 mg/ml Soy isoflavone: $\leq 0,3 \text{ g}/100 \text{ g}$
Bovine lactoferrin	Description/Definition:Bovine lactoferrin is a protein that occurs naturally in cows' milk. It is an iron-binding glycoprotein of approximately 77 kDa and consists of a single polypeptide chain of 689 amino acids.Production process: Bovine lactoferrin is isolated from skimmed milk or cheese whey via ion exchange and subsequent ultra-filtration steps.Finally, it is dried by freeze drying or spraying and the large particles are sieved out. It is a virtually odourless, light pinkish powder.Physical-Chemical properties of Bovine lactoferrin: Moisture: < 4,5 % Ash: < 1,5 %
[^{F10} Bovine milk basic whey protein isolate	Description Bovine milk basic whey protein isolate is a yellowish grey powder obtained from bovine skimmed milk via a series of isolation and purification steps. Characteristics/Composition Total protein (w/weight of product): ≥ 90 % Lactoferrin (w/weight of product): $25-75$ % Lactoperoxidase (w/weight of product): $10-40$ % Other proteins (w/weight of product): ≤ 30 % TGF- β 2: 12-18 mg/100 g Moisture: $\leq 6,0$ % pH (5 % solution w/v): $5,5 - 7,6$ Lactose: $\leq 3,0$ % Fat: $\leq 4,5$ % Ash: $\leq 3,5$ % Iron: ≤ 25 mg/100 g Heavy Metals Lead: < 0,1 mg/kg Cadmium: < 0,2 mg/kg Mercury: < 0,6 mg/kg Arsenic: < 0,1 mg/kg

Status: Point in time view as at 31/12/2020.	
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX	is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brough	t into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (S	See end of Document for details)

	Microbiological criteria:Aerobic mesophilic count: $\leq 10\ 000\ CFU/g$ Enterobacteriaceae : $\leq 10\ CFU/g$ Escherichia coli : Negative/gCoagulase positive Staphylococci: Negative/gSalmonella : Negative/25 gListeria : Negative/25 gCronobacter spp.: Negative/25 gMoulds: $\leq 50\ CFU/g$ Yeasts: $\leq 50\ CFU/g$ CFU: Colony Forming Units]
<i>Buglossoides arvensis</i> seed oil	Description/Definition: Refined Buglossoides oil is extracted from the seeds of <i>Buglossoides</i> <i>arvensis</i> (L.) I.M.Johnst Alpha-linolenic acid: ≥ 35 % w/w of total fatty acids Stearidonic acid: ≥ 15 % w/w of total fatty acids Linoleic acid: $\geq 8,0$ % w/w of total fatty acids Trans fatty acids: $\leq 2,0$ % w/w of total fatty acids Acid value: $\leq 0,6$ mg KOH/g Peroxide value (PV): $\leq 5,0$ meq O ₂ /kg Unsaponifiable content: $\leq 2,0$ % Protein content (total nitrogen): ≤ 10 µg/ml Pyrrolizidine alkaloids: Not detectable with a detection limit of 4,0 µg/kg
<i>Calanus finmarchicus</i> oil	Description/Definition: The novel food is ruby coloured, slightly viscous oil with a slight shellfish odour extracted from the crustacean (marine zooplankton) <i>Calanus finmarchicus</i> . The ingredient consists primarily of wax esters (> 85 %) with minor amounts of triglycerides and other neutral lipids. Specifications: Water: < 1,0 % Wax esters: > 85 % Total fatty acids: > 46 % Eicosapentaenoic acid (EPA): > 3,0 % Docosahexaenoic acid (DHA): > 4,0 % Total fatty alcohols: > 28 % C20:1 n-9 fatty alcohol: > 9,0 % C22:1 n-11 fatty alcohol: > 12 % Trans fatty acids: < 1,0 % Astaxanthinesters: < 0,1 % Peroxide value (PV): < 3,0 meq. O ₂ /kg
Chewing gum base (monomethoxypo glycol)	Description/Definition: The novel food ingredient is a synthetic polymer (Patent hypethydeWO2006016179). It consists of branched polymers of monomethoxypolyethylene glycol (MPEG) grafted onto polyisoprene- graft-maleic anhydride (PIP-g-MA), and unreacted MPEG (less than 35 % by weight). White to off-white colour. CAS No.: 1246080-53-4 Characteristics: Moisture: < 5,0 % Aluminium: < 3,0 mg/kg

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all c	hanges
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future	date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for	r details)

	Lithium: < 0,5 mg/kg Nickel: < 0,5 mg/kg Residual anhydride: < 15 μ mol/g Polydispersity index: < 1,4 Isoprene: < 0,05 mg/kg Ethylene oxide: < 0,2 mg/kg Free maleic anhydride: < 0,1 %Total oligomeres (less than 1 000 Dalton): \leq 50 mg/kg Ethylene glycol: < 200 mg/kg Diethylene glycol: < 30 mg/kg Monoethylene glycol methyl ether: < 3,0 mg/kg Diethylene glycol methyl ether: < 4,0 mg/kg Triethylene glycol methyl ether: < 7,0 mg/kg I,4-Dioxane: < 2,0 mg/kg
Chewing gum base (Methyl vinyl ether-maleic anhydride copolymer)	Description/Definition: Methyl vinyl ether-maleic anhydride copolymer is an anhydrous copolymer of methyl vinyl ether and maleic anhydride.Free-flowing, white to white-off powder CAS No: 9011-16-9Purity: Assay value: At least 99,5 % in dry matter Specific viscosity (1 % MEK): 2-10 Residual methyl vinyl ether: ≤ 150 ppm Residual maleic anhydride: ≤ 250 ppm Acetaldehyde: ≤ 500 ppm Dilauroyl peroxide: ≤ 15 ppm Total heavy metals: ≤ 10 ppm Microbiological criteria: Total aerobic plate count: ≤ 500 CFU/g Mould/yeast: ≤ 500 CFU/g Escherichia coli : Negative to test Salmonella : Negative to test Pseudomonas aeruginosa : Negative to test
Chia oil from <i>Salvia hispanica</i>	Description/Definition: Chia oil is produced from Chia (<i>Salvia hispanica</i> L.) seeds (99,9 % pure) by cold pressing. No solvents are used and, once pressed, the oil is held in decantation tanks and a filtration process employed to remove impurities. It can also be produced by extraction with supercritical CO ₂ . Production process: Produced by cold pressing. No solvents are used and, once pressed, the oil is held in decantation tanks and a filtration process employed to remove impurities. Acidity expressed as oleic acid: $\leq 2,0$ % Peroxide value (PV): ≤ 10 meq/kg Insoluble impurities: $\leq 0,05$ % Alpha linolenic acid: ≥ 60 % Linoleic acid: 15-20 %

Chia seeds (<i>Salvia hispanica</i>)	Description/Definition: Chia (<i>Salvia hispanica</i> L.) is a summer annual herbaceous plant belonging to the <i>Labiatae</i> family. Post-harvest the seeds are cleaned mechanically. Flowers, leaves and other parts of the plant are removed. Dry matter: 90-97 % Protein: 15-26 % Fat: 18-39 % Carbohydrate (*): 18-43 % Crude Fibre(**): 18-43 %
	(*) Carbohydrates include the fibre value
	 (**) Crude fibre is the part of fibre made mainly of indigestible cellulose, pentosans and lignin Production process: Production process of fruit juices and fruit juice blends beverages, containing Chia seeds, includes seed pre-hydration and pasteurisation steps. Microbiological controls and monitoring systems are in place.
Chitin- glucan from <i>Aspergillus</i> <i>niger</i>	 Description/Definition: Chitin-glucan is obtained from the mycelium of Aspergillus niger ; it is a slightly yellow, odourless, free-flowing powder. It has a dry matter content of 90 % or more. Chitin-glucan is composed largely of two polysaccharides: chitin, composed of repeating units of N-acetyl-D-glucosamine (CAS No: 1398-61-4), beta (1, 3)-glucan, composed of repeating units of D-glucose (CAS No: 9041-22-9). Loss on drying: ≤ 10 % Chitin-glucan: ≥ 90 % Ratio of chitin to glucan: 30:70 to 60:40 Ash: ≤ 3,0 % Lipids: ≤ 1,0 % Proteins: ≤ 6,0 %
Chitin-glucan complex from Fomes fomentarius	 Description/Definition: Chitin-glucan complex is obtained from the cell walls of the fruit bodies of the fungus <i>Fomes fomentarius</i>. It consists primarily of two polysaccharides: Chitin, composed of repeating units of N-acetyl-D-glucosamine (CAS No: 1398-61-4); Beta-(1,3)(1,6)-D-glucan, composed of repeating units of D-glucose (CAS No: 9041-22-9). The production process consists of several steps, including: cleaning, reduction in size and grinding, softening in water and heating in an alkaline solution, washing, drying. No hydrolysis is applied during the production process. Appearance: Powder, odourless, flavourless, brown Purity: Moisture: ≤ 15 % Ash: ≤ 3,0 % Chitin-glucan: ≥ 90 % Ratio of chitin to glucan: 70:20

	Total carbohydrates, excluding glucans: $\leq 0,1$ % Proteins: $\leq 2,0$ % Lipids: $\leq 1,0$ % Melanins: $\leq 8,3$ % Additives: None pH: 6,7-7,5 Heavy metals: Lead (ppm): $\leq 1,00$ Cadmium (ppm): $\leq 1,00$ Mercury (ppm): $\leq 0,03$ Arsenic (ppm): $\leq 0,20$ Microbiological criteria: Total mesophilic bacteria: $\leq 10^{3}/g$ Yeast and moulds: $\leq 10^{3}/g$
	Coliforms at 30 °C: $\leq 10^3$ /g E. coli : $\leq 10/g$
	Salmonella and other pathogenic bacteria: Absence/25 g
Chitosan extract from fungi (<i>Agaricus</i> <i>bisporus</i> ; <i>Aspergillus</i> <i>niger</i>)	Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of <i>Agaricus bisporus</i> or from the mycelium of <i>Aspergillus niger</i> . The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: (C ₆ H ₁₁ NO ₄) _n Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight):≥ 85 Glucan content (% w/w dry weight):≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i> Ash (% w/w dry weight): ≤ 3,0 Proteins (% w/w dry weight): ≤ 2,0 Particle size: > 100 nm Tapped density (g/cm ³): 0,7-1,0 Fat binding capacity 800 × (w/w wet weight): pass Heavy metals: Mercury (ppm): ≤ 0,1 Lead (ppm): ≤ 1,0 Cadmium (ppm): ≤ 0,5 Microbiological criteria: Aerobic count (CFU/g): ≤ 10 ³ Yeast and mould count (CFU/g): ≤ 10 ³ Escherichia coli (CFU/g): ≤ 10

	Enterobacteriaceae (CFU/g): ≤ 10 Salmonella : Absence/25g Listeria monocytogenes : Absence/25g
Chondroitin sulphate	Description/Definition: Chondroitin sulphate (sodium salt) is a biosynthetic product. It is obtained by chemical sulphation of chondroitin derived from fermentation by the bacterium <i>Escherichia coli</i> O5:K4:H4 strain U1-41 (ATCC 23502). Chondroitin sulphate (sodium salt) (% dry basis): 95-105 MWw (weight avg.) (kDa): 5-12 MWn (number avg.) (kDa): 4-11 Dispersity ($w_h/w_{0.05}$): $\leq 0,7$ Sulphation pattern (Δ Di-6S) (%): ≤ 85 Loss on drying (%) (105 °C to constant weight): $\leq 10,0$ Residue on ignition (% dry basis): 20-30 Protein (% dry basis): $\leq 0,5$ Endotoxins (EU/mg): ≤ 100 Total organic impurities (mg/kg): ≤ 50
Chromium Picolinate	Description/Definition:Chromium picolinate is a reddish free-flowing powder, slightly soluble in water at pH 7. The salt is also soluble in polar organic solvents.Chemical name: tris(2pyridinecarboxylato-N,O)chromium(III) or 2- pyridinecarboxylic acid chromium(III) saltCAS No.: 14639-25-9Chemical formula: $Cr(C_6 H_4 NO_2)_3$ Chemical characteristics: Chromium Picolinate: $\geq 95 \%$ Chromium (III): 12-13 % Chromium (VI): not detected Water: $\leq 4,0 \%$
<i>Cistus incanus</i> L. Pandalis herb	Description: <i>Cistus incanus</i> L. Pandalis herb; species belonging to the <i>Cistaceae</i> family and native to the Mediterranean region, Chalkidiki Peninsula. Composition: Moisture: $9-10 \text{ g}/100 \text{ g}$ herbs Protein: 6,1 g/100 g herbs Fat: 1,6 g/100 g herbs Carbohydrates: 50,1 g/100 g herbs Fiber: 27,1 g/100 g herbs Minerals: 4,4 g/100 g herbs Sodium: 0,18 g Potassium: 0,75 g Magnesium: 0,24 g Calcium: 1,0 g Iron: 65 mg Vitamin B ₁ : 3,0 µg Vitamin B ₂ : 30 µg Vitamin B ₆ : 54 µg Vitamin A: less than 0,1 mg Vitamin E: 40–50 mg Alpha-Tocopherol: 20–50 mg

	Beta and Gamma-Tocopherols: 2–15 mg Delta-Tocopherol: 0,1–2 mg
Citicoline	Description/Definition:Citicoline is produced by a microbial process.Citicoline is composed of cytosine, ribose, pyrophosphate and choline.White crystalline powderChemical name: Choline cytidine 5'-pyrophosphate, Cytidine 5'-(trihydrogen diphosphate) P'-[2-(trimethylammonio)ethyl]ester inner saltChemical formula: C $_{14}$ H $_{26}$ N $_{4}$ O $_{11}$ P $_{2}$ Molecular weight: 488,32 g/molCAS No.: 987-78-0pH (sample solution of 1 %): 2,5-3,5Purity:Assay value: \geq 98 % of dry matterLoss on drying (100 °C for 4 hours): \leq 5,0 %Ammonium: \leq 0,05 %Arsenic: Not more than 2 ppmFree phosphoric acids: \leq 0,1 %S'-Cytidylic acid: \leq 1,0 %Microbiological criteria:Total plate count: \leq 10 3 CFU/gYeast and moulds: \leq 10 2 CFU/g <i>Escherichia coli</i> : Absence in 1 g
Clostridium butyricum	Description/Definition: Clostridium butyricum (CBM-588) is a Gram-positive, spore-forming, obligate anaerobic, non-pathogenic, non-genetically modified bacterium. Depository number FERM BP-2789 Microbiological criteria: Total viable aerobic count: $\leq 10^3$ CFU/g Escherichia coli : Not detected in 1 g Staphylococcus aureus : Not
[^{F8} D-ribose	Description D-ribose is an aldopentose monosaccharide which is produced by fermentation using a transketolase-deficient strain of <i>Bacillus subtilis</i> . Chemical formula: $C_5 H_{10} O_5$ CAS No: 50-69-1 Molecular mass: 150,13 Da Characteristics/Composition Appearance: Dry with powdery texture, white to slightly yellow in colour Specific rotation $[\alpha]_D^{25}$: – 19,0° to – 21,0° D-ribose purity (% dry basis): -HPLC/RI ^h Method 98,0–102,0 % Ash: < 0,2 % Loss on drying (moisture): < 0,5 % Clarity on solution: \ge 95 % transmittance Heavy metals Lead: \le 0,1 mg/kg Arsenic: \le 0,1 mg/kg

	Cadmium: $\leq 0,1$ mg/kg Mercury: $\leq 0,1$ mg/kg Microbiological criteria Total plate count: ≤ 100 CFU ⁱ /g Yeast: ≤ 100 CFU/g Moulds: ≤ 100 CFU/g Coliforms: ≤ 10 CFU/g Salmonella sp: Negative/25 g]
Extract of defatted cocoa powder	Cocoa (<i>Theobroma cacao</i> L.) Extract Appearance: Dark brown powder free of visible impurities Physical and chemical properties: Polyphenol content: Min 55,0 % GAE Theobromine content: Max 10,0 % Ash content: Max 5,0 % Moisture content: Max 8,0 % Bulk density: 0,40-0,55 g/cm ³ pH: 5,0-6,5 Residual solvent: Max 500 ppm
Low fat cocoa extract	Low fat Cocoa (<i>Theobroma cacao</i> L.) extract Appearance: Dark red to purple powder Cocoa extract, concentrate: Min 99 % Silicon dioxide (technological aid): Max 1,0 % Cocoa flavanols: Min. 300 mg/g — Epicatechin: Min. 45 mg/g Loss on drying: Max. 5,0 %
[^{F40} Coriander seed oil from <i>Coriandrum</i> <i>sativum</i>	Description/Definition: Coriander seed oil is an oil containing glycerides of fatty acids that is produced from the seeds of the coriander plant <i>Coriandrum sativum</i> L. Slight yellow colour, bland taste CAS No: 8008-52-4 Composition of fatty acids: Palmitic acid (C16:0): 2-5 % Stearic acid (C18:0): $< 1,5 \%$ Petroselinic acid (cis-C18:1(n-12)): 60-75 % Oleic acid (cis-C18:1 (n-9)): 7-15 % Linoleic acid (C18:2): 12-19 % α -Linolenic acid (C18:3): $< 1,0 \%$ Purity: Refractive index (20 °C): 1,466-1,474 Acid value: $\le 2,5$ mg KOH/g Peroxide value (PV): $\le 5,0$ meq/kg Iodine value: 88-110 units Saponification value: 179-200 mg KOH/g Unsaponifiable matter: $\le 15 g/kg$]
[^{F12} Cranberry extract powder	Description/Definition: Cranberry extract powder is a water-soluble phenolic-rich powder extract prepared through an ethanolic extraction from the juice concentrate of

	Moisture (% w/w): ≤ 4 Proanthocyanidins — PACs (% w/w dry weight) — OSC-DMAC method ^{ee} : 55.0-60.0 or — BL-DMAC method ^{de} : 15.0-18.0 Total phenolics (GAE ^f , % w/w dry weight) ^e — Folin-Ciocalteau method: > 46.2 Solubility (water): 100 %, with no visible insoluble particles Ethanol Content (mg/kg): ≤ 100 Screen Analysis: 100 % through 30 mesh screen Appearance and aroma, as powder: Free-flowing, deep red colour. Earthy aroma with no burnt character. Heavy metals: Arsenic (ppm): ≤ 3 Microbiological criteria: Yeast: ≤ 100 CFU ^g /g Mould: ≤ 100 CFU/g Aerobic plate count: ≤ 1 000 CFU/g Coliforms: ≤ 10 CFU/g <i>Escherichia coli</i> : ≤ 10 CFU/g <i>Salmonella</i> : Absent in 375 g]
<i>Crataegus pinnatifida</i> dried fruit	Description/Definition:Dried fruits of Crataegus pinnatifida species belonging to the Rosaceaefamily and native to north China and Korea.Composition:Dry matter: 80 %Carbohydrates: 55 g/kg fresh weightFructose: 26,5–29,3 g/100 gGlucose: 25,5–28,1 g/100 gVitamin C: 29,1 mg/100 g fresh weightSodium: 2,9 g/100 g fresh weightCompotes are products obtained by thermal processing of the ediblepart of one or several species of fruits, whole or in pieces, sieved or not,without significant concentration. Sugars, water, cider, spices and lemonjuice may be used.
a-cyclodextrin	Description/Definition: A non-reducing cyclic saccharide consisting of six α -1,4-linked D-glucopyranosyl units produced by the action of cyclodextrin glucosyltransferase (CGTase, EC 2.4.1.19) on hydrolyzed starch. Recovery and purification of α -cyclodextrin may be carried out using one of the following procedures: precipitation of a complex of α -cyclodextrin with 1-decanol, dissolution in water at elevated temperature and re- precipitation, steam-stripping of thecomplexant, and crystallisation of α - cyclodextrin from the solution; or chromatography with ion-exchange or gel filtration followed by crystallisation of α -cyclodextrin from the purified mother liquor; or membrane separation methods such as ultra- filtration and reverse osmosis: Description: Virtually odourless, white or almost white crystalline solid. Synonyms: α -cyclodextrin, α -dextrin, cyclohexaamylose, cyclomaltohexaose, α -cycloamylase Chemical name: CyclohexaamyloseCAS No.: 10016-20-3 Chemical formula: (C $_6$ H $_{10}$ O $_5$) $_6$

	chromatogram of the sample corresponds to that for α -cyclodextrin in a
	chromatogram of reference α-cyclodextrin (available from <i>Consortium für Elektrochemische Industrie GmbH, München, Germany or Wacker Biochem Group, Adrian, MI, USA</i>) using the conditions described in the
	METHOD OF ASSAY Purity:
	Water: ≤ 11 % (Karl Fischer Method)
	Residual complexant: $\leq 20 \text{ mg/kg}$
	(1-decanol)
	Reducing substances: ≤ 0.5 % (as glucose)
	Sulphated ash: $\leq 0,1 \%$ Lead: $\leq 0,5 \text{ mg/kg}$
	Method of assay:
	Determine by liquid chromatography using the following conditions:
	Sample solution: Weigh accurately about 100 mg of test sample into a 10
	ml volumetric flask and add 8 ml of deionised water. Dissolve the sample
	completely using an ultra-sonification bath (10-15 min) and dilute to the mark with purified deionised water. Filter through a 0,45-micrometer
	filter
	Reference solution: Weigh accurately about 100 mg of α -cyclodextrin into a 10 ml volumetric flask and add 8 ml of deionised water. Dissolve the sample completely using an ultra-sonification bath and dilute to the mark with purified deionised water.
	Chromatography: Liquid chromatograph equipped with a refractive index
	detector and an integrating recorder. Column and packing: Nucleosil-100-NH ₂ (10 μm) (<i>Macherey & Nagel</i>
	$($ Column and nacking: Nucleosil-100-NH $_{2}$ (10 µm) (Macherev & Nagel
	Co. Düren, Germany) or similar
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v)
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min
	Co. Düren, Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows:
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S)
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where A _S and A _R are the areas of the peaks due to α -cyclodextrin for the
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where A _S and A _R are the areas of the peaks due to α -cyclodextrin for the sample solution and reference solution, respectively.
	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where A _S and A _R are the areas of the peaks due to α -cyclodextrin for the
γ-cyclodextrin	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where A _S and A _R are the areas of the peaks due to α -cyclodextrin for the sample solution and reference solution, respectively. W _S and W _R are the weights (mg) of the test sample and reference α - cyclodextrin, respectively, after correcting for water content.
γ-cyclodextrin	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where A _S and A _R are the areas of the peaks due to α -cyclodextrin for the sample solution and reference solution, respectively. W _S and W _R are the weights (mg) of the test sample and reference α -
γ-cyclodextrin	<i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µlProcedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α - CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where A _S and A _R are the areas of the peaks due to α -cyclodextrin for the sample solution and reference solution, respectively. W _S and W _R are the weights (mg) of the test sample and reference α - cyclodextrin, respectively, after correcting for water content. Description/Definition:

	glucosyltransferase (CGTase, EC 2.4.1.19) on hydrolysed starch. Recovery and purification of γ -cyclodextrin may be carried out by precipitation of a complex of γ -cyclodextrin with 8-cyclohexadecen-1- one, dissolution of the complex with water and n-decane, steam-stripping of the aqueous phase and recovery of gamma-CD from the solution by crystallisation. Virtually odourless, white or almost white crystalline solid Synonyms: γ -cyclodextrin, γ -dextrin, cyclooctaamylose, cyclomaltooctaose, γ -cycloamylase Chemical name: Cyclooctaamylose CAS number: 17465-86-0 Chemical formula: (C $_{6}$ H $_{10}$ O $_{5}$) $_{8}$ Assay: \geq 98 % (dry basis) Identification: Melting range: Decomposes above 285 °C Solubility: Freely soluble in water; very slightly soluble in ethanol Specific rotation: [α] $_{D}$ ²⁵ : between + 174 ° and + 180 ° (1 % solution) Purity: Water: \leq 11 % Residual complexant (8-cyclohexadecen-1-one (CHDC)): \leq 4 mg/kg Residual solvent (n-decane): \leq 6mg/kg Reducing substances: \leq 0,5 % (as glucose) Sulphated ash: \leq 0,1 %
[^{F13} Decorticated grains of <i>Digitaria exilis</i> (Kippist) Stapf (fonio) (Traditional food from a third country)	 Description/Definition The traditional food is the decorticated grain (bran removed) of <i>Digitaria exilis</i> (Kippist) Stapf. <i>Digitaria exilis</i> (Kippist) Stapf) is an annual herbaceous plant belonging to the <i>Poaceae</i> family. Typical nutritional components of decorticated grain of fonio Carbohydrates: 76,1 g/100 g of fonio Water: 12,4 g/100 g of fonio Protein: 6,9 g/100 g of fonio Fat: 1,2 g/100 g of fonio Fibre: 2,2 g/100 g of fonio Ash: 1,2 g/100 g of fonio
Dextran preparation produced by <i>Leuconostoc</i> <i>mesenteroides</i>	Phytate content: $\leq 2,1 \text{ mg/g}$ 1. Powdered form: Carbohydrates: 60 % with: (Dextran: 50 %, Mannitol: 0,5 %, Fructose: 0,3 %, Leucrose: 9,2 %) Protein: 6,5 % Lipid: 0,5 % Lactic acid: 10 % Ethanol: traces Ash: 13 % Moisture: 10 %2. Liquid form: Carbohydrates: 12 % with: (Dextran: 6,9 %, Mannitol: 1,1 %, Fructose: 1,9 %, Leucrose: 2,2 %) Protein: 2,0 % Lipid: 0,1 % Lactic acid: 2,0 %

	Ethanol: 0,5 % Ash: 3,4 % Moisture: 80 %
Diacylglycerol oil of plant origin	Description/Definition: Manufactured from glycerol and fatty acids derived from edible vegetable oils, in particular from soybean oil (<i>Glycine max</i>) or rapeseed oil (<i>Brassica campestris, Brassica napus</i>) using a specific enzyme. Acylglycerol Distribution: Diacylglycerols (DAG): \geq 80 % 1,3-Diacylglycerols (1,3-DAG): \geq 50 % Triacylglycerols (TAG): \leq 20 % Monoacylglycerols (MAG): \leq 5,0 % Fatty Acid Composition (MAG, DAG, TAG): Oleic acid (C18:1): 20-65 % Linoleic acid (C18:2): 15-65 % Linolenic acid (C18:3): \leq 15 % Saturated fatty acids: \leq 10 % Others: Acid value: \leq 0,5 mg KOH/g Moisture and volatile: \leq 0,1 % Peroxide value (PV): \leq 1,0 meq/kg Unsaponifiables: \leq 2,0 % MAG = monoacylglycerols, DAG = diacylglycerols, TAG = triacylglycerols
Dihydrocapsiate (DHC)	Description/Definition: Dihydrocapsiate is synthesised by enzyme-catalysed esterification of vanillyl alcohol and 8-methylnonanoic acid. Following the esterification dihydrocapsiate is extracted with n-hexane. Viscous to colourless to yellow liquid Chemical formula: C $_{18}$ H $_{28}$ O $_4$ CAS No: 205687-03-2 Physical-chemical properties: Dihydrocapsiate: > 94 % 8-Methylnonanoic acid: < 6,0 % Vanillyl acohol: < 1,0 % Other synthesis related substances: < 2,0 %
[^{F14} Dried aerial parts of <i>Hoodia</i> parviflora	Description/Definition: It is the whole dried aerial parts of <i>Hoodia parviflora</i> N.E.Br., (family <i>Apocynaceae</i>) Characteristics/Composition Plant material: Aerial parts of at least 3-year-old plants Appearance: Light green to tan fine powder Solubility (water): $> 25 \text{ mg/mL}$ Moisture: $< 5,5 \%$ $A_w: < 0,3$ pH: $< 5,0$ Protein: $< 4,5 \text{ g}/100 \text{ g}$ Fat: $< 3 \text{ g}/100 \text{ g}$ Carbohydrate (including dietary fibre): $< 80 \text{ g}/100 \text{ g}$ Dietary fibre: $< 55 \text{ g}/100 \text{ g}$

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all c	hanges
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future	date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for	r details)

	Total sugars: < 10,5 g/100 g Ash: < 20 % Hoodigosides P57: 5–50 mg/kg L: 1 000–6 000 mg/kg O: 500–5 000 mg/kg Total: 1 500–11 000 mg/kg Heavy metals: Arsenic: < 1,00 mg/kg Mercury: < 0,1 mg/kg Lead: < 0,5 mg/kg Microbiological criteria: Aerobic plate count: < 10 ⁵ CFU/g <i>Escherichia coli</i> : < 10 CFU/g <i>Staphylococcus aureus</i> : < 50 CFU/g Total coliforms: < 10 CFU/g Yeast: < 100 CFU/g Mould: < 100 CFU/g <i>Salmonella</i> species: Negative/25 g <i>Listeria monocytogenes</i> : Negative/25 g CFU: Colony Forming Units]
Dried extract of <i>Lippia</i> <i>citriodora</i> from cell cultures	Description/Definition: Dried extract of <i>Lippia citriodora</i> (Palau) Kunth from cell cultures HTN [®] Vb.
<i>Echinacea angustifolia</i> extract from cell cultures	Description/Definition: Extract of the roots of <i>Echinacea angustifolia</i> obtained from plant tissue culture which is substantially equivalent to a root extract from <i>Echinacea angustifolia</i> obtained in ethanol-water titrated to 4 % echinacoside.
[^{F15} Echinacea purpurea extract from cell cultures	Description/Definition: Dried extract of <i>Echinacea purpurea</i> from cell cultures EchiPure-PC [™]]
<i>Echium plantagineum</i> oil	Description/Definition: Echium oil is the pale yellow product obtained by refining oil extracted from the seeds of <i>Echium plantagineum</i> L. Stearidonic acid: $\geq 10 \%$ w/ w of total fatty acids Trans fatty acids: $\leq 2,0 \%$ (w/w of total fatty acids) Acid value: $\leq 0,6$ mg KOH/g Peroxide value (PV): $\leq 5,0$ meq O ₂ /kg Unsaponifiable content: $\leq 2,0 \%$ Protein content (total nitrogen): $\leq 20 \mu$ g/ml Pyrrolizidine alkaloids: Not detectable with a detection limit 4,0 µg/kg
[^{F16} Ecklonia cava phlorotannins	Description/Definition <i>Ecklonia cava</i> phlorotannins are obtained via alcohol extraction from the edible marine alga <i>Ecklonia cava</i> . The extract is a dark brown powder, rich in phlorotannins, polyphenolic compounds found as secondary metabolites in certain brown algae species. Characteristics/Composition

	Phlorotannin content: $90 \pm 5 \%$ Antioxidant activity: > 85 % Moisture: < 5 % Microbiological criteria Total viable cell count: < 3 000 CFU/g Mould/yeast: < 300 CFU/g Coliforms: Negative to test <i>Salmonella</i> spp.: Negative to test <i>Staphylococcus aureus:</i> Negative to test Heavy metals and Halogens Lead: < 3,0 mg/kg Mercury: < 0,1 mg/kg Cadmium: < 3,0 mg/kg Arsenic: < 25,0 mg/kg Inorganic Arsenic: < 0,5 mg/kg Iodine: 150,0 – 650,0 mg/kg CFU: Colony Forming Units]		
[^{F17} Egg membrane hydrolysate	Description The egg membrane hydrolysate is derived from the eggshell membranes of chicken eggs. The eggshells undergo hydro-mechanical separation in order to obtain the egg membranes, which are then further processed using a patented solubilisation method. Following the solubilisation process, the solution is filtered, concentrated, spray-dried and packaged. Characteristics/Composition		
	Chemical parameters	Methods	
	Total nitrogen- containing compounds (% w/w): ≥ 88	Combustion according to AOAC 990.03 and AOAC 992.15	
	Collagen (% w/ w): ≥ 15	Sircol TM Soluble Collagen Assay	
	Elastin (% w/	Fastin TM Elastin Assay	
	w): ≥ 20	rasun Elasun Assay	
		USP26 (chondroitin sulphate K0032 method)	
	w): ≥ 20 Total glycosaminoglyca	USP26 (chondroitin sulphate K0032 method)	

Status: Point in time view as at 31/12/2020.

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Microbiological criteria Aerobic plate count: ≤ 2500 CFU/g <i>Escherichia coli</i> : ≤ 5 MPN/g <i>Salmonella</i> : Negative (in 25 g) Coliforms: ≤ 10 MPN/g <i>Staphylococcus aureus</i> : ≤ 10 CFU/g Mesophilic spore count: ≤ 25 CFU/g Thermophilic spore count: ≤ 10 CFU/10 g Yeast: ≤ 10 CFU/g Mould: ≤ 200 CFU/g CFU: Colony Forming Units; MPN = Most Probable Number; USP: United States Pharmacopeia.]			
Epigallocatechin gallate as a purified extract from green tea leaves (<i>Camellia</i> <i>sinensis</i>)	 Description/Definition: A highly purified extract from the leaves of green tea (<i>Camellia sinensis</i> (<i>L.</i>) <i>Kuntze</i>) in the form of a fine, off-white to pale pink powder. It is composed of a minimum of 90 % epigallo-catechin gallate (EGCG), and has a melting point between approx. 210 and 215 °C Appearance: off-white to pale pink powder Chemical name: polyphenol (-) epigallocatechin-3-gallate Synonyms: epigallocatechin gallate (EGCG) CAS No.: 989-51-5 INCI name: epigallocatechin gallate Molecular mass: 458,4 g/mol Loss on drying: max 5,0 % Heavy metals: Arsenic: max 3,0 ppm Lead: max 5,0 ppm Assay: Min. 94 % EGCG (on dry material) max. 0,1 % caffeine Solubility: EGCG is fairly soluble in water, ethanol, methanol and 			
L- ergothioneine	yl)-2-(trimethylan	nmonio)-Propanoat a: C ₉ H ₁₅ N ₃ O ₂ S 229,3 Da	thioxo-2,3-dihydro- e <u><i>Method</i></u> Visual Polarimetry HPLC [Eur. Ph. 2,7 1H-NMR 1H-NMR Elemental analysis	2.29]

	Total residual solvents (methanol, ethyl acetate, isopropanol, ethanol) Loss on drying	[Eur. Ph. 01/2008:50400] < 1 000 ppm	Gas chromatography [Eur. Ph. 01/2008:20424] [Eur. Ph. 01/2008:20232]
		< 0,5 %	
	Impurities	< 0,8 %	HPLC/GPC or 1H-NMR
	Heavy metals ^{b) c}	[
	Lead	< 3,0 ppm	ICP/AES
	Cadmium	< 1,0 ppm	(Pb, Cd)
	Mercury	< 0,1 ppm	Atomic fluorescence (Hg)
	Microbiological s	specifications ^{b)}	
	Total viable aerobic count (TVAC)	$\leq 1 \times 10^{-3}$ CFU/g	[Eur. Ph. 01/2011:50104]
	Total yeast and mould count (TYMC)	\leq 1 x 10 ² CFU/g	
	Escherichia coli	Absence in 1 g	
	resonance; HPLCpermeation chromatomic emission sCFU: colony-forma)Lit. $[\alpha]_{IL}$ b)Analyse	: high-performance hatography; ICP/AF pectroscopy; hing units. $p = (+) 126,6^{\circ}$ (c = s conducted on eac m levels in accorda	
[^{F16} Extract of three herbal roots (<i>Cynanchum</i> wilfordii Hemsley, Phlomis umbrosa Turcz. and Angelica gigas Nakai)	Description/Definition The mixture of the three herbal roots is yellowish brown fine powder produced by hot-water extraction, concentration by evaporation, and spray drying Composition of the extract of mixture of the 3 herbal roots <i>Cynanchum wilfordii</i> root: $32,5 \%$ (w/w) <i>Phlomis umbrosa</i> root: $32,5 \%$ (w/w) <i>Angelica gigas</i> root: $35,0 \%$ (w/w) Specifications Loss on drying: NMT 100 mg/g Assay Cinnamic acid: $0,012 - 0,039$ mg/g Shanzhiside methyl ester: $0,20 - 1,55$ mg/g Nodakenin: $3,35 - 10,61$ mg/g Methoxsalen: < 3 mg/g		

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes

Changes to registation. Commission implementing Regulation (EO) 2017/2470, ANNEA is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Phenols: $13,0 - 40,0 \text{ mg/g}$ Coumarins: $13,0 - 40,0 \text{ mg/g}$ Iridoids: $13,0 - 39,0 \text{ mg/g}$ Saponins: $5,0 - 15,5 \text{ mg/g}$ Nutritive components Carbohydrates: $600 - 880 \text{ mg/g}$ Proteins: $70 - 170 \text{ mg/g}$ Fats: $< 4 \text{ mg/g}$ Microbiological parameters Total viable plate count: $< 5000 \text{ CFU/g}$ Total mold and yeast: $< 100 \text{ CFU/g}$ Coliform bacteria: $< 10 \text{ CFU/g}$ Salmonella : Negative/25 g <i>Escherichia coli</i> : Negative/25 g <i>Staphylococcus aureus</i> : Negative/25 g Heavy metals Lead: $< 0,65 \text{ mg/kg}$ Arsenic: $< 3,0 \text{ mg/kg}$ Mercury: $< 0,1 \text{ mg/kg}$ Cadmium: $< 1,0 \text{ mg/kg}$ CFU: Colony Forming Units]
Ferric Sodium EDTA	Description/Definition: Ferric Sodium EDTA (ethylenediaminetetraacetic acid) is an odourless free-flowing, yellow to brown powder with a chemical purity of more than 99 % (w/w). It is freely soluble in water. Chemical formula: C $_{10}$ H $_{12}$ FeN $_2$ NaO $_8$ * 3H $_2$ O Chemical characteristics: pH of 1 % solution: 3,5-5,5 Iron: 12,5-13,5 % Sodium: 5,5 % Water: 12,8 % Organic matter (CHNO): 68,4 % EDTA: 65,5-70,5 % Water insoluble matter: $\leq 0,1$ %
Ferrous ammonium phosphate	Description/Definition: Ferrous ammonium phosphate is a grey/green fine powder, practically insoluble in water and soluble in dilute mineral acids. CAS No.: 10101-60-7 Chemical formula: FeNH $_4$ PO $_4$ Chemical characteristics: pH of 5 % suspension in water: 6,8-7,8 Iron (total): $\ge 28 \%$ Iron (II): 22-30 % (w/w) Iron (III): $\le 7,0 \%$ (w/w) Ammonia: 5-9 % (w/w) Water: $\le 3,0 \%$
Fish peptides from <i>Sardinops</i> sagax	Description/Definition: The novel food ingredient is a peptide mixture, which is obtained by an alkaline protease-catalysed hydrolysis of fish (<i>Sardinops sagax</i>) muscle,

	Document Generated: 2024-06-27
known to be in for	Status: Point in time view as at 31/12/2020. ion: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes rece on or before 27 June 2024. There are changes that may be brought into force at a future date. en made appear in the content and are referenced with annotations. (See end of Document for details)
	subsequent isolation of the peptide fraction by column chromatography, concentration under vacuum and spray drying. Yellowish white powderPeptides (¹) (short chain peptides, dipeptides and tripeptides with a molecular weight of less than 2 kDa): \geq 85 g/100 g Val-Tyr (dipeptide): 0,1-0,16 g/100 g Ash: \leq 10 g/100 g Moisture: \leq 8 g/100 g
	(¹) Kjeldahl method
Flavonoids from <i>Glycyrrhiza</i> glabra	Description/Definition: Flavonoids derived from the roots or rootstock of <i>Glycyrrhiza glabra</i> L. are extracted with ethanol followed by further extraction of this ethanolic extract with medium-chain triglycerides. It is a dark-brown coloured liquid, containing 2,5 % to 3,5 % of glabridin. Moisture: < 0,5 % Ash: < 0,1 % Peroxide value (PV): < 0,5 meq/kg Glabridin: 2,5-3,5 % of fat Glycyrrhizinic acid: < 0,005 % Fat including polyphenol-type substances: \geq 99 % Protein: < 0,1 % Carbohydrates: not detectable
[^{F19} Fruit pulp, pulp juice, concentrated pulp juice from <i>Theobroma</i> <i>cacao</i> L. (Traditional food from a third country)	Description/Definition The traditional food is the fruit pulp from the cocoa (<i>Theobroma cacao</i> L) plant, which is the 'aqueous, mucilaginous and acidic substance in which the seeds are embedded'. Cocoa fruit pulp is obtained by splitting cocoa pods followed by separation from husks and beans; the pulp is then subject to pasteurisation and freezing. Cocoa pulp juice and/or cocoa concentrated pulp juice are produced following processing (enzymatic treatment, pasteurization, filtration, and concentration). Typical compositional data of cocoa fruit pulp, pulp juice, concentrated pulp juice Protein (g/100 g): 0,0 to 2,0 Total fat (g/100 g): 0,0 to 0,2 Total sugars (g/100 g): > 11,0
	Brix level (° Brix): ≥ 14 pH: 3,3 to 4,0 Microbiological criteria Total Plate Count (aerobic): < 10 000 cfu ⁱ /g Enterobacteriaceae: ≤ 10 cfu/g <i>Salmonella</i> : Absence in 25 g]
Fucoidan extract from the seaweed <i>Fucus</i> <i>vesiculosus</i>	Description/Definition: Fucoidan from the seaweed <i>Fucus vesiculosus</i> is extracted using aqueous extraction in acidic solution and filtration processes without the use of organic solvents. The resulting extract is concentrated and dried to yield the fucoidan extract with the following specifications: Off-white to brown powder Odour and Taste: Bland odour and taste Moisture: < 10 % (105 °C for 2 hours) pH value: 4,0-7,0 (1 % suspension at 25 °C)

Status: Point in time view as at 31/12/2020.

	Heavy metals: Arsenic (inorganic): < 1,0 ppm Cadmium: < 3,0 ppm Lead: < 2,0 ppm
Fucoidan extract from the seaweed Undaria pinnatifida	Description/Definition: Fucoidan from seaweed Undaria pinnatifida is extracted using aqueous extraction in acidic solution and filtration processes without the use of organic solvents. The resulting extract is concentrated and dried to yield the fucoidan extract with the following specifications: Off-white to brown powder Odour and Taste: Bland odour and tasteMoisture: < 10 % (105 °C for 2 hours) pH value: 4,0-7,0 (1 % suspension at 25 °C) Heavy metals: Arsenic (inorganic): < 1,0 ppm Cadmium: < 3,0 ppm Lead: < 2,0 ppm Microbiology: Total aerobic microbial count: < 10 000 CFU/g Yeast and mould count: < 100 CFU/g Total enterobacteria count: Absence/g <i>Escherichia coli</i> : Absence/g Salmonella : Absence/10 g Staphylococcus aureus : Absence/g Composition of the two permitted types of extracts, based on the level of fucoidan:

	Extract 1: Fucoidan: 75-95 % Alginate: 2,0-6,5 % Polyphloroglucinol: 0,5-3,0 % Mannitol: 1-10 % Natural salts/Free Minerals: 0,5-1,0 % Other carbohydrates: 0,5-2,0 % Protein: 2,0-2,5 % Extract 2: Fucoidan: 50-55 % Alginate: 2,0-4,0 % Polyphloroglucinol: 1,0-3,0 % Mannitol: 25-35 % Natural salts/Free Minerals: 8-10 % Other carbohydrates: 0,5-2,0 % Protein: 1,0-1,5 %
2'- Fucosyllactose (synthetic)	Definition: Chemical name: α-L-Fucopyranosyl- $(1\rightarrow 2)$ -β-D-galactopyranosyl- $(1\rightarrow 4)$ - D-glucopyranose Chemical formula: C ₁₈ H ₃₂ O ₁₅ CAS No: 41263-94-9 Molecular weight: 488,44 g/mol Description: 2'-fucosyllactose is a white to off-white powder that is produced by a chemical synthesis process. Purity: 2'-Fucosyllactose: $\geq 95 \%$ D-Lactose: $\leq 1,0 \text{ w/w }\%$ L-Fucose: $\leq 1,0 \text{ w/w }\%$ Difucosyl- D-lactose isomers: $\leq 1,0 \text{ w/w }\%$ 2'-Fucosyl- D-lactose isomers: $\leq 1,0 \text{ w/w }\%$ pH (20 °C, 5 % solution): 3,2-7,0 Water (%): $\leq 9,0 \%$ Ash, sulphated: $\leq 0,2 \%$ Acetic acid: $\leq 0,3 \%$ Residual solvents (methanol, 2-propanol, methyl acetate, acetone): $\leq 50,0$ mg/kg singly, $\leq 200,0 \text{ mg/kg}$ in combination Residual proteins: $\leq 0,01 \%$ Heavy Metals: Palladium: $\leq 0,1 \text{ mg/kg}$ Nickel: $\leq 3,0 \text{ mg/kg}$ Microbiological criteria: Aerobic mesophilic bacteria total count: $\leq 500 \text{ CFU/g}$ Yeasts and Moulds: $\leq 10 \text{ CFU/g}$ Residual endotoxins: $\leq 10 \text{ EU/mg}$
2'- Fucosyllactose (microbial source)	[^{F41} Definition: Chemical name: α-L-Fucopyranosyl-(1→2)-β-D-galactopyranosyl-(1→4)-D-glucopyranose Chemical formula: C ₁₈ H ₃₂ O ₁₅ CAS No: 41263-94-9 Molecular weight: 488,44 g/mol

Status: Point in time view as at 31/12/2020.

Source:	Source:
Genetically modified strain of	Genetically modified strain of
<i>Escherichia coli</i> K-12	<i>Escherichia coli</i> BL21
Description: 2'-Fucosyllactose is a white to off- white powder that is produced by a microbial process. Purity: 2'-Fucosyllactose: \geq 83 % D-Lactose: \leq 10,0 % L-Fucose: \leq 2,0 % Difucosyl-D-lactulose: \leq 1,5 % Sum of saccharides (2'- Fucosyl-D-lactulose): \geq 90 % pH (20 C, 5 % solution): 3,0-7,5 Water: \leq 9,0 % Sulphated ash: \leq 2,0 % Acetic acid: \leq 1,0 % Residual proteins: \leq 0,01 % Microbiological criteria: Aerobic mesophilic bacteria total count: \leq 3 000 CFU/g Yeasts: \leq 100 CFU/g Moulds: \leq 100 CFU/g Endotoxins: \leq 10 EU/mg	Description: 2'-Fucosyllactose is a white to off white powder and the liquid concentrate ($45 \% \pm 5 \%$ w/v) aqueous solution is a colourless to slight yellow clear aqueous solution. 2'-Fucosyllactose is produced by a microbiological process. Purity: 2'-Fucosyllactose: $\geq 90 \%$ Lactose: $\leq 5,0 \%$ Fucose: $\leq 3,0 \%$ 3-Fucosyllactose: $\leq 5,0 \%$ Fucosylgalactose: $\leq 5,0 \%$ Glucose: $\leq 3,0 \%$ Glucose: $\leq 3,0 \%$ Glucose: $\leq 3,0 \%$ Galactose: $\leq 3,0 \%$ Galactose: $\leq 3,0 \%$ Water: $\leq 9,0 \%$ (powder) Ash, sulphated: $\leq 0,5 \%$ (powder and liquid) Residual proteins: $\leq 0,01 \%$ (powder and liquid) Heavy Metals: Lead: $\leq 0,02 \text{ mg/kg}$ (powder and liquid) Arsenic: $\leq 0,2 \text{ mg/kg}$ (powder and liquid) Mercury: $\leq 0,5 \text{ mg/kg}$ (powder a liquid) Microbiological criteria: Total plate count: $\leq 10^4 \text{ CFU/g}$ (powder), $\leq 5 000 \text{ CFU/g}$ (liquid Yeasts and Moulds: $\leq 100 \text{ CFU/g}$ (powder), $\leq 5 000 \text{ CFU/g}$ (liquid Yeasts and Moulds: $\leq 100 \text{ CFU/g}$ (powder), $\leq 50 \text{ CFU/g}$ (liquid) Enterobacteriaceae/Coliforms: absence in 11 g (powder and liquid) <i>Salmonella</i> : negative/200 ml (liquid <i>Cronobacter</i> : negative/100 g (powder), negative/200 ml (liquid Endotoxins: $\leq 100 \text{ EU/g}$ (powder $\leq 100 \text{ EU/ml}$ (liquid) Aflatoxin M1: $\leq 0,025 \text{ µg/kg}$ (powder and liquid)]

[^{F20} 2'- Fucosyllactose/ Difucosyllactose mixture (' 2'- FL/DFL ') (microbial source)	Description/Definition: 2'-Fucosyllactose/Difucosyllactose mixture is a purified, white to off- white amorphous powder that is produced by a microbial process. After purification, the 2'-Fucosyllactose/Difucosyllactose mixture is isolated by spray drying. Source: Genetically modified strain of Escherichia coli strain K-12 DH1 Characteristics/Composition Appearance: White to off white powder or agglomerates Sum of 2'-Fucosyllactose, Difucosyllactose, Lactose and Fucose (% of dry matter): ≥ 92,0 % (w/w) Sum of 2'-fucosyllactose and difucosyllactose (% of dry matter): ≥ 85,0 % (w/w) 2'-Fucosyllactose (% of dry matter): ≥ 75,0 % (w/w) Difucosyllactose (% of dry matter): ≥ 5,0 % (w/w) Difucosyllactose (% of dry matter): ≥ 5,0 % (w/w) 2'-Fucosyl-D-lactulose: ≤ 2,0 % (w/w) Sum of other carbohydrates ^k : ≤ 6,0 % (w/w) Sum of other carbohydrates ^k : ≤ 6,0 % (w/w) Moisture: ≤ 6,0 % (w/w) PH (20 °C, 5 % solution): 4,0-6,0 Residual protein: ≤ 0,01 % (w/w) Microbiological criteria: Aerobic mesophilic bacteria total plate count: ≤ 1000 CFU/g <i>Enterobacteriaceae</i> : ≤ 10 CFU/g Salmonella sp.: Negative/25 g Yeast: ≤ 100 CFU/g Mould: ≤ 100 CFU/g Residual endotoxins: ≤ 10 EU/mg CFU: Colony Forming Units; EU: Endotoxin Units]
Galacto- oligosaccharide	Description/Definition: Galacto-oligosaccharide is produced from milk lactose by an enzymatic process using β-galactosidases from Aspergillus oryzae, Bifidobacterium bifidum, Pichia pastoris, Sporobolomyces singularis, Kluyveromyces lactis, Bacillus circulans, and Papiliotrema terrestris. GOS: min 46 % Dry Matter (DM) Lactose: max 40 % DM Glucose: max 27 % DM Galactose: min 0,8 % DM Ash: max 4,0 % DM Protein: max 4,5 % DM Nitrite: max. 2 mg/kg
Glucosamine HCl from Aspergillus niger and genetically modified strain of <i>E. coli</i> K-12	White crystalline odourless powder Molecular formula: C ₆ H ₁₃ NO ₅ · HCl Relative molecular mass: 215,63 g/mol D-Glucosamine HCl 98,0-102,0 % of reference standard (HPLC) Specific rotation + 70,0 ° - + 73,0 °

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes

known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

Glucosamine sulphate KCl from <i>Aspergillus</i> <i>niger</i> and genetically modified strain of <i>E. coli</i> K-12	White crystalline odourless powder Molecular formula: $(C_6 H_{14} NO_5)_2 SO_4 \cdot 2KCl$ Relative molecular mass: 605,52 g/mol D-Glucosamine Sulphate 2KCl 98,0-102,0 % of reference standard (HPLC) Specific Rotation +50,0 ° to +52,0 °
Glucosamine sulphate NaCl from <i>Aspergillus</i> <i>niger</i> and genetically modified strain of <i>E. coli</i> K-12	White crystalline odourless powder Molecular formula: $(C_6 H_{14} NO_5)_2 SO_4 \cdot 2NaCl$ Relative molecular mass: 573,31 g/mol D-Glucosamine HCl: 98-102 % of reference standard (HPLC) Specific Optical Rotation: +52 ° - +54 °
Guar Gum	Description/Definition: Native guar gum is the ground endosperm of seeds from natural strains of guar <i>Cyamopsis tetragonolobus</i> L. Taub. (<i>Leguminosae</i> family). It consists of a high molecular weight polysaccharide, primarily composed of galactopyranose and mannopyranose units combined through glycosidic linkages, which may be described chemically as a galactomannan (galactomannan content not less than 75 %). Appearance: White to yellowish powder Molecular weight: Between 50 000 – 8 000 000 Daltons CAS number: 9000-30-0 Einces Number: 232-536-8 Purity: As specified by Commission Regulation (EU) No 231/2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council [®] & by Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins ^b . Physico-chemical properties: Powder Shelf-life: 2 years Colour: White Odour: Light Average diameter of particles: 60-70µm Moisture: Max 15 % Viscosity * at 1 hour —Viscosity * at 2 hours: Min 3 600 mPa.s Viscosity * at 24 hours: Min 4 000 mPa.s Solubility: Soluble in hot and cold water pH for 10g/L, at 25 °C - 6-7,5 Flakes Useful life: 1 year Colour: White/off white with absence or minimal presence of black spots Odour: Light Average diameter of particles: 1-10 mm Moisture: Max 15 %

	Document Generated: 2024-06-27	
Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)		
	Viscosity * at 1 hour: Min 3 000 mPa.s Viscosity * at 2 hours — Viscosity * at 24 hours — Solubility — Soluble in hot and cold water pH for 10g/L, at 25 °C - 5-7,5	
	(*) The measurements of viscosity are carried out under the following conditions: 1 %, 25 °C, 20 rpm	
Heat-treated milk products fermented with <i>Bacteroides</i> <i>xylanisolvens</i>	Description/Definition: Heat-treated fermented milk products are produced with <i>Bacteroides xylanisolvens</i> (DSM 23964) as starter culture. Semi-skimmed milk (between 1,5 % and 1,8 % fat) or skimmed milk (0,5 % fat or less) is pasteurised or ultra-heat-treated before starting the fermentation with <i>Bacteroides xylanisolvens</i> (DSM 23964). The resulting fermented milk product is homogenised and then heat-treated to inactivate <i>Bacteroides xylanisolvens</i> (DSM 23964). The final product does not contain viable cells of <i>Bacteroides xylanisolvens</i> (DSM 23964). (¹).	
	(¹) Modified DIN EN ISO 21528-2.	
Hydroxytyrosol	Description/Definition:Hydroxytyrosol is a pale yellow viscous liquid obtained by chemical synthesisMolecular formula: $C_8 H_{10} O_3$ Molecular weight: 154,6 g/molCAS No: 10597-60-1Moisture $\leq 0,4 \%$ Odour: CharacteristicTaste: Slightly bitterSolubility (water): Miscible with waterpH: 3,5-4,5Refractive Index: 1,571-1,575Purity:Hydroxytyrosol: $\geq 99 \%$ Acetic acid: $\leq 0,4 \%$ Hydroxytyrosol acetate: $\leq 0,3 \%$ Sum of homovanillic acid, iso-homovanilic acid, and 3- methoxy-4hydroxyphenylglycol: $\leq 0,3 \%$ Heavy MetalsLead: $\leq 0,03 mg/kg$ Cadmium: $\leq 0,01 mg/kg$ Mercury: $\leq 0,01 mg/kg$ Residual SolventsEthyl acetate: $\leq 2,50 mg/kg$ Isopropanol: $\leq 2,50 mg/kg$ Tetrahydrofuran: $\leq 0,01 mg/kg$	
Ice Structuring Protein type III HPLC 12	Description/Definition: The Ice Structuring Protein (ISP) preparation is a light-brown liquid produced by submerged fermentation of a genetically-modified strain of food-grade baker's yeast (<i>Saccharomyces cerevisiae</i>) in which a synthetic gene for the ISP has been inserted into the yeast's genome.	

	The protein is expressed and secreted into the growth medium where it is separated from the yeast cells by micro-filtration and concentrated by ultra-filtration. As a result, the yeast cells are not transferred into the ISP preparation as such or under an altered form. The ISP preparation consists of native ISP, glycosylated ISP and proteins and peptides from the yeast and sugars as well as acids and salts commonly found in food. The concentrate is stabilised with 10 mM citric acid buffer. Assay: ≥ 5 g/l active ISP pH: 2,5-3,5 Ash: $\leq 2,0$ % DNA: Not detectable
Aqueous extract of dried leaves of <i>Ilex</i> guayusa	Description/Definition: Dark brown liquid. Aqueous extracts of dried leaves of <i>Ilex guayusa</i> . Composition: Protein: < 0,1 g/100 ml Fat: < 0,1 g/100 ml Carbohydrate: 0,2–0,3 g/100 ml Total sugars: < 0,2 g/100 ml Caffeine: 19,8–57,7 mg/100 ml Theobromine: 0,14–2,0 mg/100 ml Chlorogenic acids: 9,9–72,4 mg/100ml
[^{F21} Infusion from coffee leaves of <i>Coffea</i> <i>arabica</i> L. and/or <i>Coffea</i> <i>canephora</i> Pierre ex A. Froehner (Traditional food from a third country)	Description/Definition:The traditional food consists of an infusion of leaves from Coffea arabica L. and/or Coffea canephora Pierre ex A.Froehner (family: Rubiaceae).The traditional food is prepared by mixing a maximum of 20 g of dried leaves from Coffea arabica L. and/or Coffea canephora Pierre ex A.Froehner with 1 L of hot water. Leaves are removed and the infusion is then subjected to pasteurization (at least 71 °C for 15 seconds).Composition: Visual: Brown green liquid Odour and taste: Characteristic Chlorogenic acid (5-CQA): < 100 mg/L Caffeine: < 80 mg/L Epigallocatechin gallate (EGCG): < 700 mg/L Microbiological criteria: Total plate count: < 500 CFU/g Total veast and mould count: < 100 CFU/g Total coliforms: < 100 CFU/g Escherichia coli : Absence in 1 g Salmonella : Absence in 25 g Heavy metals: Lead (Pb): < 3,0 mg/L Arsenic (As): < 2,0 mg/L Cadmium (Cd): < 1,0 mg/L CFU: Colony Forming Units]
Isomalto- oligosaccharide	Powder:Solubility (water) (%): > 99Glucose (% dry basis): $\leq 5,0$ Isomaltose + DP3 to DP9 (% dry basis): ≥ 90 Moisture (%): $\leq 4,0$

Lactitol	Description/Definition:
	(¹) Food and Nutrition Paper 5 Rev. 2 — Guide to specifications for general notices, general analytical techniques, identification tests, test solutions and other reference materials (JECFA), 1991, 322 pp., English, ISBN 92-5-102991-1.
	Determine using an atomic absorption technique appropriate to the specified level. The selection of sample size and method of sample preparation may be based on the principles of the method described in FNP $5(^{1})$, 'Instrumental methods'
	Lead: $\leq 0.1 \text{ mg/kg}$
	Heavy metals:
	Assay: \geq 98 % on the dry basis Loss on drying: \leq 6,5 % (60 °C, 5 hours)
	Purity:
	Formula weight: 360,3 (monohydrate)
	OH HO OH
	HOM OH
	O OH · H ₂ O
	0,0
	ОН
	Structural formula
	Chemical formula: $C_{12}H_{22}O_{11} \cdot H_2O$
	fructofuranose, monohydrate CAS No.: 13718-94-0
	crystals with a sweet tasteChemical name: 6-O-α-D-glucopyranosyl-D-
	sucrose by an enzymatic process. The commercial product is the monohydrate. Appearance: Virtually odourless, white or almost white
	A reducing disaccharide that consists of one glucose and one fructose moiety linked by an alpha-1,6-glycosidic bond. It is obtained from
Isomaltulose	Description/Definition: A reducing disappheride that consists of one glucose and one fructose
	Arsenic (mg/kg) : $\leq 0,5$
	Heavy metals: Lead (mg/kg): ≤ 0.5
	pH: 4 - 6 Sulphated $ash(g/100 g)$: $\leq 0,3$
	Isomaltose + DP3 to DP9 (% dry basis): \geq 90
	Dried solids (g/100 g): > 75 Glucose (% dry basis): $\le 5,0$
	Arsenic (mg/kg): ≤ 0.5 Syrup:
	Heavy metals: Lead (mg/kg): ≤ 0.5
	neavy metais:

Status: Point in time view as at 31/12/2020.

	Crystalline powder or colourless solution manufactured via catalytic hydrogenation of lactose. Crystalline products occur in anhydrous, monohydrate and dihydrate forms. Nickel is used as a catalyst. Chemical name: 4-O- β -D-Galactopyranosyl-D-glucitol Chemical formula: C ₁₂ H ₂₄ O ₁₁ Molecular weight: 344,31 g/mol CAS No: 585-86-4 Purity: Solubility (in water): Very soluble in water Specific rotation [α] _D ²⁰ = + 13 ° to + 16 ° Assay: \geq 95 % d.b (d.b — expressed on the dry weight basis) Water: \leq 10,5 % Other polyols: \leq 2,5 % d.b Reducing sugars: \leq 0,2 % d.b Chlorides: \leq 100 mg/kg d.b Sulphated ash: \leq 0,1 % d.b Nickel: \leq 2,0 mg/kg d.b Arsenic: \leq 3,0 mg/kg d.b Lead: \leq 1,0 mg/kg d.b
Lacto- N - neotetraose (synthetic)	Definition: Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 4)$ -2-acetamido-2- deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ - D- glucopyranose Chemical formula: C $_{26}$ H $_{45}$ NO $_{21}$ CAS No: 13007-32-4 Molecular weight: 707,63 g/mol Description: Lacto- <i>N</i> -neotetraose is a white to off-white powder. Produced by a chemical synthesis process and is isolated by crystallisation. Purity: Assay (water free): $\geq 96 \%$ D-Lactose: $\leq 1,0 \%$ Lacto-N-neotetraose fructose isomer: $\leq 0,6 \%$ pH (20 °C, 5 % solution): 5,0-7,0 Water: $\leq 9,0 \%$ Ash, sulphated: $\leq 0,4 \%$ Acetic acid: $\leq 0,3 \%$ Residual solvents (methanol, 2-propanol, methyl acetate, acetone): ≤ 50 mg/kg singly, ≤ 200 mg/kg in combination Residual proteins: $\leq 0,01 \%$ Palladium: $\leq 0,1$ mg/kg Nickel: $\leq 3,0$ mg/kg Microbiological criteria: Aerobic mesophilic bacteria total count: ≤ 500 CFU/g Yeasts: ≤ 10 CFU/g Moulds: ≤ 10 CFU/g Residual endotoxins: ≤ 10 EU/mg
[^{F42} Lacto N neotetraose (microbial source)	Definition: Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 4)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Chemical formula: C ₂₆ H ₄₅ NO ₂₁

	CACN 12007 20 4
	CAS No: 13007-32-4
	Molecular weight: 707,63 g/mol
	Source:
	Genetically modified strain of <i>Escherichia coli</i> K-12
	Description: Lacto- <i>N</i> -neotetraose is a white to off-white powder that is produced by a
	microbiological process.
	Purity:
	Assay (water free): $\geq 80 \%$
	D-Lactose: $\leq 10,0\%$
	Lacto- N -triose II: $\leq 3,0\%$
	para -Lacto- N -neohexaose: $\leq 5,0\%$
	Lacto- N -neotetraose fructose isomer: $\leq 1,0\%$
	Sum of saccharides (Lacto- N -neotetraose, D-Lactose, Lacto- N -triose
	II, <i>para</i> -Lacto- <i>N</i> -neohexaose, Lacto- <i>N</i> -neotetraose fructose isomer):
	$\geq 92\%$
	pH (20 C, 5 % solution): 4,0-7,0
	Water: $\leq 9,0\%$
	Ash, sulphated: ≤ 0.4 %
	Residual solvents (methanol): $\leq 100 \text{ mg/kg}$
	Residual proteins: $\leq 0.01 \%$
	Microbiological criteria:
	Aerobic mesophilic bacteria total count: \leq 500 CFU/g
	Yeasts: ≤ 10 CFU/g
	Moulds: $\leq 10 \text{ CFU/g}$
	Residual endotoxins: $\leq 10 \text{ EU/mg}$
	CFU: Colony Forming Units; EU: Endotoxin Units.]
[F23] a sta N/	
$[^{F23}Lacto-N-$	Definition:
tetraose ('LNT')	Definition: Chemical formula: C ₂₆ H ₄₅ O ₂₁
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26}H_{45}O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -
tetraose ('LNT')	Definition: Chemical formula: $C_{26}H_{45}O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26}H_{45}O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description:
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26}H_{45}O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process.
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1 \rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1 \rightarrow 3)$ - β -D-galactopyranosyl- $(1 \rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition:
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1 \rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1 \rightarrow 3)$ - β -D-galactopyranosyl- $(1 \rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 \%$ (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1 \rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1 \rightarrow 3)$ - β -D-galactopyranosyl- $(1 \rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 \%$ (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): $\geq 70,0 \%$ (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1 \rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1 \rightarrow 3)$ - β -D-galactopyranosyl- $(1 \rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): \geq 90,0 % (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): \geq 70,0 % (w/w) D-Lactose: \leq 12,0 % (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: C $_{26}$ H $_{45}$ O $_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): \geq 90,0 % (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): \geq 70,0 % (w/w) D-Lactose: \leq 12,0 % (w/w) Lacto- <i>N</i> -tetraose II: \leq 10,0 % (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 \%$ (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): $\geq 70,0 \%$ (w/w) D-Lactose: $\leq 12,0 \%$ (w/w) Lacto- <i>N</i> -tetraose II: $\leq 10,0 \%$ (w/w) <i>Para</i> -lacto- <i>N</i> -hexaose-2: $\leq 3,5 \%$ (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 \%$ (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): $\geq 70,0 \%$ (w/w) D-Lactose: $\leq 12,0 \%$ (w/w) Lacto- <i>N</i> -tetraose II: $\leq 10,0 \%$ (w/w) Para -lacto- <i>N</i> -hexaose-2: $\leq 3,5 \%$ (w/w) Lacto- <i>N</i> -tetraose fructose isomer: $\leq 1,0 \%$ (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 \%$ (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): $\geq 70,0 \%$ (w/w) D-Lactose: $\leq 12,0 \%$ (w/w) Lacto- <i>N</i> -tetraose II: $\leq 10,0 \%$ (w/w) Para -lacto- <i>N</i> -hexaose-2: $\leq 3,5 \%$ (w/w) Lacto- <i>N</i> -tetraose fructose isomer: $\leq 1,0 \%$ (w/w) Sum of other carbohydrates: $\leq 5,0 \%$ (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 \%$ (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): $\geq 70,0 \%$ (w/w) D-Lactose: $\leq 12,0 \%$ (w/w) Lacto- <i>N</i> -tetraose II: $\leq 10,0 \%$ (w/w) Lacto- <i>N</i> -tetraose fuctose isomer: $\leq 1,0 \%$ (w/w) Sum of other carbohydrates: $\leq 5,0 \%$ (w/w) Moisture: $\leq 6,0 \%$ (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β -D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 %$ (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): $\geq 70,0 %$ (w/w) D-Lactose: $\leq 12,0 %$ (w/w) Lacto- <i>N</i> -tetraose II: $\leq 10,0 %$ (w/w) Lacto- <i>N</i> -tetraose fuctose isomer: $\leq 1,0 %$ (w/w) Sum of other carbohydrates: $\leq 5,0 %$ (w/w) Moisture: $\leq 6,0 %$ (w/w)
tetraose ('LNT') (microbial	Definition: Chemical formula: $C_{26} H_{45} O_{21}$ Chemical name: β -D-Galactopyranosyl- $(1\rightarrow 3)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1\rightarrow 3)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Molecular mass: 707,63 Da CAS No 14116-68-8 Description: Lacto- <i>N</i> -tetraose is a purified, white to off-white amorphous powder that is produced by a microbial process. Source: Genetically modified strain of <i>Escherichia coli</i> strain K-12 DH1 Characteristics/Composition: Appearance: White to off-white powder Sum of lacto- <i>N</i> -tetraose, D-Lactose and lacto- <i>N</i> -tetraose II (% of dry matter): $\geq 90,0 \%$ (w/w) Lacto- <i>N</i> -tetraose (% of dry matter): $\geq 70,0 \%$ (w/w) D-Lactose: $\leq 12,0 \%$ (w/w) Lacto- <i>N</i> -tetraose II: $\leq 10,0 \%$ (w/w) Lacto- <i>N</i> -tetraose fuctose isomer: $\leq 1,0 \%$ (w/w) Sum of other carbohydrates: $\leq 5,0 \%$ (w/w) Moisture: $\leq 6,0 \%$ (w/w)

Status: Point in time view as at 31/12/2020. **Changes to legislation:** Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes

known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Microbiological criteria:Aerobic mesophilic bacteria total plate count: $\leq 1\ 000\ CFU/g$ Enterobacteriaceae : $\leq 10\ CFU/g$ Salmonella sp.: Negative/25 gYeast: $\leq 100\ CFU/g$ Mould: $\leq 100\ CFU/g$ Residual endotoxins: $\leq 10\ EU/mg$ CFU: Colony Forming Units; EU: Endotoxin Units.]
[^{F24} Lonicera caerulea L. berries (haskap) (Traditional food from a third country)	Description/Definition: The traditional food are fresh and frozen berries from <i>Lonicera caerulea</i> var. edulis. <i>Lonicera caerulea</i> L. is a deciduous shrub belonging to the <i>Caprifoliaceae</i> family. Typical nutritional components of haskap berries (given in fresh berries): Carbohydrates: 12,8 % Fibre: 2,1 % Lipids: 0,6 % Proteins: 0,7 % Ash: 0,4 % Water: 85,5 %]
Lucerne leaf extract from <i>Medicago sativa</i>	Description/Definition: The Lucerne (<i>Medicago sativa</i> L.) is processed within 2 hours after harvest. It is chopped and crushed. By passing through an oleaginous- type press, the Lucerne provides a fibrous residue and press juice (10 % of dry matter). The dry matter of this juice contains about 35 % of crude protein. The press juice (pH 5,8-6,2) is neutralised. Preheating and vapour injection allows coagulation of proteins associated with carotenoid and chlorophyll pigments. The protein precipitate is separated by centrifugation and thereafter dried. After adding ascorbic acid the Lucerne protein concentrate is granulated and stored in inert gas or in cold storage. Composition: Protein: 45-60 % Fat: 9-11 % Free carbohydrates (soluble fibre): 1-2 % Polysaccharides (insoluble fibre): 11-15 % including cellulose: 2-3 % Minerals: 8-13 % Saponins: ≤ 1,4 % Isoflavones: ≤ 350 mg/kg Coumestrol: ≤ 100 mg/kg Phytates: ≤ 200 mg/kg L-canavanine: ≤ 4,5 mg/kg
Lycopene	Description/Definition: Synthetic lycopene is produced by the Wittig condensation of synthetic intermediates commonly used in the production of other carotenoids used in food. Synthetic lycopene consists of \geq 96 % lycopene and minor quantities of other related carotenoid components. Lycopene is presented either as a powder in a suitable matrix or an oily dispersion. The colour is dark red or red-violet. Antioxidative protection has to be assured.

	Chemical name: Lycopene CAS No.: 502-65-8 (<i>all</i> -trans lycopene) Chemical formula: C ₄₀ H ₅₆ Formula weight: 536,85 Da
Lycopene from <i>Blakeslea</i> trispora	Description/Definition: The purified lycopene from <i>Blakeslea trispora</i> consists of \geq 95 % lycopene and \leq 5 % other carotenoids. It is presented either as a powder in a suitable matrix or an oily dispersion. The colour is dark red or redviolet. Anti-oxidative protection has to be assured. Chemical name: Lycopene CAS No.: 502-65-8 (all trans lycopene) Chemical formula: C ₄₀ H ₅₆ Formula weight: 536,85 Da
Lycopene from tomatoes	Description/Definition: The purified lycopene from tomatoes (<i>Lycopersicon esculantum</i> L.) consists of \geq 95 % lycopene and \leq 5 % other carotenoids. It is presented either as a powder in a suitable matrix or an oily dispersion. The colour is dark red or red-violet. Anti-oxidative protection has to be assured. Chemical name: Lycopene CAS No.: 502-65-8 (all trans lycopene) Chemical formula: C ₄₀ H ₅₆ Formula weight: 536,85 Da
Lycopene oleoresin from tomatoes	Description/Definition: Lycopene oleoresin from tomatoes is obtained by solvent extraction of ripe tomatoes (<i>Lycopersicon esculentum Mill.</i>) with subsequent removal of the solvent. It is a red to dark brown viscous, clear liquid. Total lycopene: 5-15 % Thereof trans-lycopene: 90-95 % Total carotenoids (calculated as lycopene): 6,5-16,5 % Other carotenoids: 1,75 % (Phytoene/phytofluene/ β -carotene): (0,5-0,75/0,4-0,65/0,2-0,35 %) Total tocopherols: 1,5-3,0 % Unsaponifiable matter: 13-20 % Total fatty acids: 60-75 % Water (Karl Fischer): \leq 0,5 %
[^{F16} Hen egg white lysozyme hydrolysate	Description/Definition Hen egg white lysozyme hydrolysate is obtained from hen egg white lysozyme by an enzymatic process, using subtilisin from <i>Bacillus</i> <i>licheniformis</i> . The product is a white to light yellow powder. Specification Protein (TN(*) x 5,30): 80-90 % Tryptophan: 5-7 % Ratio Tryptophan/LNAA(**): 0,18-0.25 Degree of hydrolysis: 19-25 % Moisture: < 5 % Ash: < 10 % Sodium: < 6 % Heavy metals Arsenic: < 1 ppm

Status: Point in time view as at 3	31/12/2020.
------------------------------------	-------------

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes	
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.	
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)	
	-

	Lead: < 1 ppm Cadmium: < 0,5 ppm Mercury: < 0,1 ppm Microbiological criteria Total aerobic count: < 10^{3} CFU/g Total combined yeasts/moulds count: < 10^{2} CFU/g Enterobacteria: < 10 CFU/g Salmonella spp: Absence in 25 g Escherichia coli : Absence in 10 g Staphylococcus aureus : Absence in 10 g * TN: total nitrogen ** LNAA: large neutral amino acids]
Magnesium citrate malate	Description/Definition: Magnesium citrate malate is a white to yellowish-white, amorphous powder.Chemical formula: Mg 5 (C 6 H 5 O 7) 2 (C 4 H 4 O 5) 2 Chemical name: Pentamagnesium di-(2-hydroxybutanedioate)-di-(2- hydroxypropane-1,2,3-tricarboxylate) CAS No.: 1259381-40-2 Molecular weight: 763,99 Daltons (anhydrous) Solubility: Freely soluble in water (about 20 g in 100 ml) Description of the physical state: Amorphous powder Assay magnesium: 12,0-15,0 % Loss on drying (120 °C/4 hours): ≤ 15 % Colour (solid): White to yellowish-white Colour (solid): White to yellowish-white Colour (20 % aqueous solution): Colourless to yellowish Appearance (20 % aqueous solution): Clear solution pH (20 % aqueous solution): Approx. 6,0 Impurities: Chloride: ≤ 0,05 % Sulphate: ≤ 0,05 % Sulphate: ≤ 0,05 % Arsenic: ≤ 3,0 ppm Lead: ≤ 2,0 ppm Cadmium: ≤ 1 ppm Mercury: ≤ 0,1 ppm
Magnolia Bark Extract	Description/Definition: Magnolia bark extract is obtained from the bark of the plant <i>Magnolia</i> <i>officinalis</i> L. and produced with supercritical carbon dioxide. The bark is washed and oven dried to reduce moisture content before being crushed and extracted with supercritical carbon dioxide. The extract is dissolved in medical-grade ethanol and re-crystallised to yield magnolia bark extract. Magnolia bark extract is mainly composed of two phenolic compounds, magnolol and honokiol. Appearance: Light brownish powder Purity: Magnolol: $\geq 85,2 \%$ Honokiol: $\geq 0,5 \%$ Magnolol & Honokiol: $\geq 94 \%$ Total Eudesmol: $\leq 2 \%$

Maize-germ oil high in unsaponifiable matter	Moisture: 0,50 % Heavy metals: Arsenic (ppm): $\leq 0,5$ Lead (ppm): $\leq 0,5$ Methyl eugenol (ppm): ≤ 10 Tubocurarine (ppm): $\leq 2,0$ Total Alkaloid (ppm): ≤ 100 Description/Definition: Maize-germ oil high in unsaponifiable matter is produced by vacuum distillation and it is different from refined maize-germ oil in the concentration of the unsaponifiable fraction (1,2 g in refined maize-germ oil and 10 g in 'maize-germ oil high in unsaponifiable matter'). Purity:
	Unsaponifiable matter: > 9,0 g/100 g Tocopherols: \geq 1,3 g/100 g
	α-tocopherol (%): 10-25 % β-tocopherol (%): < 3,0 %
	γ-tocopherol (%): 68-89 %
	δ-tocopherol (%): < 7,0 % Starola, tritornania alaphala, mathylatarola: > 6.5 g/100 g
	Sterols, triterpenic alcohols, methylsterols: > 6,5 g/100 g Fatty acids in triglycerides:
	palmitic acid: 10,0-20,0 % stearic acid: < 3,3 %
	oleic acid: 20,0-42,2 %
	linoleic acid: 34,0-65,6 %
	linolenic acid: $< 2,0 \%$ Acid value: $\le 6,0 \text{ mg KOH/g}$
	Peroxide value (PV): $\leq 10 \text{ mEq O}_2/\text{kg}$
	Heavy metals: Iron (Fe): < 1 500 μg/kg
	Copper (Cu): $< 100 \ \mu g/kg$
	Impurities: Polycyclic aromatic hydrocarbons (PAH) Benzo(a)pyrene: < 2 µg/kg Treatment with active carbon is required to ensure that polycyclic aromatic hydrocarbons (PAH) are not enriched in the production of 'maize-germ oil high in unsaponifiable matter'
Methylcellulose	Description/Definition: Methyl cellulose is cellulose obtained directly from natural strains of fibrous plant material and partially etherified with methyl groups. Chemical name: Methyl ether of cellulose Chemical formula: The polymers contain substituted anhydroglucose
	units with the following general formula: C6H7O2(OR1)(OR2)(OR3) where R1, R2, R3 each may be one of the following:
	$\begin{array}{c} - & CH_3 \text{ or} \\ - & CH_2 CH_3 \end{array}$
	Molecular weight: Macromolecules: from about 20 000 (n about 100) up to about 380 000 g/mol (n about 2 000) Assay: Content not less than 25 % and not more than 33 % of methoxyl groups (-OCH ₃) and not more than 5 % of hydroxyethoxyl groups (-
	OCH ₂ CH ₂ OH)

Status: Point in time view as at 31/12/2020.	
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes	
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.	
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)	

	Slightly hygroscopic white or slightly yellowish or greyish odourless and tasteless, granular or fibrous powder. Solubility: Swelling in water, producing a clear to opalescent, viscous, colloidal solution. Insoluble in ethanol, ether and chloroform. Soluble in glacial acetic acid. Purity: Loss on drying: $\leq 10 \% (105 \text{ °C}, 3 \text{ hours})$ Sulphated Ash: $\leq 1,5 \%$ determined at $800 \pm 25 \text{ °C}$ pH: $\geq 5,0$ and $\leq 8,0 (1 \%$ colloidal solution) Heavy metals: Arsenic: $\leq 3,0 \text{ mg/kg}$ Lead: $\leq 2,0 \text{ mg/kg}$ Mercury: $\leq 1,0 \text{ mg/kg}$
[^{F25} 1- Methylnicotinam	Definition:
chloride	Chemical formula: C ₇ H ₉ N ₂ OCl
	CAS No: 1005-24-9 Molecular weight: 172,61 Da
	Description
	1-Methylnicotinamide chloride is white or off-white, crystalline solid produced by a chemical synthesis process.
	Characteristics/Composition
	Appearance: White – off-white, crystalline solid
	Purity: \geq 98,5 % Trigonelline: \leq 0,05 %
	Nicotinic Acid: ≤ 0.10 %
	Nicotinamide: $\leq 0,10$ %
	Largest unknown impurity: $\leq 0.05 \%$ Sum of unknown impurities: $\leq 0.20 \%$
	Sum of all impurities: $\leq 0,20\%$
	Solubility: soluble in water and methanol. Practically insoluble in 2-
	propanol and dichloromethane Moisture: $\leq 0.3 \%$
	Loss on drying: $\leq 1,0$ %
	Residue on ignition: $\leq 0,1$ %
	Residual Solvents and Heavy Metals Methanol: $\leq 0.3 \%$
	Heavy metals: $\leq 0.002 \%$
	Microbiological criteria:
	Total aerobic microbial count: $\leq 100 \text{ CFU/g}$ Mould/yeast: $\leq 10 \text{ CFU/g}$
	Enterobacteriaceae: absence in 1 g
	Pseudomonas aeruginosa : absence in 1 g
	Staphylococcus aureus : absent in 1 g CFU: Colony Forming Units]
(68)-5-	Description/Definition:
• •	• Colligencial name: N-[4-[[[(6S)-2-amino-1,4,5,6,7,8-hexahydro-5-methyl-4-
acid, glucosamine	oxo-6-pteridinyl]methyl]amino]benzoyl]-L-glutamic acid, glucosamine salt
salt	Chemical formula: C ₃₂ H ₅₁ N ₉ O ₁₆
	Molecular weight: 817,80 g/mol (anhydrous)

<i>Status: Point in time view as at 31/12/2020.</i>
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	CAS No.: 1181972-37-1 Appearance: Creamy to light-brown powder Purity: Diastereoisomeric purity: At least 99 % of (6S)-5-methyltetrahydrofolic acid Glucosamine assay: 34-46 % in dry basis 5-Methyltetrahydrofolic acid assay: 54-59 % in dry basis Water: $\leq 8,0$ % Heavy metals: Lead: $\leq 2,0$ ppm Cadmium: $\leq 1,0$ ppm Mercury: $\leq 0,1$ ppm Arsenic: $\leq 2,0$ ppm Boron: ≤ 10 ppm Microbiological criteria: Total aerobic microbial count: ≤ 100 CFU/g Yeasts and moulds: ≤ 100 CFU/g <i>Escherichia coli</i> : Absence in 10g
Monomethylsilar	deriver in the second s
(Organic Silicon)	Chemical name: Silanetriol, 1-methyl- Chemical formula: CH $_6$ O $_3$ Si Molecular weight: 94,14 g/mol CAS No: 2445-53-6 Purity: Organic Silicon (monomethylsilanetriol) preparation (aqueous solution): Acidity (pH): 6,4-6,8 Silicon: 100-150 mg Si/l Heavy metals: Lead: $\leq 1,0 \ \mu g/l$ Mercury: $\leq 1,0 \ \mu g/l$ Cadmium: $\leq 1,0 \ \mu g/l$ Arsenic: $\leq 3,0 \ \mu g/l$ Solvents: Methanol: $\leq 5,0 \ mg/kg$ (residual presence)
Mycelial extract from Shiitake mushroom (<i>Lentinula</i> edodes)	Description/Definition:The novel food ingredient is a sterile aqueous extract obtained from themycelium of Lentinula edodes cultivated in a submerged fermentation.It is a light brown, slightly turbid liquid.Lentinan is a β -(1-3) β -(1-6)-D-glucan which has a molecular weight ofapproximately 5×10^5 Daltons, a degree of branching of 2/5 and a triplehelical tertiary structure.Purity/Composition of the mycelial extract from Lentinula edodes :Moisture: 98 %Dry matter: 2 %Free glucose: < 20 mg/mlTotal protein(1): < 0,1 mg/mlN-containing constituents(2): < 10 mg/mlLentinan: $0,8 - 1,2$ mg/ml\(1)Bradford method(2)Kjeldahl method

[^{F26} Nicotinamide riboside chloride	Description/Definition: The novel food is a synthetic form of nicotinamide riboside. The novel food contains ≥ 90 % nicotinamide riboside chloride, predominantly in its β form, the remaining components being residual solvents, reaction by-products and degradation products. Nicotinamide riboside chloride: CAS number: 23111-00-4 EC number: 807-820-5 IUPAC name: 1-[(2R,3R,4S,5R)-3,4-dihydroxy-5- (hydroxymethyl)oxolan-2-yl]pyridin-1-ium-3-carboxamide;chloride Chemical formula: C ₁₁ H ₁₅ N ₂ O ₅ Cl Molecular weight: 290,7 g/mol Characteristics/Composition: Colour: White to light brown Form: Powder Identification: Conforms by NMR (nuclear magnetic resonance) Nicotinamide riboside chloride: ≥ 90 % Water content: ≤ 2 % Residual solvents: Acetone: ≤ 5 000 mg/kg Methanol: ≤ 1 000 mg/kg Methanol: ≤ 1 000 mg/kg Acetonitrile: ≤ 50 mg/kg Methyl tert-butyl ether: ≤ 500 mg/kg Acetonici ≤ 27 mg/kg Acetic acid: ≤ 5 000 mg/kg Acetate: ≤ 1 000 mg/kg Acetic acid: ≤ 5 000 mg/kg Heavy metals: Arsenic: ≤ 1 mg/kg Microbiological criteria: Total Plate Count: ≤ 1 000 CFU/g <i>Escherichia coli</i> : Absence in 10 g]
Noni fruit juice (<i>Morinda</i> <i>citrifolia</i>)	Description/Definition: Noni fruits (fruits of <i>Morinda citrifolia</i> L.) are pressed. The obtained juice is pasteurised. An optional fermentation step before or after the pressing may occur. Rubiadin: $\leq 10 \ \mu g/kg$ Lucidin: $\leq 10 \ \mu g/kg$
Noni fruit juice powder (<i>Morinda</i> <i>citrifolia</i>)	Description/Definition: Seeds and skin of the sun-dried fruits of <i>Morinda citrifolia</i> are separated. The obtained pulp is filtered to separate juice from the flesh. Desiccation of the produced juice occurs in one or two ways: Either by atomisation using maize maltodextrins, this mixture is obtained by keeping the rates of inflow of the juice and maltodextrins constant Or by zeodratation or drying and then mixing with an excipient, this process allows the juice to be dried initially and then mixed with maltodextrins (same amount as used in atomisation).
Noni fruit puree and concentrate	Description/Definition: The fruits of <i>Morinda citrifolia</i> are harvested by hand. Seeds and skin may be separated mechanically from the pureed fruits. After

(Morinda citrifolia)	pasteurisation, the puree is packaged in aseptic containers and stored under cold conditions.no_br Morinda citrifolia concentrate is prepared from M. citrifolia puree by treatment with pectinolytic enzymes (50– 60 °C for 1-2 h). Then the puree is heated to inactivate the pectinases and then immediately cooled. The juice is separated in a decanter centrifuge. Afterwards the juice is collected and pasteurised, prior to being concentrated in a vacuum evaporator from a brix of 6 to 8 to a brix of 49 to 51 in the final concentrate. Composition: Puree: Moisture: 89-93 % Protein: < 0.6 g/100 g Fat: ≤ 0.4 g/100 g Total carbohydrates: 5-10 g/100 g Fructose: 0,5-3,82 g/100 g Glucose: 0,5-3,14 g/100 g Dietary fibre: < 0,5-3 g/100 g 5,15-dimethylmorindol (1): $\leq 0.254 \mu g/ml$ Lucidin (1): Not detectable Alizarin (1): Not detectable Rubiadin (1): Not detectable Rubiadin (1): Not detectable Concentrate: Moisture: 48-53 % Protein: 3-3,5 g/100 g Fat: < 0,04 g/100 g Fat: < 0,04 g/100 g Fat: < 0,04 g/100 g Glucose: 9-11 g/100 g Dietary fibre: 1,5-5,0 g/100 g Fructose: 9-11 g/100 g Dietary fibre: 1,5-5,0 g/100 g S,15-dimethylmorindol (¹): $\leq 0.254 \mu g/ml$ (¹) By an HPLC-UV method developed and validated for the analysis of anthraquinones in Morinda citrifolia puree and concentrate. Limits of detection: 2,5 ng/ml (5,15 dimethylmorindol); 50,0 ng/ml (lucidin); 6,3 ng/ml (alizarin) and 62,5 ng/ml (rubiadin).
Noni leaves (<i>Morinda</i> <i>citrifolia</i>)	Description/Definition: After cutting, the leaves of <i>Morinda citrifolia</i> are subject to drying and roasting steps. The product has a particle size ranging from broken leaves to coarse powder with fines. It is of greenish brown to brown colour. Purity/Composition: Moisture: $< 5,2 \%$ Protein: 17- 20 % Carbohydrate: 55-65 % Ash: 10-13 % Fat: 4-9 % Oxalic acid: $< 0,14 \%$ Tannic acid: $< 2,7 \%$ 5,15-dimethylmorindol: $< 47 \text{ mg/kg}$ Rubiadin: non detectable, $\le 10 \mu\text{g/kg}$

Status: Point in time view as at 31/12/2020. Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.

Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

Noni fruit powder (<i>Morinda</i> <i>citrifolia</i>)	Lucidin: non detectable, $\leq 10 \ \mu g/kg$ Description/Definition: Noni fruit powder is made from pulped noni (<i>Morinda citrifolia L.</i>) fruits by freeze-drying. Fruits are pulped and seeds are removed. After freeze-drying during which water is removed from noni fruits, the remaining noni pulp is milled to a powder and encapsulated. Purity/Composition Moisture: 5,3-9 % Protein: 3,8-4,8 g/100 g Fat: 1-2 g/100 g Ash: 4,6-5,7 g/100 g Total carbohydrates: 80-85 g/100 g Fructose: 20,4-22,5 g/100 g Glucose: 22-25 g/100 g Dietary fibre: 15,4-24,5 g/100 g 5,15-dimethylmorindol (¹): $\leq 2,0 \ \mu g/ml$ (¹) By an HPLC-UV method developed and validated for the analysis of anthraquinones in Morinda citrifolia fruit powder. Limits of detection: 2,5 ng/ml (5,15 dimethylmorindol)	
<i>Odontella aurita</i> microalgae	Silicon: 3,3 % Crystalline silica: max 0,1-0,3 % as impurity	
Oil enriched with phytosterols/ phytostanols	Crystalline silica: max 0,1-0,3 % as impurity Description/Definition: Oil enriched with phytosterols/phytostanols is composed of an oil fraction and a phytosterol fraction. Acylglycerol Distribution: Free fatty acids (expressed as oleic acid): $\leq 2,0$ % Monoacylglycerols (MAG): ≤ 10 % Diacylglycerols (DAG): ≤ 25 % Triacylglycerols (TAG): Making up the balance Phytosterol fraction: β -sitosterol: ≤ 80 % β -sitostanol: ≤ 15 % campesterol: ≤ 40 % campesterol: ≤ 40 % stigmasterol: $\leq 3,0$ % brassicasterol $\leq 3,0$ % other sterols/stanols: $\leq 3,0$ % Others: Moisture and volatile: $\leq 0,5$ % Peroxide value (PV): $< 5,0$ meq/kg Trans fatty acids: ≤ 1 % Contamination/Purity (GC-FID or equivalent method) of phytosterols/ phytostanols: Phytosterols and phytostanols extracted from sources other than vegetable oil suitable for food have to be free of contaminants, best	
Oil extracted from squids	Acid value: ≤ 0.5 KOH/g oil Peroxide value (PV): ≤ 5 meq O ₂ /kg oil p-Anisidine value: ≤ 20	

	Cold test at 0 °C: Moisture: $\leq 0,1 \%$ Unsaponifiable m Docosahexaeonic Eicosapentaenoic	a (w/w) atter: $\leq 5,0$ %Trans acid: ≥ 20 %	s fatty acids: $\leq 1,0$ S	%
[^{F5} Partially defatted chia seed (<i>Salvia</i> <i>hispanica</i>) powders	 Description/Definition: The novel foods are partially defatted chia seed (<i>Salvia</i> powders obtained by pressing and grinding of the whole <i>hispanica</i> L. Physical-sensorial: Foreign matter: 0,1 % 			
		Powder with high protein content	Powder with high fibre content	
	Particle size	\leq 130 μ m	\leq 400 μ m	
	Chemical compo	sition:		
		Salvia hispanica powder with high protein content	Salvia hispanica powder with high fibre content	
	Moisture	≤9,0 %	≤9,0 %	
	Protein	≥ 40,0 %	≥24,0 %	
	Fat	≤17 %	≤ 12 %	
	Fibre	\leq 30 %	≥ 50 %	
	Coliforms: < 100 Enterobacteriacea Bacillus cereus : Escherichia coli :	$\leq 10\ 000\ CFU/g$ U/g FU/g $ureus : \leq 10\ CFU/g$ $me: \leq 100\ CFU/g$ $\leq 50\ CFU/g$ $\leq 10\ MPN/g$ $\log enes : Absence/g$ Absence in 25 g m ppm $\leq 4\ ppb$		
Pasteurised	Parameter	Target	Comments	
fruit-based preparations produced using	Fruit storage before high-	Minimum 15 days at – 20 °C	Fruit harvested an conjunction with	

high-pressure treatment	pressure treatment		agricultural and manufacturing practices
	Fruit added	40 % to 60 % of thawed fruit	Fruit homogenised and added to other ingredients
	pН	3,2 to 4,2	
	° Brix	7 to 42	Assured by added sugars
	a w	< 0,95	Assured by added sugars
	Final storage	60 days maximum at + 5 °C maximum	Equivalent to storage regimen for conventionally processed product

[^{F27}PhenylcapsaiciPescription/Definition:

	Phenylcapsaicin (N -[(4-hydroxy-3-methoxyphenyl)methyl]-7- phenylhept-6-ynamide, C ₂₁ H ₂₃ NO ₃ , CAS no: 848127-67-3), is synthesized chemically via a two step synthesis process involving in a first step the production of the acetylenic acid intermediate through a reaction of phenyl acetylene with a carboxylic acid derivative, and in a second step a series of reactions of the acetylenic acid intermediate with vanillylamine derivative to produce phenylcapsaicin. Characteristics/Composition: Purity (% of dry matter): \geq 98 % Moisture: \leq 0,5 % Total synthesis related production by-products: \leq 1,0 % N,N -dimethyl formamide: \leq 880 mg/kg Dichloromethane: \leq 600 mg/kg Ethyl acetate: \leq 0,5 % Other solvents: \leq 0,5 % Heavy metals: Lead: \leq 1,0 mg/kg Mercury: \leq 0,1 mg/kg Microbiological criteria: Total plate count: \leq 10 CFU/g Coliforms: \leq 10 CFU/g Coliforms: \leq 10 CFU/g Yeast and mould: \leq 10 CFU/g CFU: Colony Forming Units]
Phosphated maize starch	Description/Definition: Phosphated maize starch (phosphated distarch phosphate) is a chemically modified resistant starch derived from high amylose starch by combining chemical treatments to create phosphate cross-links between carbohydrate residues and esterified hydroxyl groups. The novel food ingredient is a white or nearly white powder. CAS No: 11120-02-8 Chemical formula: $(C_6 H_{10} O_5)_n [(C_6 H_9 O_5)_2 PO_2 H]x [(C_6 H_9 O_5)_2 PO_3 H_2]y$

	Document Generated: 2024-06-
known to be in f	Status: Point in time view as at 31/12/2020. ation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes force on or before 27 June 2024. There are changes that may be brought into force at a future date. een made appear in the content and are referenced with annotations. (See end of Document for details)
Changes that have b	een made appear in the content and are rejerenced with annotations. (See end of Document for details)
	n = number of glucose units; x, y = degrees of substitution The chemical characteristics of phosphated distarch phosphate: Loss on drying: 10-14 % pH: 4,5-7,5 Dietary fibre: \geq 70 % Starch: 7-14 % Protein: \leq 0,8 % Lipids: \leq 0,8 % Residual bound phosphorus: \leq 0,4 % (as phosphorus) 'high amylose maize' as source
from fish phospholipids	rinDescription/Definition: The novel food ingredient is yellow to brown powder. Phosphatidylserine is obtained from fish phospholipids by an enzymatic transphosphorylation with the amino acid L-serine. Specification of the phosphatidylserine product manufactured from fish phospholipids: Moisture: $< 5,0 \%$ Phospholipids: $\geq 75 \%$ Phosphatidylserine: $\geq 35 \%$ Glycerides: $< 4,0 \%$ Free L-serine: $< 1,0 \%$ Tocopherols: $< 0,5 \% (^1)$ Peroxide value (PV): $< 5.0 \text{ meg } \Omega_{2}/kg$
	Peroxide value (PV): $< 5,0 \text{ meq O}_2/\text{kg}$
	(¹) Tocopherols may be added as antioxidants according to Commission Regulation (EU) No 1129/2011
Phosphatidylse	rinDescription/Definition:
from soya phospholipids	The novel food ingredient is off-white to light yellow powder. It is also available in liquid form with a clear brown to orange colour. The liquid form contains medium chain triacylglycerides (MCT) as a carrier. It contains lower levels of Phosphatidylserine due to the fact that it includes significant amounts of oil (MCT). Phosphatidylserine from soya phospholipids is obtained through enzymatic transphosphatidylation of high-phosphatidylcholine soybean lecithin with the amino acid L-serine. Phosphatidylserine consists of a glycerophosphate skeleton conjugated with two fatty acids and L-serine via a phosphodiester linkage.
	Characteristics of Phosphatidylserine from soya phospholipids:
	Powder form:
	Moisture: $< 2,0 \%$
	Phospholipids: $\geq 85\%$
	Phosphatidylserine: $\geq 61 \%$
	Glycerides: < 2,0 % free L-serine: < 1,0 %
	Tocopherols: $< 0.3 \%$
	Phytosterols: $< 0.2 \%$
	Liquid form:
	Moisture: < 2,0 %
	Phospholipids: ≥ 25 %Phosphatidylserine: ≥ 20 %
	Glycerides: not applicable
	free L-serine: < 1,0 %
	Tocopherols: $< 0.3 \%$
	1000000000000000000000000000000000000

Phytosterols: $< 0,2 \%$)
--------------------------	---

	1 hytosterois. < 0,2 /0
Phospholipid product containing equal amounts of phosphatidylseri and phosphatidic acid	Description/Definition: The product is manufactured through enzymatic conversion of soy lecithin. The phospholipid product is a highly concentrated, yellow- brown powder form of phosphatidylserine and phosphatidic acid at an equal level. nSpecification of the product: Moisture: $\leq 2,0 \%$ Total phospholipids: $\geq 70 \%$ Phosphatidylserine: $\geq 20 \%$ Phosphatidic acid: $\geq 20 \%$ Glycerides: $\leq 1,0 \%$ Free L-serine: $\leq 1,0 \%$ Tocopherols: $\leq 0,3 \%$ Phytosterols: $\leq 2,0 \%$ Silicon dioxide is used with a maximum content of 1,0 %
Phospholipides from egg yolk	85 % and 100 % pure Phospholipides from egg yolk
Phytoglycogen	Description: White to off-white powder which is an odourless, colourless, flavourless polysaccharide derived from non-GM sweet corn using conventional food processing techniques Definition: Glucose polymer (C $_6$ H $_{12}$ O $_6$)n with linear linkages of $\alpha(1 - 4)$ glycosidic bonds branched every 8 to 12 glucose units by $\alpha(1 - 6)$ glycosidic bonds Specifications: Carbohydrates: 97 % Sugars: 0,5 % Fibre: 0,8 % Fat: 0,2 % Protein: 0,6 %
Phytosterols/ phytostanols	Description/Definition: Phytosterols and phytostanols are sterols and stanols that are extracted from plants and may be presented as free sterols and stanols or esterified with food grade fatty acids. Composition (with GC-FID or equivalent method): β -sitosterol: < 81 % β -sitostanol: < 35 % campesterol: < 40 % campestanol: < 15 % stigmasterol: < 30 % brassicasterol: < 3,0 % other sterols/stanols: < 3,0 % Contamination/Purity (GC-FID or equivalent method): Phytosterols and phytostanols extracted from sources other than vegetable oil suitable for food have to be free of contaminants, best ensured by a purity of more than 99 % of the phytosterol/phytostanol ingredient.
Plum kernel oil	Description/Definition: Plum kernel oil is a vegetable oil obtained by cold pressing of plum (<i>Prunus domestica</i>) kernels.

Potato proteins (coagulated) and hydrolysates	Composition:Oleic acid (C18:1): 68 %Linoleic acid (C18:2): 23 % γ -Tocopherol:80 % of total tocopherols β -Sitosterol: 80-90 % of total sterolsTriolein: 40-55 % of triglyceridesCyanhydric acid: maximum 5 mg/kg oilDry substance: \geq 800 mg/gProtein (N * 6,25): \geq 600 mg/g (dry substance)Ash: \leq 400 mg/g (dry substance)Glycoalkaloid (total): \leq 150 mg/kg
thereof	Lysinoalanine (total): \leq 500 mg/kg Lysinoalanine (free): \leq 10 mg/kg
Prolyl oligopeptidase (enzyme preparation)	Specification of the enzyme: Systematic name: Prolyl oligopeptidase Synonyms: Prolyl endopeptidase, proline-specific endopeptidase, endoprolylpeptidase Molecular weight: 66 kDa Enzyme Commission number: EC 3.4.21.26 CAS number: 72162-84-6 Source: A genetically modified strain of <i>Aspergillus niger</i> (GEP-44) Description: Prolyl oligopeptidase is available as an enzyme preparation containing approximately 30 % maltodextrin. Specifications of the enzyme preparation of prolyl oligopeptidase: Activity: > 580 000 PPI(1)/g (> 34,8 PPU(2)/g) Appearance: Microgranulate Colour: Off-white to orange yellowish. The colour may change from batch to batch Dry Matter: > 94 % Gluten: < 20 ppm Heavy metals: Lead: $\leq 1,0$ mg/kg Arsenic: $\leq 1,0$ mg/kg Mercury: $\leq 0,1$ mg/kg Microbiological criteria: Total aerobic plate count: $\leq 10^3$ CFU/g Total yeasts and moulds: $\leq 10^2$ CFU/g
	Sulphite reducing anaerobes: $\leq 30 \text{ CFU/g}$ Enterobacteriaceae : $< 10 \text{ CFU/g}$ Salmonella : Absence in 25 g Escherichia coli : Absence in 25 g Staphylococcus aureus : Absence in 10 g Pseudomonas aeruginosa : Absence in 10 g Listeria monocytogenes : Absence in 25 g Antimicrobial activity: AbsentMycotoxins: Below limits of detection: Aflatoxin B1, B2, G1, G2 ($< 0.25 \mu \text{g/kg}$), total Aflatoxins ($< 2.0 \mu \text{g/kg}$), Ochratoxin A ($< 0.20 \mu \text{g/kg}$), T-2 Toxin ($< 5 \mu \text{g/kg}$), Zearalenone ($< 2.5 \mu \text{g/kg}$), Fumonisin B1 and B2 ($< 2.5 \mu \text{g/kg}$)
	(¹) PPI – Protease Picomole International

(²) PPU – Prolyl Peptidase Units or Proline Protease Units

[^{F28} Protein extract from pig kidneys	Description/Definition: The protein extract is obtained from homogenised pig kidneys through a combination of salt precipitation and high speed centrifugation. The obtained precipitate contains essentially proteins with 7 % of the enzyme diamine oxidase (enzyme nomenclature E.C. 1.4.3.22) and is resuspended in a physiologic buffer system. The obtained pig kidney extract is formulated as encapsulated enteric coated pellets or enteric coated tablets to reach the active sites of digestion. Basic Product: Specification: pig kidney protein excerpt with natural content of Diamine oxidase (DAO): Physical condition: liquid Colour: brownish Appearance: slightly turbid solution pH value: 6,4–6,8 Enzymatic activity: > 2 677 kHDU DAO/ml (DAO REA (DAO Radioextractionassay)) Microbiological criteria: Brachyspira spp:: negative (Real Time PCR) Listeria monocytogenes : negative (Real Time PCR) Staphylococcus aureus : < 100 CFU/g Influenza A: negative (Reverse Transcription Real Time PCR) Escherichia coli : < 10 CFU/g Total aerobic microbiological count: < 10 ⁵ CFU/g Yeasts/moulds count: < 10 ⁵ CFU/g Salmonella : Absence/10g Bile salt resistant enterobacteriaceae: < 10 ⁴ CFU/g Final product: Specification pig kidney protein excerpt with natural content of DAO (E.C. 1.4.3.22) in an enteric coated formulation: Physical condition: solid Colour: yellow grey Appearance: micropellets or tablets Enzymatic activity: 110-220 kHDU DAO/g pellet or g tablet (DAO REA (DAO REA (DAO REA) (DAO REA)
	Bile salt resistant enterobacteriaceae: < 10 ⁴ CFU/g Final product:
	(E.C. 1.4.3.22) in an enteric coated formulation:
	Colour: yellow grey
	Enzymatic activity: 110-220 kHDU DAO/g pellet or g tablet (DAO REA
	Acid stability 15 min 0,1M HCl followed by 60 min Borat pH = 9,0: > 68 kHDU DAO/g pellet or g tablet (DAO REA (DAO
	Radioextractionassay)) Humidity: < 10 %
	Staphylococcus aureus : < 100 CFU/g Escherichia coli : < 10 CFU/g
	Total aerobic microbiological count: $< 10^4$ CFU/g
	Total combined yeasts/moulds count: $< 10^{-3}$ CFU/g Salmonella : Absence/10g
	Bile salt resistant enterobacteriaceae: $< 10^{2}$ CFU/g]
[^{F29} Pyrroloquinol quinone disodium salt	iDefinition: Chemical name: disodium 9-carboxy-4,5-dioxo-1 H -pyrrolo[5,4-f]quinoline-2,7-dicarboxylate Chemical formula: C $_{14}$ H $_4$ N $_2$ Na $_2$ O $_8$

	CAS No: 122628-50-6 Molecular weight: 374,17 Da Description Pyrroloquinoline quinone disodium salt is a reddish-brown powder produced by the non-genetically modified bacterium <i>Hyphomicrobium</i> <i>denitrificans</i> strain CK-275. Characteristics/Composition Appearance: Reddish-brown powder Purity: \geq 99,0 % (dry weight) UV absorbance (A322/A259): 0,56 \pm 0,03 UV absorbance (A233/A259): 0,90 \pm 0,09 Moisture: \leq 12,0 % Residual Solvent Ethanol: \leq 0,05 % Heavy metals Lead: $<$ 3 mg/kg Arsenic: $<$ 2 mg/kg Microbiological criteria: Total viable cell count: \leq 300 CFU/g Mould/yeast: \leq 12 CFU/g Coliforms: absent in 1 g <i>Hyphomicrobium denitrificans</i> : \leq 25 CFU/g CFU: Colony Forming Units]
Rapeseed oil high in unsaponifiable matter	Description/Definition: Rapeseed oil high in unsaponifiable matter' is produced by vacuum distillation and it is different from refined rapeseed oil in the concentration of the unsaponifiable fraction (1 g in refined rapeseed oil and 9 g in 'rapeseed oil high in unsaponifiable matter'). There is a minor reduction of triglycerides containing monounsaturated and polyunsaturated fatty acids. Purity: Unsaponifiable matter: > 7,0 g/100 g Tocopherols: > 0,8 g/100 g α -tocopherol (%): 30-50 % γ -tocopherol (%): 50-70 % δ -tocopherol (%): 50-70 % δ -tocopherol (%): < 6,0 % Sterols, triterpenic alcohols, methylsterols: > 5,0 g/100 g Fatty acids in triglycerides: palmitic acid: 3-8 % stearic acid: 0,8-2,5 % oleic acid: 50-70 % linoleic acid: 15-28 % linolenic acid: 6-14 % erucic acid: 2,0 % Acid value: \leq 6,0 mg KOH/g Peroxide value (PV): \leq 10 mEq O 2/kg Heavy metals: Iron (Fe): < 1 000 µg/kg Copper (Cu): < 100 µg/kg Copper (Cu): < 100 µg/kg Polycyclic aromatic hydrocarbons (PAH) Benzo(a)pyrene: < 2 µg/kg

	Treatment with active carbon is required to ensure that polycyclic aromatic hydrocarbons (PAH) are not enriched in the production of 'rapeseed oil high in unsaponifiable matter.
Rapeseed Protein	Definition:Rapeseed protein is an aqueous protein-rich extract from rapeseed press cake originating from non-genetically modified <i>Brassica napus</i> L. and <i>Brassica rapa</i> L.Description:White to off-white, spray dried powder Total protein: ≥ 90 % Soluble protein: ≥ 85 % Moisture: $\leq 7,0$ % Carbohydrates: $\leq 7,0$ % Fat: $\leq 2,0$ % Ash: $\leq 4,0$ % Fibre: $\leq 0,5$ %
[^{F30} Refined shrimp peptide concentrate	DescriptionRefined shrimp peptide concentrate is a peptide mixture obtained from northern shrimp (<i>Pandalus borealis</i>) shells and heads via a series of purification steps following enzymatic proteolysis using a protease from <i>Bacillus licheniformis</i> and/or <i>Bacillus amyloliquefaciens</i> .Characteristics/Composition Total Dry matter (%): $\geq 95,0\%$ Peptides (w/weight dry matter): $\geq 87,0\%$ of which peptides with molecular weight < 2 kDa: $\geq 99,9\%$

	Coagulase positive <i>Staphylococcus aureus</i> : ≤ 200 CFU/g <i>Pseudomonas aeruginosa</i> : ND/25g Mould/yeast: ≤ 20 CFU/g
	CFU: Colony Forming UnitsND: Not Detectable]
Trans- resveratrol	Description/Definition: Synthetic <i>Trans</i> -resveratrol is off-white to beige crystals. Chemical name: 5-[(E)-2-(4-hydroxyphenyl)ethenyl]benzene-1,3-diol Chemical formula: $C_{14} H_{12} O_3$ Molecular weight: 228,25 Da CAS No: 501-36-0 Purity: <i>Trans</i> -resveratrol: \geq 98 %-99 % Total by-products (related substances): \leq 0,5 % Any single related substance: \leq 0,1 % Loss on drying: \leq 0,5 % Heavy metals: Lead: \leq 1,0 ppm Mercury: \leq 0,1 ppm Arsenic: \leq 1,0 ppm Impurities: Diisopropylamine: \leq 50 mg/kg <i>Microbial source</i> : A genetically modified strain of <i>Saccharomyces</i> <i>cerevisiae</i> Appearance: Off-white to slight yellow powder Particle size: 100 % less than 62,23 µm Trans-resveratrol content: Min. 98 % w/w (dry weight basis) Ash: Max. 0,5 % w/w Moisture: Max. 3 % w/w
Rooster comb extract	Description/Definition: Rooster comb extract is obtained from <i>Gallus gallus</i> by enzymatic hydrolysis of rooster comb and by subsequent filtration, concentration and precipitation steps. The principal constituents of rooster comb extract are the glycosaminoglycans hyaluronic acid, chondroitin sulphate A and dermatan sulphate (chondroitin sulphate B). White or almost white hygroscopic powder. Hyaluronic acid: 60-80 % Chondroitin sulphate A: \leq 5,0 %Dermatan sulphate (chondroitin sulphate B): \leq 25 % pH: 5,0-8,5 Purity: Chlorides: \leq 1,0 % Nitrogen: \leq 8,0 % Loss on drying: (105 °C for 6 hours): \leq 10 % Heavy metals: Mercury: \leq 0,1 mg/kg Cadmium: \leq 1,0 mg/kg Chromium: \leq 10 mg/kg Lead: \leq 0,5 mg/kg

	Microbiological criteria:
	Total viable aerobic count: $\leq 10^2$ CFU/g <i>Escherichia coli</i> : Absence in 1 g <i>Salmonella</i> : Absence in 1 g <i>Staphylococcus aureus</i> : Absence in 1 g <i>Pseudomonas aeruginosa</i> : Absence in 1g
Sacha Inchi oil from <i>Plukenetia</i> <i>volubilis</i>	Description/Definition: Sacha inchi oil is a 100 % cold pressed vegetable oil obtained from the seeds of <i>Plukenetia volubiis</i> L. It is a transparent, fluid (liquid) and shiny oil at room temperature. It has a fruity, light, green vegetable taste without undesirable flavours. Aspect, limpidity, shine, colour: Fluid at room temperature, clean, shiny yellow gold Odour and taste: Fruity, vegetable without non acceptable taste or odour Purity: Water and Volatiles: < 0,2 g/100 g Impurities insoluble in hexane: < 0,05 g/100 g Oleic acidity: < 2,0 g/100 g Peroxide value (PV): < 15 meq O ₂ /kg Trans fatty acids: < 1,0 g/100 g Total unsaturated fatty acids: > 90 %Omega 3 alpha linolenic acid (ALA): > 45 % Saturated fatty acids: < 10 % No trans fatty acids (< 0,5 %) No erucic acid (< 0,2 %) More than 50 % of tri-linolenin and di-linolenin-triglycerides Phytosterols composition and level No cholesterol (< 5,0 mg/100 g)
Salatrims	Description/Definition: Salatrim is the internationally recognised acronym for (short and long chain acyl triglyceride molecules). Salatrim is prepared by non-enzymatic inter-esterification of triacetin, tripropionin, tributyrin, or their mixtures with hydrogenated canola, soybean, cottonseed, or sunflower oil. Description: Clear, slightly amber liquid to a light coloured waxy solid at room temperature. Free of particulate matter and of foreign or rancid odour. Glycerol ester disribution: Triacylglycerols: ≥ 87 % Diacylglycerols: ≥ 10 % Monoacylglycerols: ≤ 2,0 % Fatty acid composition: MOLE % LCFA (long chain fatty acids): 33-70 % MOLE % SCFA (short chain fatty acids): 30-67 % Saturated long chain fatty acids: < 70 % by weight Trans fatty acids: ≤ 1,0 % Free fatty acids as oleic acid: ≤ 0,5 % Triacylglycerol profile: Triesters (short/long of 0,5 to 2,0): ≥ 90 % Triesters (short/long = 0): ≤ 10 % Unsaponifiable material: ≤ 1,0 %

Status: Point in time view as at 31/12/2020.
Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Ash: $\leq 0,1 \%$ Colour: $\leq 3,5 \text{ Red (Lovibond)}$ Peroxide value (PV): $\leq 2,0 \text{ Meq/Kg}$
<i>Schizochytrium</i> <i>sp.</i> oil rich in DHA and EPA	Acid value: $\leq 0,5 \text{ mg KOH/g}$ Peroxide value (PV): $\leq 5,0 \text{ meq/kg oil}$ Oxidative stability: All food products containing <i>Schizochytrium sp.</i> oil rich in DHA and EPA should demonstrate oxidative stability by appropriate and recognised national/international test methodology (e.g. AOAC) Moisture and volatiles: $\leq 0,05 \%$ Unsaponifiables: $\leq 4,5 \%$ Trans-fatty acids: $\leq 1 \%$ DHA content: $\geq 22,5 \%$ EPA content: $\geq 10 \%$
<i>[^{F31}Schizochytrium</i> sp. (ATCC PTA-9695) oil	The novel food is obtained from the strain ATCC PTA-9695 of the microalgae <i>Schizochytrium</i> sp. Peroxide value (PV): \leq 5,0 meq/kg oil Unsaponifiables: \leq 3,5 % Trans-fatty acids: \leq 2,0 % Free fatty acids: \leq 0,4 % Docosapentaenoic acid (DPA) n-6: \leq 7,5 % DHA content: \geq 35 %]
<i>Schizochytrium</i> sp. oil	Acid value: $\leq 0.5 \text{ mg KOH/g}$ Peroxide value (PV): $\leq 5.0 \text{ meq/kg oil}$ Moisture and volatiles: $\leq 0.05 \%$ Unsaponifiables: $\leq 4.5 \%$ Trans-fatty acids: $\leq 1.0 \%$ DHA content: $\geq 32.0 \%$
<i>[^{F43}Schizochytrium sp.</i> (T18) oil	Acid value: ≤ 0.8 mg KOH/g Peroxide value (PV): ≤ 5.0 meq/kg oil Moisture and volatiles: ≤ 0.05 % Unsaponifiables: ≤ 3.5 % Trans-fatty acids: ≤ 2.0 % Free fatty acids: ≤ 0.4 % DHA content: ≥ 35 %]
[^{F33} Syrup from Sorghum bicolor (L.) Moench. (Traditional food from a third country)	Description/Definition The traditional food is syrup from <i>Sorghum bicolor</i> (L.) Moench (genus, <i>Sorghum</i> ; family, <i>Poaceae</i> (alt. <i>Gramineae</i>)). The syrup is obtained from stalks of <i>S. bicolor</i> , after applying production processes such as crushing, extraction, and evaporation including a heat treatment in order to obtain a minimum of 74 °Brix syrup Compositional data of syrup from <i>Sorghum bicolor</i> (L.) Moench Water: 22,7 g/100 g Ash: 2,4 Sugars, total: > 74,0 g/100 g]
Fermented soybean extract	Description/Definition: Fermented soybean extract is an odourless milk-white coloured powder. It is comprised of 30 % fermented soybean extract powder and 70 %

	resistant dextrin (as carrier) from corn-starch, which is added during the processing. Vitamin K $_2$ is removed during the manufacturing process. Fermented soybean extract contains nattokinase isolated from natto, a foodstuff produced by the fermentation of non-genetically modified soybeans (<i>Glycine max</i> (L.)) with a selected strain of <i>Bacillus subtilis</i> var. natto. Nattokinase activity: 20 000 -28 000 Fibrin degradation unit/g(¹) Identity: Confirmable Condition: No offensive taste or smell Loss on drying: $\leq 10 \%$ Vitamin K $_2$: $\leq 0,1$ mg/kg Heavy metals: Lead: $\leq 5,0$ mg/kg Microbiological criteria: Total viable aerobic count: $\leq 10^3$ CFU(³)/g Yeast and mould: $\leq 10^2$ CFU/g Coliforms: ≤ 30 CFU/g Spore-forming bacteria: ≤ 10 CFU/g <i>Escherichia coli</i> : Absence/25 g <i>Listeria</i> : Absence/25 g (¹) Assay method as described by Takaoka et al. (2010).
[^{F44} Spermidine- rich wheat	Description/Definition: Spermidine-rich wheat germ extract is obtained from non-fermented,
germ extract	non-sprouting wheat germs (<i>Triticum aestivum</i>) by the process of solid- liquid extraction targeting specifically, but not exclusively polyamines.
(Triticum aestivum)	Spermidine:(N-(3-aminopropyl)butane-1,4-diamine):0,8-2,4 mg/g
,	Spermine: 0,4-1,2 mg/g
	Spermidine trichloride $< 0,1 \ \mu g/g$ Putrescine: $< 0.3 \ mg/g$
	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\le 16,0 \text{ µg/g}$
	Putrescine: $< 0.3 \text{ mg/g}$ Cadaverine: $\le 16.0 \mu\text{g/g}$ Mycotoxins:
	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\le 16,0 \mu\text{g/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu\text{g/kg}$
	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\le 16,0 \mu\text{g/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu\text{g/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10\ 000\ \text{CFU/g}$
	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\leq 16,0 \text{ µg/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \text{ µg/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$
	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\le 16,0 \mu\text{g/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu\text{g/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$ <i>Escherichia coli:</i> $< 10 \text{ CFU/g}$
	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\leq 16,0 \text{ µg/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \text{ µg/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$
Sucromalt	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\leq 16,0 \mu\text{g/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu\text{g/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$ <i>Escherichia coli:</i> $< 10 \text{ CFU/g}$ <i>Salmonella:</i> Absence/25g <i>Listeria monocytogenes:</i> Absence/25g] Description/Definition:
Sucromalt	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\leq 16,0 \mu g/g$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu g/kg$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$ <i>Escherichia coli:</i> $< 10 \text{ CFU/g}$ <i>Salmonella:</i> Absence/25g <i>Listeria monocytogenes:</i> Absence/25g] Description/Definition: Sucromalt is a complex mixture of saccharides which is produced
Sucromalt	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\le 16,0 \mu\text{g/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu\text{g/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$ <i>Escherichia coli:</i> $< 10 \text{ CFU/g}$ <i>Salmonella:</i> Absence/25g <i>Listeria monocytogenes:</i> Absence/25g] Description/Definition: Sucromalt is a complex mixture of saccharides which is produced from sucrose and a starch hydrolysate by means of an enzymatic reaction. In this process, glucose units are attached to saccharides
Sucromalt	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\le 16,0 \mu\text{g/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu\text{g/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$ <i>Escherichia coli:</i> $< 10 \text{ CFU/g}$ <i>Salmonella:</i> Absence/25g <i>Listeria monocytogenes:</i> Absence/25g] Description/Definition: Sucromalt is a complex mixture of saccharides which is produced from sucrose and a starch hydrolysate by means of an enzymatic reaction. In this process, glucose units are attached to saccharides from the starch hydrolysate by means of an enzyme produced by the
Sucromalt	Putrescine: < 0,3 mg/g Cadaverine: $\leq 16,0 \ \mu g/g$ Mycotoxins: Aflatoxins (total): < 0,4 $\mu g/kg$ Microbiological criteria: Total aerobic bacteria: < 10 000 CFU/g Yeast and moulds: < 100 CFU/g <i>Escherichia coli:</i> < 10 CFU/g <i>Salmonella:</i> Absence/25g <i>Listeria monocytogenes:</i> Absence/25g] Description/Definition: Sucromalt is a complex mixture of saccharides which is produced from sucrose and a starch hydrolysate by means of an enzymatic reaction. In this process, glucose units are attached to saccharides from the starch hydrolysate by means of an enzyme produced by the bacterium <i>Leuconostoc citreum</i> or by means of a recombinant strain
Sucromalt	Putrescine: $< 0,3 \text{ mg/g}$ Cadaverine: $\le 16,0 \mu\text{g/g}$ Mycotoxins: Aflatoxins (total): $< 0,4 \mu\text{g/kg}$ Microbiological criteria: Total aerobic bacteria: $< 10 000 \text{ CFU/g}$ Yeast and moulds: $< 100 \text{ CFU/g}$ <i>Escherichia coli:</i> $< 10 \text{ CFU/g}$ <i>Salmonella:</i> Absence/25g <i>Listeria monocytogenes:</i> Absence/25g] Description/Definition: Sucromalt is a complex mixture of saccharides which is produced from sucrose and a starch hydrolysate by means of an enzymatic reaction. In this process, glucose units are attached to saccharides from the starch hydrolysate by means of an enzyme produced by the

Status: Point in time view as at 31/12/2020.

<i>Changes to legislation:</i> Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)	
	these oligosaccharides, contains mainly fructose but also the disaccharide leucrose and other disaccharides. Total solids: 75-80 % Moisture: 20-25 % Sulphatase: Max 0,05 % pH: 3,5-6,0 Conductivity < 200 (30 %) Nitrogen < 10 ppm Fructose: 35-45 % d.w. Leucrose: 7-15 % d.w. Other disaccharides: Max 3 % Higher saccharides: 40-60 % d.w
Sugar cane fibre	Description/Definition: Sugar Cane Fibre is derived from the dry cell wall or fibrous residue remaining after expression or extraction of sugar juice from sugar cane, of the Saccharum genotype. It consists primarily of cellulose and hemicellulose. The production process consists of several steps, including: chipping, alkaline digestion, removal of lignins and other non- cellulosic components, bleaching of purified fibres, acid washing and neutralization.Moisture: $\leq 7,0 \%$ Ash: $\leq 0,3 \%$ Total Dietary Fibre (AOAC) dry basis (all insoluble): $\geq 95 \%$ of which: Hemicellulose (20-25 %) and cellulose (70-75 %) Silica (ppm): ≤ 200 Protein: 0,0 % Fat: Trace pH: 4-7 Heavy metals: Mercury (ppm): $\leq 0,1$ Lead (ppm): $\leq 1,0$ Cadmium (ppm): $\leq 0,1$ Microbiological criteria: Yeast and moulds (CFU/g): $\leq 1 000$ Salmonella : Absence Listeria monocytogenes : Absence
[^{F34} Sugars obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp	Description/Definition: Sugars are obtained from the concentrated cocoa pulp (<i>Theobroma</i> <i>cacao</i> L.) juice either via a drying process or via a purification process to produce high purity glucose or fructose. Sugars produced by a drying process Nutritional composition: Total sugars (g/100g): > 80 Moisture (%): < 5 Microbiological criteria: Total Plate Count (aerobic) (cfu/g): < 10 ⁴ Moulds and Yeasts (cfu/g): < 50 Enterobacteriaceae (cfu/g): < 10 <i>Salmonella</i> spp.: Absence in 25 g <i>Alicyclobacillus:</i> Absence in 50 g Thermo-acidophilic bacteria: Absence in 50 g

Status: Point in time view as at 31/12/2020.

	Sugars produced by a purification process Nutritional composition of Glucose obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp: Glucose content (%): > 93 Ash (%): < 0,2 Moisture (%): < 1,0 Nutritional composition of Fructose obtained from cocoa (<i>Theobroma</i> <i>cacao</i> L.) pulp: Fructose content (%): > 98 Glucose content (%): < 0,5 % Ash (%): < 0,2 Moisture (%):< 0,5 Microbiological criteria for glucose and fructose obtained from cocoa (<i>Theobroma</i> cacao L.) pulp: Total Plate Count (aerobic) (cfu/g): < 10 ⁴ <i>Salmonella</i> spp.: Absence in 25 g]
Sunflower oil extract	Description/Definition: The sunflower extract is obtained by a concentration factor of 10 of the unsaponifiable fraction of refined sunflower oil extracted from the seeds of the sunflower, <i>Helianthus Annuus</i> L. Composition: Oleic acid (C18:1): 20 % Linoleic acid (C18:2): 70 % Unsaponifiable matter: 8,0 % Phytosterols: 5,5 % Tocopherols: 1,1 %
Dried <i>Tetraselmis</i> <i>chuii</i> microalgae	Description/Definition: The dried product is obtained from the marine microalgae <i>Tetraselmis chuii</i> , belonging to the <i>Chlorodendraceae</i> family, cultivated in sterile sea water in closed photobioreactors insulated from the outside air. Purity/Composition: Identified by means of nuclear marker rDNA 18 S (sequence analysed no less than 1 600 base pairs) in the National Centre for Biotechnology information (NCBI) database: Not less than 99,9 % Humidity: $\leq 7,0 \%$ Proteins: 35-40 % Ashes: 14-16 % Carbohydrates: 30-32 % Fibre: 2-3 % Fat: 5-8 % Saturated fatty acids: 29-31 % of total fatty acids Monounsaturated fatty acids: 21-24 % of total fatty acids Polyunsaturated fatty acids: 44-49 % of total fatty acids Iodine: $\leq 15 \text{ mg/kg}$
<i>Therapon barcoo /</i> Scortum	Description/Definition: Scortum/ <i>Therapon barcoo</i> is a species of fish in the family Terapontidae. It is an endemic fresh water species from Australia. It is now reared in fish farms. Taxonomic Identification: Class: Actinopterygii > order: Perciformes > family: Terapontidae > genus: <i>Therapon</i> or <i>Scortum barcoo</i> Composition of fish flesh:

	Protein (%): 18-25 Moisture (%): 65-75 Ash (%): 0,5-2,0 Energy (KJ/Kg): 6000-11500 Carbohydrates (%): 0,0 Fat (%): 5-15 Fatty acids (mg FA/g fillet): Σ PUFA n-3: 1,2-20,0 Σ PUFA n-6: 0,3-2,0 PUFA n-6: 1,5-15,0 Total omega 3 acids: 1,6-40,0 Total omega 6 acids: 2,6-10,0
D-Tagatose	Description/Definition: Tagatose is produced by isomerization of galactose by means of chemical or enzymatic conversion, or by epimerization of fructose by means of enzymatic conversion. These are single-step conversions. Appearance: White or almost white crystals Chemical name: D-tagatose Synonym: D- <i>lyxo</i> -Hexulose CAS number: 87-81-0 Chemical formula: C ₆ H ₁₂ O ₆ Formula weight: 180,16 (g/mol) Purity: Assay: \geq 98 % on a dry weight basis Loss on drying: \leq 0,5 % (102 °C, 2 hours) Specific Rotation: $[\alpha]_D^{-20}$: -4 to -5 ,6° (1 % aqueous solution)(¹) Melting range: 133–137 °C Heavy metals: Lead: \leq 1,0 mg/kg(*)
	(*) Determine using an atomic absorption technique appropriate to the specified level. The selection of sample size and method of sample preparation may be based on the principles of the method described in FNP 5. 'Instrumental methods'(¹).
	(¹) Food and nutrition paper 5 Rev 2 – Guide to specifications for general notices, general analytical techniques, identification tests, test solutions and other reference materials (JECFA) 1991, 307 p.; English – ISBN 92-5-102991-1
[^{F18} Taxifolin- rich extract]	Description: Taxifolin-rich extract from the wood of Dahurian Larch (<i>Larix gmelinii</i> (Rupr.) Rupr) is a white to pale-yellow powder that crystallizes from hot aqueous solutions. [^{F18} Definition: Chemical name: [(2R,3R)-2-(3,4 dihydroxyphenyl)-3,5,7-trihydroxy-2,3-dihydrochromen-4-one, also called (+) trans (2R,3R)- dihydroquercetin] and with no more than 2 % of the cis-form] Specifications: <i>Physical parameter</i> Moisture: ≤ 10 % <i>Compound analysis</i> Taxifolin (m/m): $\geq 90,0$ % of the dry weight <i>Heavy Metals, Pesticide</i>

Status: Point in time view as at 31/12/2020.

	Residual solvents Ethanol: < 5 000 r Microbiological c Total Plate Count Enterobacteria: ≤ Yeast and Mould: Escherichia coli : Salmonella : Abse Staphylococcus au Pseudomonas : Ab	g/kg ng/kg g/kg trichloroethane (DDT): $\leq 0,05$ mg/kg ng/kg riteria (TPC): $\leq 10^4$ CFU/g 100/g ≤ 100 CFU/g Absence/1 g unce/10 g ureus : Absence/1 g
	Extract component	Content, usual observed range (%)
	Taxifolin	90 - 93
	Aromadendrin	2,5 - 3,5
	Eriodictyol	0,1-0,3
	Quercetin	0,3 - 0,5
	Naringenin	0,2 - 0,3
	Kaempferol	0,01 - 0,1
	Pinocembrin	0,05 - 0,12
	Unidentified flavonoids	1 – 3
	Water(*)	1,5
	crystal.	n in its hydrated form and during the drying process is a This results on the inclusion of water of crystallisation ntity of 1,5 %.
Trehalose	by an α -1,1-glucos sucrose by a multi dihydrate. Virtuall taste Synonyms: α,α -tree Chemical name: α CAS No.: 6138-23 Chemical formula Formula weight: 3 Assay: \geq 98 % on Determine using a	is accharide that consists of two glucose moieties linkes sidic bond. It is obtained from liquefied starch or from is tep enzymatic process. The commercial product is the ly odourless, white or almost white crystals with a sweet ehalose a-D-glucopyranosyl- α -D-glucopyranoside, dihydrate 3-4 (dihydrate) : C $_{12}$ H $_{22}$ O $_{11} \cdot 2$ H $_{2}$ O (dihydrate) 378,33 (dihydrate)

	preparation may be based on the principles of the method described in FNP 5 (1), 'Instrumental methods'
	Method of assay: Principle: trehalose is identified by liquid chromatography and quantified by comparison to a reference standard containing standard trehalose Preparation of sample solution: weigh accurately about 3 g of dry sample into a 100 ml volumetric flask and add about 80 ml of purified, deionised water. Bring sample to complete dissolution and dilute to mark with purified deionised water. Filter through a 0,45 micron filter Preparation of standard solution: dissolve accurately weighed quantities of dry standard reference trehalose in water to obtain a solution having known concentration of about 30 mg of trehalose per ml. Apparatus: liquid chromatography equipped with a refractive index detector and integrating recorder Conditions:
	Column: Shodex Ionpack KS-801 (Showa Denko Co.) or equivalent — length: 300 mm — diameter: 10 mm — temperature: 50 °C Mobile phase: water flow rate: 0,4 ml/min Injection volume: 8 μ l Procedure: inject separately equal volumes of the sample solution and the standard solution into the chromatograph. Record the chromatograms and measure the size of response of the trehalose peak Calculate the quantity, in mg, of trehalose in 1 ml of the sample solution by the following formula:% trehalose = $100 \times (R_U/R_S) (W_S/W_U)$ where
	R _S = peak area of trehalose in the standard preparation R _U = peak area of trehalose in the sample preparation W _S = weight in mg of trehalose in the standard preparation W _U = weight of dry sample in mg Characteristics: Identification: Solubility: Freely soluble in water, very slightly soluble in ethanol Specific rotation: $[α]_D^{20} = +179^\circ$ (5% aqueous solution, dihydrate), $+199^\circ$ (5% aqueous solution, anhydrous substance) Melting point: 97 °C (dihydrate) Purity: Loss on drying: ≤ 1,5% (60 °C, 5h) Total ash: ≤ 0,05% Heavy metals: Lead: ≤ 1,0 mg/kg
[^{F18} UV-treated mushrooms (Agaricus bisporus)	Description/Definition Commercially grown <i>Agaricus bisporus</i> to which UV light treatment is applied to harvested mushrooms. UV radiation: a process of radiation in ultraviolet light within the wavelength of 200-800 nm. Vitamin D ₂

Status:	Point in	time view as	at 31/12/2020.

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Chemical name: $(3\beta,5Z,7E,22E)$ -9,10-secoergosta-5,7,10(19),22- tetraen-3-ol Synonym: Ergocalciferol CAS No: 50-14-6 Molecular weight: 396,65 g/mol Contents Vitamin D ₂ in the final product: 5-20 µg/100 g fresh weight at the expiration of shelf life.]
[^{F18} UV-treated baker's yeast (<i>Saccharomyces</i> <i>cerevisiae</i>)	Description/Definition Baker's yeast (<i>Saccharomyces cerevisiae</i>) is treated with ultraviolet light to induce the conversion of ergosterol to vitamin D ₂ (ergocalciferol). Vitamin D ₂ content in the yeast concentrate varies between 800 000-3 500 000 IU vitamin D/100 g (200-875 μ g/g). The yeast may be inactivated. The yeast concentrate is blended with regular baker's yeast in order not to exceed the maximum level in the pre-packed fresh or dry yeast for home baking. Tan-coloured, free-flowing granules. Vitamin D ₂ Chemical name: (5Z,7E,22E)-(3S)-9,10-secoergosta-5,7,10(19),22- tetraen-3-ol Synonym: Ergocalciferol CAS No.: 50-14-6 Molecular weight: 396,65 g/mol Microbiological criteria for the yeast concentrate Coliforms: $\leq 10^3/g$ <i>Escherichia coli</i> : $\leq 10/g$ <i>Salmonella</i> : Absence in 25 g]
UV-treated bread	Description/Definition: UV-treated bread is yeast leavened bread and rolls (without toppings) to which a treatment with ultraviolet radiation is applied after baking in order to convert ergosterol to vitamin D ₂ (ergocalciferol). UV radiation: A process of radiation in ultraviolet light within the wavelength of 240-315 nm for maximum of 5 seconds with energy input of 10-50 mJ/cm ² . Vitamin D ₂ : Chemical name: (5Z,7E,22E)-3S-9,10-secoergosta-5,7,10(19),22- tetraen-3-ol Synonym: Ergocalciferol CAS No: 50-14-6 Molecular weight: 396,65 g/mol Contents: Vitamin D ₂ (ergocalciferol) in the final product: 0,75-3 µg/100 g(¹) Yeast in dough: 1-5 g/100 g (²) (¹) EN 12821, 2009, European Standard. (²) Recipe calculation.
UV-treated milk	Description/Definition:

	UV-treated milk is cow's milk (whole and semi-skimmed) to which a treatment with ultraviolet (UV) radiation via turbulent flow is applied after pasteurisation. The treatment of the pasteurised milk with UV radiation results in an increase in the vitamin D ₃ (cholecalciferol) concentrations by conversion of 7-dehydrocholesterol to vitamin D ₃ . UV radiation: A process of radiation in ultraviolet light within the wavelength of 200-310 nm with energy input of 1 045 J/l. Vitamin D ₃ : Chemical name: (1S,3Z)-3-[(2E)-2-[(1R,3aS,7aR)-7a-methyl-1-[(2R)-6-methylheptan-2-yl]-2,3,3a,5,6,7-hexahydro-1H-inden-4-ylidene]ethylidene]-4-methylidenecyclohexan-1-ol Synonym: Cholecalciferol CAS No: 67-97-0 Molecular weight: 384,6377 g/mol Contents: Vitamin D ₃ in the final product: Whole milk(¹)0,5-3,2 µg/100 g(²) Semi-skimmed milk(1): 0,1–1,5 µg/100 g(²)
	 (1) As defined by Regulation (EU) No 1308/2013 of the European Parliament and of the Council of 17 December 2013 establishing a common organisation of the markets in agricultural products and repealing Council Regulations (EEC) No 922/72, (EEC) No 234/79, (EC) No 1037/2001 and (EC) No 1234/2007 (OJ L 347, 20.12.2013, p. 671). (2) HPLC
[^{F7} Vitamin D 2 mushroom powder	Description/Definition Vitamin D ₂ mushroom powder is a granular powder made from homogenised <i>Agaricus bisporus</i> mushrooms that have been exposed to UV light. The mushrooms are washed, homogenised and suspended in water to produce a mushroom slurry. The mushroom slurry is passed under a UV lamp. The slurry is then filtered, dried and ground, producing vitamin D ₂ mushroom powder. UV radiation: A process of radiation in ultraviolet light within a range of wavelength similar to those UV-treated novel foods authorised under the novel food regulation. Characteristics/Composition Vitamin D ₂ content: 1 000–1 300 µg/g of mushroom powder ^I Moisture: $\leq 10,0 \%$ Ash: $\leq 13,5 \%$ Heavy Metals Lead (as Pb): $\leq 0,5 \text{ mg/kg}$ Mercury: $\leq 0,1 \text{ mg/kg}$ Arsenic: $\leq 0,3 \text{ mg/kg}$ Mycotoxins Aflatoxins (sum of B1+B2+G1+G2): $< 4 \mu g/kg$ Microbiological criteria: Total plate count: $\leq 5 000 \text{ CFU}^{g}$

	Yeast and mould: $\leq 100 \text{ CFU/g}$ Salmonella sp.: Absent in 25 g Staphylococcus aureus : $\leq 10 \text{ CFU/g}$ Escherichia coli : $\leq 10 \text{ CFU/g}$ Coliforms: $\leq 10 \text{ CFU/g}$ Enterobacteriaceae : $\leq 10 \text{ CFU/g}$ Listeria monocytogenes : Absent in 25 g]
Vitamin K ₂ (menaquinone)	This novel food is produced by a synthetic or microbiological process. Vitamin K ₂ (2-methyl-3-all-trans-polyprenyl-1,4-naphthoquinones), or the menaquinone series, is a group of prenylated naphthoquinone derivatives. The number of isoprene residues, where 1 isoprene unit consists of 5 carbons comprising the side chain, is used to characterise the menaquinone homologues containing primarily MK-7 and to a smaller extent MK-6. Vitamin K ₂ (menaquinones) series with menaquinone-7 (MK-7)(n = 6) being C ₄₆ H ₆₄ O ₂ , menaquinone-6 (MK-6)(n = 5) being C ₄₁ H ₅₆ O ₂ and menaquinone-4 (MK-4)(n = 3) being C ₃₁ H ₄₀ O ₂ . Chemical Name: (all-E)-2-(3,7,11,15,19,23,27- Heptamethyl-2,6,10,14,18,22,26-octacosaheptaenyl)-3-methyl-1,4- naphtalenedione CAS Number: 2124-57-4 Molecular formula: C ₄₆ H ₆₄ O ₂ Molecular weight: 649 g/mol
	Specification of synthetic Vitamin K 2 (menaquinone-7)Appearance: Yellow powderPurity: Max 6,0 % cis-isomer, max 2,0 % other impuritiesContent: 97-102 % Menaquinone-7 (including at least 92 % all-transMenaquinone-7)Specifications of microbiologically produced Vitamin K 2(menaquinone-7)Source: Bacillus subtilis spp. natto and Bacillus licheniformisAppearance: Yellow powder or oil suspension
Wheat bran extract	Description/Definition: White crystalline powder obtained by enzymatic extraction from <i>Triticum aestivum</i> L. bran, rich in arabinoxylan oligosaccharides Dry matter: Min. 94 % Arabinoxylan oligosaccharides: Min 70 % of dry matter

	Average degree of polymerisation of arabinoxylan oligosaccharides: 3-8 Ferulic acid (bound to arabinoxylan oligosaccharides): 1-3 % of dry matter Total poly/oligosaccharides: Min 90 % Protein: Max 2 % of dry matter Ash: Max 2 % of dry matter Microbiological parameters: Mesophilic bacteria – total count: Max 10 000/g Yeasts: Max 100/g Fungi: Max 100/g <i>Salmonella</i> : Absence in 25g <i>Bacillus cereus</i> : Max 1000/g <i>Clostridium perfringens</i> : Max 1000/g					
[^{F45} Xylo- oligosaccharides	obtained from cor xylanase from <i>Tri</i>	cription: novel food is a mixture of xylo-oligosaccharides (XOS) which are ined from corncobs (<i>Zea mays</i> subsp. <i>mays</i>) via hydrolysis by a nase from <i>Trichoderma reesei</i> followed by a purification process. racteristics/Composition				
	Parameter	Powder form	Powder form 2	Syrup form		
	Moisture (%)	≤ 5,0	≤ 5,0	70-75		
	Protein (g/100 g)	< 0,2				
	Ash (%)	≤ 0,3				
	pН	3,5-5,0				
	Total carbohydrate content (g/100 g)	≥97	≥ 95	≥ 70		
	XOS content (dry basis) (g/100 g)	≥ 95	≥70	≥ 70		
	Other carbohydrates (g/100 g) (^a)	2,5-7,5	2-16	1,5-31,5		
	Monosaccharides total (g/100 g)	0-4,5	0-13	0-29		
	Glucose (g/100 g)	0-2	0-5	0-4		
	Arabinose (g/100 g)	0-1,5	0-3	0-10		
	Xylose (g/100 g)	0-1,0	0-5	0-15		
	Disaccharides total (g/100 g)	27,5-48	25-43	26,5-42,5		

Xylobiose (XOS DP2) (g/100 g)	25-45	23-40	25-40	
Cellobiose (g/100 g)	2,5-3	2-3	1,5-2,5	
Oligosaccharides total (g/100 g)	41-77	36-72	32-71	
xylotriose (XOS DP3) (g/100 g)	27-35	18-30	18-30	
xylotetraose (XOS DP4) (g/100 g)	10-20	10-20	8-20	
xylopentaose (XOS DP5) (g/100 g)	3-10	5-10	3-10	
xylohexaose (XOS DP6) (g/100 g)	1-5	1-5	1-5	
Xyloheptaose (XOS DP7) (g/100 g)	0-7	2-7	2-6	
Maltodextrin (g/100 g) (^b)	0	20-25	0	
Copper (mg/kg)	< 5,0			
Lead (mg/kg)	< 0,5			
Arsenic (mg/kg)	< 0,3			
Salmonella (CFU (°)/25 g)	Negative			
<i>E, coli</i> (MPN (^d)/100 g)	Negative			
Yeast (CFU/g)	< 10			
Mould (CFU/g)	< 10			
DP :	Degree of poly	rmerization		
	rbohydrates ind inose) and cello		arides (glucose, xyloso	
	^b) Maltodextrin content is calculated according to the amount added in the process.			
(^c) CFU: Colony Forming Units.				
(^d) MPN: M	lost Probable N	umber l		

<i>I^{F36}Yarrowia</i> <i>lipolytica</i> yeast biomass	Description/Definition: The novel food is the dried and heat-killed biomass of the yeast <i>Yarrowia lipolytica</i> .
	Characteristics/Composition: Protein: 45-55 g/100 g
	Dietary fibre: 24-30 g/100 g Sugars: < 1,0 g/100 g
	Fat: 7-10 g/100 g
	Total ash: $\leq 12 \%$
	Water content: $\leq 5 \%$
	Dry matter content: \geq 95 %
	Microbiological criteria:
	Total Aerobic Microbial Count: $\leq 5 \times 10^{-3}$ CFU/g
	Total Yeast and Mould Count: $\leq 10^2$ CFU/g Viable <i>Yarrowia lipolytica</i> cells ⁱ : < 10 CFU/g (i.e. limit of detection) Coliforms: ≤ 10 CFU/g
	Salmonella spp.: Absence in 25 g]
X 7 / X /	
Yeast beta- glucans	Description/Definition: Beta-glucans are complex, high molecular mass (100–200 kDa) polysaccharides, found in the cell wall of many yeasts and cereals. The chemical name for 'yeast beta-glucans' is $(1-3),(1-6)$ - β -D-glucans. Beta-glucans consist of a backbone of β -1-3-linked glucose residues that are branched by β -1-6-linkages, to which chitin and mannoproteins are
	linked by β-1-4-bonds.
	Beta-glucans are isolated from yeast <i>Saccharomyces cerevisiae</i> . The tertiary structure of the glucan cell wall of <i>Saccharomyces cerevisiae</i> consists of chains of β-1,3-linked glucose residues, branched by β-1,6-linkages, forming a backbone to which are linked chitin via β-1,4- bonds, β-1,6-glucans and some mannoproteins. This novel food is available in three different forms: soluble, insoluble and insoluble in water, but dispersible in many liquid matrices. Chemical characteristics yeast (<i>Saccharomyces cerevisiae</i>) beta-
	glucans:
	Soluble form: Tatal contradiction > 75.0 /
	Total carbohydrates: $> 75 \%$ Beta-glucans (1,3/1,6): $> 75 \%$
	Ash: < 4,0 %
	Moisture: < 8,0 %
	Protein: < 3,5 %
	Fat: < 10 %
	Insoluble form: Tatal contradiction > 70.9 /
	Total carbohydrates: $> 70 \%$ Beta-glucans (1,3/1,6): $> 70 \%$
	Ash: $\leq 12 \%$
	Moisture: < 8,0 %
	Protein: < 10 %
	Fat: < 20 %
	Insoluble in water, but dispersible in many liquid matrices:
	(1,3)-(1,6)-β-D-Glucans: > 80 % Ash: < 2,0 %
	Moisture: < 6,0 %

Status: Point in time view as at 31/12/2020	
---	--

Changes to legislation: Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes
known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date.
Changes that have been made appear in the content and are referenced with annotations. (See end of Document for details)

	Protein: < 4,0 % Total fat: < 3,0 % <i>Microbiological data for insoluble in water, but dispersible in many</i> <i>liquid matrices:</i> Total plate count: < 1 000 CFU/g Enterobacteriaceae: < 100 CFU/g Total coliforms: < 10 CFU/g Yeast: < 25 CFU/g Mould: < 25 CFU/g <i>Salmonella</i> : Absence in 25 g <i>Escherichia coli</i> : Absence in 1 g <i>Bacillus cereus</i> : < 100 CFU/g <i>Staphylococcus aureus</i> : Absence in 1 g <i>Heavy metals for insoluble in water, but dispersible in many liquid</i> <i>matrices:</i> [^{F15} Lead: < 0,2 mg/kg Arsenic: < 0,2 mg/kg Mercury: < 0,1 mg/kg]
Zeaxanthin	Description/Definition: Zeaxanthin is a naturally occurring xanthophyll pigment, it is an oxygenated carotenoid. The synthetic zeaxanthin is presented either as a spray-dried powder of gelatin or starch base ('beadlets') with added α -tocopherol and ascorbyl palmitate or as a corn oil suspension with added α -tocopherol. Synthetic zeaxanthin is produced by a multi-step chemical synthesis from smaller molecules. Orange-red crystalline powder with little or no odour. Chemical formula: C ₄₀ H ₅₆ O ₂ CAS No: 144-68-3 Molecular weight: 568,9 daltons Physical-chemical properties: Loss on drying: < 0,2 % <i>All</i> -trans zeaxanthin: > 96 % Cis-zeaxanthin: < 2,0 % Other carotenoids: < 1,5 % Triphenylphosphine oxid (CAS No 791-28-6): < 50 mg/kg
Zinc L-pidolate	Description/Definition: Zinc L-pidolate is a white to off-white powder, with characteristic odour. International non-proprietary name (INN): L-pyroglutamic acid, Zinc salt Synonyms: Zinc 5-oxoproline, Zinc pyroglutamate, Zinc pyrrolidone carboxylate, Zinc PCA, L-Zinc pidolate CAS No.: 15454-75-8 Molecular formula: (C $_5$ H $_6$ NO $_3$) $_2$ Zn Relative anhydrous molecular mass: 321,4 Appearance: White to slightly white powder Purity: Zinc L-pidolate (purity): \ge 98 % pH (10 % aqueous sol.): 5,0-6,0 Specific rotation: 19,6 ° - 22,8 ° Water: \le 10,0 %

> Glutamic acid: < 2,0 % **Heavy metals:** Lead: $\leq 3,0$ ppm Arsenic: $\leq 2,0$ ppm Cadmium: $\leq 1,0$ ppm Mercury: $\leq 0,1$ ppm **Microbiological criteria:** Total viable mesophilic count: $\leq 1\ 000\ CFU/g$ Yeasts and moulds: $\leq 100\ CFU/g$ Pathogen: Absence

- a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).
- **b** Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).
- c [^{F12}OSC-DMAC (4-dimethylaminocinnamaldehyde) method (Ocean Spray Cranberries, Inc) Martin MA, Ramos S, Mateos R, Marais JPJ, Bravo-Clemente, L, Khoo C and Goya L. Food Res Intl 2015 71: 68-82. Modified from Cunningham DG, Vannozzi S, O'Shea E, Turk R (2002) In: Ho C-T, Zheng QY (eds) Quality Management of Nutraceuticals ACS Symposium series 803, Washington DC. *Quantitation of PACs by DMAC Color Reaction pp* 151-166.
- **d** BL-DMAC 4-dimethylaminocinnamaldehyde) method (Brunswick Lab) Multi-laboratory validation of a standard method for quantifying proanthocyanidins in cranberry powders. Prior RL, Fan E, Ji H, Howell A, Nio C, Payne MJ, Reed J. *J Sci Food Agric.* 2010 Jul;90(9):1473-8.
- e The different values for these three parameters are due to the different methods used.
- f GAE: Gallic Acid Equivalents.
- g [^{F38}CFU: Colony Forming Units.]]
- h [^{F8}HPLC/RI: High-performance liquid chromatography coupled with refractive index detection.
- i CFU: Colony-forming unit.]
- **j** [^{F36}To be tested immediately after the heat-treatment step. Measures have to be in place to prevent cross-contamination with viable *Yarrowia lipolytica* cells during packaging and/or storage of the NF.]
- k [^{F20}3'-Fucosyllactose, 2'-Fucosyl-galactose, Glucose, Galactose, Mannitol, Sorbitol, Galactitol, Trihexose, Allo-lactose and other structurally related carbohydrates.]
- I [^{F7}Converted from International Units (IU) using the conversion factor of $0,025 \ \mu g = 1 \ \text{IU.}$]

Textual Amendments

- **F38** Substituted by Commission Implementing Regulation (EU) 2020/1163 of 6 August 2020 authorising the placing on the market of vitamin D2 mushroom powder as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F39** Substituted by Commission Implementing Regulation (EU) 2019/108 of 24 January 2019 authorising the change of specifications of the novel food ingredient lipid extract from Antarctic Krill (Euphausia superba) under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

F40 Substituted by Commission Implementing Regulation (EU) 2019/2165 of 17 December 2019 authorising the change of the specifications of the novel food coriander seed oil from Coriandrum sativum under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

F41 Substituted by Commission Implementing Regulation (EU) 2019/388 of 11 March 2019 authorising the change of the specifications of the novel food 2'-fucosyllactose produced with Escherichia coli

K-12 under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

- **F42** Substituted by Commission Implementing Regulation (EU) 2019/1314 of 2 August 2019 authorising the change of the specifications of the novel food Lacto-N-neotetraose produced with Escherichia coli K-12 under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F43** Substituted by Commission Implementing Regulation (EU) 2020/478 of 1 April 2020 correcting Implementing Regulation (EU) 2017/2470 establishing the Union list of novel foods (Text with EEA relevance).
- **F44** Substituted by Commission Implementing Regulation (EU) 2020/443 of 25 March 2020 authorising the change of the specifications of the novel food spermidine-rich wheat germ extract (Triticum aestivum) under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).
- **F45** Inserted by Commission Implementing Regulation (EU) 2018/1648 of 29 October 2018 authorising the placing on the market of xylo-oligosaccharides as a novel food under Regulation (EU) 2015/2283 of the European Parliament and of the Council and amending Commission Implementing Regulation (EU) 2017/2470 (Text with EEA relevance).

Status:

Point in time view as at 31/12/2020.

Changes to legislation:

Commission Implementing Regulation (EU) 2017/2470, ANNEX is up to date with all changes known to be in force on or before 27 June 2024. There are changes that may be brought into force at a future date. Changes that have been made appear in the content and are referenced with annotations.