Commission Implementing Regulation (EU) 2018/1023 of 23 July 2018 correcting Implementing Regulation (EU) 2017/2470 establishing the Union list of novel foods (Text with EEA relevance)

COMMISSION IMPLEMENTING REGULATION (EU) 2018/1023

of 23 July 2018

correcting Implementing Regulation (EU) 2017/2470 establishing the Union list of novel foods

(Text with EEA relevance)

THE EUROPEAN COMMISSION,

Having regard to the Treaty on the Functioning of the European Union,

Having regard to Regulation (EU) 2015/2283 of the European Parliament and of the Council on novel foods, amending Regulation (EU) No 1169/2011 of the European Parliament and of the Council and repealing Regulation (EC) No 258/97 of the European Parliament and of the Council and Commission Regulation (EC) No 1852/2001⁽¹⁾, and in particular Article 8 thereof,

Whereas:

- Pursuant to Article 8 of Regulation (EU) 2015/2283, the Commission is to establish, by 1 January 2018, the Union list of novel foods authorised or notified under Regulation (EC) No 258/97 of the European Parliament and of the Council⁽²⁾.
- (2) The Union list of novel foods authorised or notified under Regulation (EC) No 258/97 was established by Commission Implementing Regulation (EU) 2017/2470⁽³⁾.
- (3) Pursuant to Article 36 of Regulation (EU) 2015/2283, the new novel food Regulation applies from 1 January 2018. A number of products were authorised or notified under Regulation (EC) No 258/97 during the period between the Standing Committee vote on the Union list on 6 December 2017 and the date of application of Regulation (EU) 2015/2283 on 1 January 2018. These products should therefore be included in the Union list established through Implementing Regulation (EU) 2017/2470.
- (4) On 19 December 2017, the company Demethra Biotech S.r.l. notified the Commission that it placed the novel food '*Echinacea purpurea* extract from cell cultures' on the Union market pursuant to Article 5 of Regulation (EC) No 258/97. This novel food was not included in the Union list. Therefore, a new entry should be added to Tables 1 and 2 of the Annex to Implementing Regulation (EU) 2017/2470.
- (5) On 21 and 22 December 2017, two companies, DuPont Nutrition & Biosciences ApS and FrieslandCampina Nederland BV, notified the Commission that they placed the novel food '2'-Fucosyllactose (microbial source)' on the Union market pursuant to Article 5 of Regulation (EC) No 258/97. '2'-Fucosyllactose (microbial source)' was already included in the Annex to Implementing Regulation (EU) 2017/2470. Those new notifications modify the numerical values of several parameters listed in the

specifications of this novel food and therefore, the entry '2'-Fucosyllactose (microbial source)' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 should be corrected accordingly.

- (6) On 20 December 2017, the company c-LEcta GmbH notified the Commission that it placed the novel food 'Trehalose' on the Union market pursuant to Article 5 of Regulation (EC) No 258/97. 'Trehalose' was included in the Annex to Implementing Regulation (EU) 2017/2470. That new notification concerns a new source of trehalose, sucrose. Therefore, the specifications of the entry 'Trehalose' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 should be corrected accordingly.
- (7) After the publication of Implementing Regulation (EU) 2017/2470, a number of errors or omissions were noted concerning the specifications or the conditions of use of a number of authorised novel foods. Therefore, the Union list established in the Annex to Implementing Regulation (EU) 2017/2470 should be corrected.
- (8) The novel food 'L-Alanyl-L-Glutamine' was authorised under certain conditions of use pursuant to Article 5 of Regulation (EC) No 258/97. The category 'Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen' was erroneously omitted. Therefore, a correction adding 'Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen' as allowed food category in the entry 'L-Alanyl-L-Glutamine' in Table 1 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (9) The novel food 'Glucosamine HCl' was authorised under certain conditions of use pursuant to Article 5 of Regulation (EC) No 258/97. The food category 'Milk-based drinks and similar products intended for young children' was added erroneously and should be deleted from this entry. A correction in the entry 'Glucosamine HCl' in Table 1 of the Annex to Implementing Regulation (EU) 2017/2470 is therefore necessary.
- (10) The novel food 'Lacto-N-neotetraose' was authorised under certain conditions of use and maximum levels by Commission Implementing Decision (EU) 2016/375⁽⁴⁾. The wording 'at concentrations up to 1,2 g/l' was added erroneously and should be removed from the food category 'Milk-based drinks and similar products intended for young children' for this novel food. Therefore, a correction of the entry 'Lacto-N-neotetraose' in Table 1 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (11) The novel food 'Spermidine-rich wheat germ extract (*Triticum aestivum*)' was authorised under certain conditions of use for 'adult population excluding pregnant and lactating women' pursuant to Article 5 of Regulation (EC) No 258/97. However, the exclusion of pregnant and lactating women erroneously did not feature in the Union list. In consequence, the correction of the entry 'Spermidine-rich wheat germ extract (*Triticum aestivum*)' in Table 1 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (12) The novel food 'Antarctic Krill oil from *Euphausia superba*' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 erroneously omitted the following requirement, which should be added: 'Oxidative stability: all food products containing Antarctic Krill oil from *Euphausia superba* should demonstrate oxidative stability

by appropriate and recognised national/international test methodology (e.g. AOAC)'. Therefore, a correction of this entry in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.

- (13) The novel food 'Antarctic Krill oil rich in phospholipids from *Euphausia superba*' was authorised under certain conditions of use by the Finnish competent authorities⁽⁵⁾. The specifications erroneously added the following requirement: 'Oxidative stability: all food products containing Antarctic Krill oil rich in phospholipids from *Euphausia superba* should demonstrate oxidative stability by appropriate and recognised national/ international test methodology (e.g. AOAC)'. This requirement should be removed. Therefore, a correction of the entry 'Antarctic Krill oil rich in phospholipids from *Euphausia superba*' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (14) The novel food 'Chia seeds (*Salvia hispanica*)' was initially authorised under certain conditions of use by Commission Decision 2009/827/EC⁽⁶⁾. The specifications erroneously added the following requirement: '(EU: carbohydrates are available = sugar + starch)'. This requirement should be removed. Therefore, a correction of the entry 'Chia seeds (*Salvia hispanica*)' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (15) The novel food 'Chitosan extract from fungi (*Agaricus bisporus*; *Aspergillus niger*)' was initially authorised under certain conditions of use pursuant to Article 5 of Regulation (EC) No 258/97. The specifications erroneously added the following requirement: 'Fat binding capacity 800 x 9 w/wet weight): pass'. This requirement should be replaced by 'Fat binding capacity 800 x (w/w wet weight): pass'. Therefore, a correction of the entry 'Chitosan extract from fungi (*Agaricus bisporus*; *Aspergillus niger*)' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (16) The novel food 'Citicoline' was authorised under certain conditions of use by Commission Implementing Decision 2014/423/EU⁽⁷⁾. In Table 2 of the Annex to Implementing Regulation (EU) 2017/2470, the specifications of the novel food 'Citicoline' refer to citicoline produced via either a synthetic or a microbial process. After the publication of that Regulation, it became clear that microbial process for the production of citicoline, also involved synthetic process. Thus, the specifications concerning 'Citicoline' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 should be corrected to include only the microbial production process.
- (17) The novel food 'Echinacea angustifolia extract from cell cultures' was initially authorised under certain conditions of use pursuant to Article 5 of Regulation (EC) No 258/97. The specifications erroneously omitted the wording 'description/definition'. Therefore, a correction of the entry 'Echinacea angustifolia extract from cell cultures' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (18) The novel food 'Galacto-oligosaccharide' is included in the Union list established by Implementing Regulation (EU) 2017/2470. The following microbial sources *Pichia pastoris*, *Kluyveromyces lactis*, *Sporobolomyces singularis* and *Papiliotrema terrestris* of the enzyme 'β-galactosidase' were erroneously omitted in the specifications.

Therefore, these sources of β -galactosidase should be added to the entry 'Galactooligosaccharide' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470.

- (19) The novel food 'Vitamin K₂ (menaquinone)' was initially authorised under certain conditions of use by Commission Decision 2009/345/EC⁽⁸⁾. The chemical definition of Vitamin K₂ was added to 'specifications of microbiologically produced vitamin K₂ (menaquinone-7)' but erroneously not added to 'specifications of synthetic vitamin K₂ (menaquinone-7)'. Therefore, a correction of the entry 'Vitamin K₂ (menaquinone)' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (20) The novel food 'Yeast beta-glucans' was authorised under certain conditions of use by Commission Implementing Decision 2011/762/EU⁽⁹⁾. In the specifications, 'Microbiological data' and 'heavy metals' erroneously refer to the three forms of Yeast beta-glucans instead of to the form 'Insoluble in water but dispersible in many liquid matrices'. Therefore, a correction of the entry 'Yeast beta-glucans' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (21) The novel food 'Phytosterols/phytostanols' was authorised under certain conditions of use by Commission Decision 2004/333/EC⁽¹⁰⁾. On 14 April 2016, the company BASF SE Human Nutrition, ENS/HR notified the Commission that it placed the novel food 'Phytosterols/phytostanols' on the Union market in the category 'Food supplement' pursuant to Article 5 of Regulation (EC) No 258/97. The category 'Food supplement' was erroneously omitted. Therefore, a correction adding 'Food supplement' as allowed food category in the entry 'Phytosterols/phytostanols' in Table 1 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (22) The novel food 'Arachidonic acid-rich oil from the fungus *Mortierella alpina*' was authorised under certain conditions of use by Commission Decision 2008/968/EC⁽¹¹⁾. The following non-genetically modified strain 'CBS 210.32' of the fungus *Mortierella alpina* was erroneously not included in the specifications. Therefore, this strain should be added to the entry 'Arachidonic acid-rich oil from the fungus *Mortierella alpina*' in Table 2 of the Annex to Implementing Regulation (EU) 2017/2470.
- (23) The novel food 'Epigallocatechin gallate as a purified extract from green tea leaves (*Camellia sinensis*)' was initially authorised under certain conditions of use pursuant to Article 5 of Regulation (EC) No 258/97. The food category 'Foods fortified in accordance with Regulation (EC) No 1925/2006' was added erroneously and should be deleted from this entry. Furthermore, a correction adding 'Foods' to 'food supplements as defined in Directive 2002/46/EC' as the allowed food category in the entry 'Epigallocatechin gallate as a purified extract from green tea leaves (*Camellia sinensis*)' in Table 1 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.
- (24) The novel food 'Lycopene from tomatoes' was authorised under certain conditions of use pursuant to Article 5 of Regulation (EC) No 258/97. The food category 'Food supplement' was omitted erroneously and should be added to this entry. Therefore, a correction adding 'Food supplement' as allowed food category in the entry 'Lycopene from tomatoes' in Table 1 of the Annex to Implementing Regulation (EU) 2017/2470 is necessary.

- (25) In addition, after the publication of Implementing Regulation (EU) 2017/2470, several typographical errors have been identified in the Annex. While such typographic errors are usually corrected by a corrigendum, for the sake of clarity for economic operators and enforcement authorities, the correction of those typographical errors should be included in this correcting act.
- (26) Given the number of corrections, it is appropriate to replace the whole Annex to Implementing Regulation (EU) 2017/2470.
- (27) The measures provided for in this Regulation are in accordance with the opinion of the Standing Committee on Plants, Animals, Food and Feed,

HAS ADOPTED THIS REGULATION:

Article 1

The Annex to Implementing Regulation (EU) 2017/2470 is replaced by the Annex to this Regulation.

Article 2

This Regulation shall enter into force on the twentieth day following that of its publication in the *Official Journal of the European Union*.

This Regulation shall be binding in its entirety and directly applicable in all Member States.

Done at Brussels, 23 July 2018.

For the Commission The President Jean-Claude JUNCKER

ANNEX

UNION LIST OF NOVEL FOODS

Content of the list

- 1. The Union list shall consist of Tables 1 and 2.
- 2. Table 1 includes the authorised novel foods and contains the following information:

Column 1	: Authorised novel food
Column 2	: Conditions under which the novel food may be used. This column is
	further subdivided into two: Specified food category and Maximum levels
Column 3	: Additional specific labelling requirements
Column 4	: Other requirements

3. Table 2 includes the specifications on novel foods and contains the following information:

Column 1	:	Authorised novel food
Column 2	:	Specifications

TABLE 1: AUTHORISED NOVEL FOODS

Authorised novel food		Conditions under which the novel food may be used		Other requirements
<i>N</i> -Acetyl-D- neuraminic acid	Specified food category Infant and follow-on formulae as defined by Regulation (EU) No 609/2013 ^a Processed cereal-based foods and baby foods for infants and young children as defined by Regulation (EU) No 609/2013	Maximum levels0,05 g/L of reconstituted formula0,05 g/kg for solid foods	requirementsThe designationof the novel foodon the labellingof the foodstuffscontaining itshall be 'N-acetyl-D-neuraminic acid'FoodsupplementscontainingN-acetyl-D-neuraminicacid shall beara statementthat the foodsupplements	
	Foods for special medical purposes for infants and young children as defined by Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the infants and young children for whom the	given to infants, young children and children under 10 years of age where they consume breast milk or other foods with	

Total diet replacement foods for weight control as defined by Regulation (EU) No 609/2013	products are intended but in any case not higher than the maximum levels specified for the category mentioned in the table corresponding to the products. 0,2 g/L (drinks) 1,7 g/kg (bars)	added <i>N</i> -acetyl- D-neuraminic acid within the same twenty four hour period.
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014 ^b	1,25 g/kg	
Unflavoured pasteurised and sterilised (including UHT) milk-based products	0,05 g/L	
Unflavoured fermented milk- based products, heat treated after fermentation, flavoured fermented milk products including heat- treated products	0,05 g/L (beverages) 0,4 g/kg (solids)	
Dairy analogues, including beverage whiteners	0,05 g/L (beverages) 0,25 g/kg (solids)	

	Cereal bars	0,5 g/kg]	
	Table top sweeteners	8,3 g/kg		
	Fruit and vegetable-based drinks	0,05 g/L		
	Flavoured drinks	0,05 g/L		
	Speciality coffee, tea, herbal and fruit infusions, chicory; tea, herbal and fruit infusions and chicory extracts; tea, plant, fruit and cereal preparations for infusions	0,2 g/kg		
	Food Supplements as defined in Directive 2002/46/EC ^e	300 mg/day for general population older than 10 years 55 mg/day for infants 130 mg/day for young children 250 mg/day for children between 3 to 10 years of age		
<i>Adansonia digitata</i> (Baobab) dried fruit pulp	Not specified	·	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Baobab fruit pulp'	
Ajuga reptans	Specified food	Maximum levels		
extract from cell cultures	<i>category</i> Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of a similar extract of the flowering aerial parts of <i>Ajuga reptans</i>		

L-Alanyl-L- Glutamine	Specified food category	Maximum levels		
	Food Supplements as defined in Directive 2002/46/EC			
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013 excluding foods for infants and young children			
	Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen			
Algal oil from the microalgae	Specified food category	Maximum levels of DHA	The designation of the novel food	
Ulkenia sp.	Bakery products (breads, rolls and sweet biscuits)	200 mg/100 g	on the labelling of the foodstuffs containing it shall be 'Oil from the micro-	
	Cereal bars	500 mg/100 g		
	Non-alcoholic beverages (including milk based beverages)	60 mg/100 ml	algae <i>Ulkenia</i> sp.'	
<i>Allanblackia</i> seed oil	Specified food category	Maximum levels	The designation of the novel food	
	Yellow fat spreads and cream based spreads	20 g/100 g	on the labelling of the foodstuffs containing it shall be <i>Allanblackia</i> seed oil'	
<i>Aloe macroclada</i> Baker leaf	Specified food category	Maximum levels		
extract	Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of the similar gel derived <i>from</i>		

Antarctic	Specified food	<i>Aloe vera</i> (L.) Burm. <i>Maximum</i>	The designation
Krill oil from Euphausia superba	category	levels of combined DHA and EPA	of the novel food on the labelling of the foodstuffs
	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g	containing it shall be 'Lipid extract from the crustacean Antarctic Krill (<i>Euphausia</i> superba)'
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g	
	Non-alcoholic beverages Milk-based drinks Dairy analogue drinks	80 mg/100 ml	
	Spreadable fat and dressings	600 mg/100 g	
	Cooking fats	360 mg/100 ml	-
	Breakfast cereals	500 mg/100 g	-
	Bakery products (breads, rolls and sweet biscuits)	200 mg/100 g	
	Nutrition bars/ cereal bars	500 mg/100 g	-
	Food Supplements as defined in Directive 2002/46/EC	3 000 mg/day for the general population 450 mg/day for pregnant and lactating women	
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended	
	Total diet replacement for weight control	250 mg/meal	

	as defined in Regulation (EU) No 609/2013 and meal replacements for weight control Processed cereal-based food and baby food intended for infants and young children covered by Regulation (EU) No 609/2013	200 mg/100 ml		
	Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen			
	Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014			
Antarctic Krill oil rich in phospholipids from <i>Euphausia</i>	Specified food category	Maximum levels of combined DHA and EPA	The designation of the novel food on the labelling of the foodstuffs	
superba	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g	containing it shall be 'Lipid extract from the crustacean	
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g	Antarctic Krill (Euphausia superba)'	
	Non-alcoholic beverages Milk-based drinks	80 mg/100 ml		

Dairy analogue drinks	
Spreadable fat and dressings	600 mg/100 g
Cooking fats	360 mg/100 ml
Breakfast cereals	500 mg/100 g
Bakery products (breads, rolls and sweet biscuits)	200 mg/100 g
Nutrition bars/ cereal bars	500 mg/100 g
Food Supplements as defined in Directive 2002/46/EC	3 000 mg/day for the general population 450 mg/day for pregnant and lactating women
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal
Processed cereal-based food and baby food intended for infants and young children covered by Regulation (EU) No 609/2013 Foods intended	200 mg/100 ml
to meet the expenditure of	

	intense muscular effort, especially for sportsmen Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014			
Arachidonic acid-rich oil	Specified food category	Maximum levels	The designation of the novel food	
from the fungus Mortierella alpina	Infant formula and follow- on formula as defined in Regulation (EU) No 609/2013	In accordance with Regulation (EU) No 609/2013	on the labelling of the foodstuffs containing it shall be 'Oil from <i>Mortierella</i> <i>alpina</i> ' or ' <i>Mortierella</i>	
	Foods for special medical purposes for premature infants as defined in Regulation (EU) No 609/2013	In accordance with Regulation (EU) No 609/2013	alpina oil'	
Argan oil from Argania spinosa	Specified food category	Maximum levels	The designation of the novel food	
9 i	As seasonings	Not specified	on the labelling	
	Food Supplements as defined in Directive 2002/46/EC	In line with normal food use of vegetable oils	of the foodstuffs containing it shall be 'Argan oil' and if used as seasoning 'Vegetable oil only for seasoning' shall be mentioned on the label	
Astaxanthin- rich oleoresin	Specified food category	Maximum levels	The designation of the novel food	
from Haematococcus pluvialis algae	Food Supplements as defined in Directive 2002/46/EC	$\frac{40-80 \text{ mg/day}}{\text{of oleoresin,}}$ resulting in $\leq 8 \text{ mg}$	on the labelling of the foodstuffs containing it shall be 'Astaxanthin'	

Basil seeds (Ocimum basilicum)	Specified food category Fruit juice and fruit/vegetable blend beverages	astaxanthin per day <i>Maximum</i> <i>levels</i> 3 g/200 ml for addition of whole basil seeds (<i>Ocimum</i> <i>basilicum</i>)	_	
Fermented black bean extract	Specified food category Food	Maximum levels4,5 g/day	The designation of the novel food on the labelling of the foodstuffs	
	Supplements as defined in Directive 2002/46/EC		of the foodstuffs containing it shall be 'Fermented black bean (Soya) extract" or 'Fermented Soya extract'	
Bovine lactoferrin	Specified food category	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Lactoferrin from cows' milk'	
lactoierrin	Infant formula and follow- on formula as defined in Regulation (EU) No 609/2013 (ready to drink)	100 mg/100 ml		
	Foods on dairy basis intended for young children (ready to eat/drink)	200 mg/100 g		
	Processed cereal food (solid)	670 mg/100 g		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	Depending on the needs of the individual up to 3 g/day		
	Beverages based on milk	200 mg/100 g		
	Powdered drink mixes based on milk (ready to drink)	330 mg/100 g		

	Beverages based on fermented milk (including yoghurt drinks) Non-alcoholic drinks Products based on yoghurt Products based on cheese Ice cream Cakes and	50 mg/100 g 120 mg/100 g 80 mg/100 g 2 000 mg/100 g 130 mg/100 g 1 000 mg/100 g		
	pastries	1 000 mg/100 g		
	Candies	750 mg/100 g		
	Chewing gum	3 000 mg/100 g		
<i>Buglossoides</i> <i>arvensis</i> seed oil	Specified food category	Maximum levels of stearidonic acid (STA)	The designation of the novel food on the labelling of the foodstuffs	
	Dairy products	250 mg/100 g	containing it shall be 'Refined	
	and analogues	75 mg/100 g for drinks	Buglossoides oil'	
	Cheese and cheese products	750 mg/100 g		
	Butter and other fat and oil emulsions including spreads (not for cooking or frying purposes)	750 mg/100 g		
	Breakfast cereals	625 mg/100 g		
	Food supplements as defined in Directive 2002/46/EC, excluding food supplements for infants and young children	500 mg/day		
	Foods for special medical purposes as defined in	In accordance with the particular nutritional		

Regulation (EU) No 609/2013, excluding foods for special medical purposes intended for infants and young children	requirements of the persons for whom the products are intended		
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal		
Specified food	Maximum	The designation	
Food supplements as defined in Directive 2002/46/EC	2,3 g/day	of the novel food on the labelling of the foodstuffs containing it shall be 'oil from <i>Calanus</i> <i>finmarchicus</i> (crustacean)'	
Specified food	Maximum levels	The designation	
Cathying g um	8%	on the labelling of the foodstuffs containing it shall be 'Gum base (including 1,3-butadiene, 2-methyl- homopolymer, maleated, esters with polyethylene glycol mono-Me ether)' or 'Gum base (including CAS No: 1246080-53-4)'	
<i>Specified food</i> <i>category</i> Chewing gum	Maximum levels 2 %	The designation of the novel food on the labelling of the foodstuffs	
	No 609/2013, excluding foods for special medical purposes intended for infants and young children Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control Specified food category Food supplements as defined in Directive 2002/46/EC Specified food category Vativieg gum	No 609/2013, excluding foods for special medical purposes intended for infants and young childrenof the persons for whom the products are intendedTotal diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control250 mg/mealSpecified food categoryMaximum levelsSpecified food category2,3 g/daySpecified food categoryMaximum levelsSpecified food categoryMaximum levelsSpecified food categoryMaximum levelsSpecified food categoryMaximum levelsSpecified food categoryMaximum levelsSpecified food categoryMaximum levelsSpecified food categoryMaximum levels	No 609/2013, excluding foods for special medical purposes intended for infants and young childrenof the persons for whom the products are intendedTotal diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control250 mg/mealSpecified food category2,3 g/dayThe designation of the novel food on the labelling of the foodstuffs containing it shall be 'oil from Calanus finmarchicus (crustacean)'Specified food categoryMaximum levelsThe designation of the foodstuffs containing it shall be 'oil from Calanus finmarchicus (crustacean)'Specified food categoryMaximum levelsThe designation of the foodstuffs containing it shall be 'oil from Calanus finmarchicus (crustacean)'Specified food category8 %The designation of the foodstuffs containing it shall be 'Gum base (including 1,3-butadiene, 2-methyl- homopolymer, maleated, esters with polyethylene glycol mono-Me ether)' or 'Gum base (including CAS No: 1246080-53-4)'Specified food categoryMaximum levelsThe designation of the novel food

			base (including methyl vinyl ether-maleic anhydride copolymer)' or 'Gum base (including CAS No 9011-16-9)'
Chia oil from <i>Salvia hispanica</i>	Specified food category Fats and oils Pure chia oil Food Supplements as defined in Directive 2002/46/EC	Maximum levels 10 % 2 g/day 2 g/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chia oil (<i>Salvia</i> <i>hispanica</i>)'
Chia seeds (Salvia hispanica)	Specified food categoryBread productsBaked productsBaked productsBreakfast cerealsFruit, nut and seed mixesFruit juice and fruit/vegetable blend beverages	Maximum levels 5 % (whole or ground chia seeds) 10 % whole chia seeds 10 % whole chia seeds 10 % whole chia seeds 15 g/day for addition of whole, mashed or ground chia seeds	 The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chia seeds (<i>Salvia hispanica</i>)' Pre-packaged
	Pre-packaged Chia seed as such Fruit spreads Yoghurt	 15 g/day whole chia seeds 1 % whole chia seeds 1,3 g whole chia seeds per 100 g of yoghurt or 4,3 g whole chia seeds per 330 g of yoghurt (portion) 	Chia (Salvia hispanica) seeds shall carry additional labelling to inform the consumer that the daily intake is no

	Sterilised ready to eat meals based on cereal grains, pseudocereals grains and/or pulses	5 % whole chia seeds	more than 15 g.	
Chitin- glucan from <i>Aspergillus</i> <i>niger</i>	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels 5 g/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chitin- glucan from <i>Aspergillus</i> <i>niger</i> '	
Chitin-glucan complex from <i>Fomes</i> <i>fomentarius</i>	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels 5 g/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chitin-glucan from <i>Fomes</i> <i>fomentarius</i> '	
Chitosan extract from fungi (<i>Agaricus</i> <i>bisporus</i> ; <i>Aspergillus</i> <i>niger</i>)	Specified food category Food Supplements as defined in Directive 2002/46/EC	<i>Maximum</i> <i>levels</i> In line with normal use in food supplements of chitosan from crustaceans	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chitosan extract from <i>Agaricus</i> <i>bisporus</i> ' or 'Chitosan extract from <i>Aspergillus</i> <i>niger</i> '	
Chondroitin sulphate	Specified food category Food supplements as defined in Directive 2002/46/EC for adult population, excluding pregnant and lactating women	<i>Maximum levels</i> 1 200 mg/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Chondroitin sulphate derived from microbial fermentation and sulphation'	

Chromium Picolinate	Specified food category	Maximum levels of total chromium	The designation of the novel food on the labelling
	Foods covered by Regulation (EU) No 609/2013	250 μg/day	of the foodstuffs containing it shall be 'Chromium
	Foods fortified in accordance with Regulation (EC) No 1925/2006 ^d		Picolinate'
<i>Cistus incanus</i> L. Pandalis	Specified food category	Maximum levels	The designation of the novel food
herb	Herbal infusions	Intended daily intake: 3 g herbs/ day (2 cups/day)	on the labelling of the foodstuffs containing it shall be ' <i>Cistus</i> <i>incanus</i> L. Pandalis herb'
Citicoline	Specified food category	Maximum levels	1. The designation
	Food Supplements as defined in Directive 2002/46/EC	500 mg/day	of the novel food on the labelling
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	250 mg per serving and a maximum daily consumption level of 1 000 mg	of the foodstuffs containing it shall be 'Citicoline' 2. The labelling of foods containing citicoline shall bear a statement that the product is not intended to be consumed by children

Clostridium butyricum	Specified food category	Maximum levels	The designation of the novel food	
-	Food Supplements as defined in Directive 2002/46/EC	1,35 × 10 ⁸ CFU/ day	on the labelling of the foodstuffs containing it shall be ' <i>Clostridium</i> <i>butyricum</i> MIYAIRI 588 (CBM 588)' or ' <i>Clostridium</i> <i>butyricum</i> (CBM 588)'	
Extract of defatted cocoa	Specified food category	Maximum levels	Consumers shall be instructed not	
powder	Nutrition bars Milk based beverages	1 g/day and 300 mg polyphenols corresponding	to consume more than 600 mg polyphenols corresponding to	
	Any other foods (including food supplements as defined in Directive 2002/46/ EC) which have become established vehicles for functional ingredients and which are typically positioned for consumption by health conscious adults	to not more than 550 mg of extract of defatted cocoa powder in one portion of food (or food supplement)	1,1 g of extract of defatted cocoa powder per day	
Low fat cocoa extract	Specified food category	Maximum levels	Consumers shall be instructed not	
	Foods including food supplements as defined in Directive 2002/46/EC	730 mg per serving and around 1,2 g/day	to consume more than 600 mg of cocoa flavanols per day	
Coriander seed oil from	Specified food category	Maximum levels	The designation of the novel food	
Coriandrum sativum	Food Supplements as defined	600 mg/day	on the labelling of the foodstuffs containing it shall be	

	in Directive 2002/46/EC		'Coriander seed oil'	
Crataegus pinnatifida	Specified food category	Maximum levels	The designation of the novel food on the labelling	
dried fruit	Herbal infusions	In line with		
	Jams and jellies in accordance with Directive 2001/113/EC ^e	normal food use of <i>Crataegus</i> <i>laevigata</i>	of the foodstuffs containing it shall be ' <i>Crataegus</i> <i>pinnatifida</i> dried	
	Compotes		fruit'	
a-cyclodextrin	Not specified	, 	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Alpha- cyclodextrin' or ' α -cyclodextrin'	
γ-cyclodextrin	Not specified		The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Gamma- Cyclodextrin' or ' γ -Cyclodextrin'	
Dextran preparation	Specified food category	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Dextran'	
produced by Leuconostoc mesenteroides	Bakery products	5 %		
Diacylglycerol oil of plant	Specified food category	Maximum levels	The designation of the novel food	
origin	Cooking oils		on the labelling of the foodstuffs	
	Fat spreads		containing	
	Salad dressings		it shall be 'Diacylglycerol	
	Mayonnaise		oil of plant	
	Meal replacement for weight control (as drinks)		origin (at least 80 % diacylglycerols)'	
	Bakery products			

	Yoghurt type products			
Dihydrocapsiate (DHC)	Specified food category	Maximum levels	1.	The designation
(2110)	Cereal bars	9 mg/100 g		of the
	Biscuits, cookies and crackers	9 mg/100 g		novel food on the
	Rice based snacks	12 mg/100 g	of the	labelling of the foodstuffs
	Carbonated drinks, dilutable drinks, fruit juice based beverages	1,5 mg/100 ml	_	containing it shall be 'Dihydrocapsiate'
	Vegetable drinks	2 mg/100 ml	2.	Food supplements
	Coffee based drinks, tea based drinks	1,5 mg/100 ml	_	containing synthetic dihydrocapsiate
	Flavoured water — still	1 mg/100 ml	will be labelled	
	Precooked oatmeal cereal	2,5 mg/100 g		intended for
	Other cereals	4,5 mg/100 g		children up to
	Ice cream, dairy desserts	4 mg/100 g		4.5 years'
	Pudding mixes (ready to eat)	2 mg/100 g		
	Products based on yoghurt	2 mg/100 g		
	Chocolate confectionery	7,5 mg/100 g		
	Hard candy	27 mg/100 g		
	Sugar-free gum	115 mg/100 g		
	Whitener/ creamer	40 mg/100 g		
	Sweeteners	200 mg/100 g		
	Soup (ready to eat)	1,1 mg/100 g		
	Salad dressing	16 mg/100 g		
	Vegetable protein	5 mg/100 g		

	Ready to eat meals Meal replacements for weight control Meal replacement for weight control (as drinks)	3 mg/meal 3 mg/meal 1 mg/100 ml		
	Food Supplements as defined in Directive 2002/46/EC	3 mg/single intake 9 mg/day		
	Non-alcoholic powdered drink mixes	14,5 mg/kg equivalent to 1,5 mg/100 ml		
Dried extract of <i>Lippia citriodora</i>	Specified food category	Maximum levels	The designation of the novel food	
from cell cultures	Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of a similar extract from the leaves of <i>Lippia</i> <i>citriodora</i>	on the labelling of the foodstuffs containing it shall be 'dried extract of <i>Lippia</i> <i>citriodora</i> from cell cultures HTN [®] Vb'	
Echinacea angustifolia	Specified food category	Maximum levels		
extract from cell cultures	Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of a similar extract from the root of <i>Echinacea</i> <i>angustifolia</i>		
Echinacea purpurea	Specified food category	Maximum levels	The designation of the novel food	
extract from cell cultures	Food Supplements as defined in Directive 2002/46/EC	In line with normal use in food supplements of a similar extract from florets within the flower head	on the labelling of the foodstuffs containing it shall be 'dried extract of <i>Echinacea</i> <i>purpurea</i> from cell cultures HTN [®] Vb'	

		of Echinacea purpurea		
<i>Echium plantagineum</i> oil	Specified food category	Maximum levels of stearidonic acid (STA)	The designation of the novel food on the labelling of the foodstuffs	
	Milk-based products and drinkable yoghurt products delivered in a single dose	250 mg/100 g; 75 mg/100 g for drinks	containing it shall be 'Refined echium oil'	
	Cheese preparations	750 mg/100 g		
	Spreadable fat and dressings	750 mg/100 g		
	Breakfast cereals	625 mg/100 g		
	Food supplements as defined in Directive 2002/46/EC	500 mg/day		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal		
Epigallocatechin gallate as a	Specified food category	Maximum levels	The labelling shall bear a	
purified extract from green tea leaves (<i>Camellia</i> <i>sinensis</i>)	Foods including food supplements as defined in Directive 2002/46/EC	150 mg of extract in one portion of food or food supplement	statement that consumers should not consume more than 300 mg of extract per day	

L-ergothioneine	Specified food category	Maximum levels	The designation of the novel food	
	Food supplements as defined in Directive 2002/46/EC	30 mg/day for general population (excluding pregnant and lactating women) 20 mg/day for children older than 3 years	on the labelling of the foodstuffs containing it shall be 'L- ergothioneine'	
Ferric Sodium EDTA	Specified food category	Maximum levels (expressed as anhydrous EDTA)	The designation of the novel food on the labelling of the foodstuffs containing it	
	Food supplements as defined in Directive 2002/46/EC	18 mg/day for children 75 mg/day for adults	shall be 'Ferric Sodium EDTA'	
	Foods covered by Regulation (EU) No 609/2013	12 mg/100 g		
	Foods fortified in accordance with Regulation (EC) No 1925/2006			
Ferrous ammonium	Specified food category	Maximum levels	The designation of the novel food	
phosphate	Food supplements as defined in Directive 2002/46/EC	To be used in compliance with Directive 2002/46/EC, Regulation	on the labelling of the foodstuffs containing it shall be 'Ferrous ammonium phosphate'	
	Foods covered by Regulation (EU) No 609/2013	(EU) No 609/2013 and/or Regulation (EC) No 1925/2006		
	Foods fortified in accordance with Regulation (EC) No 1925/2006			

Fish peptides from <i>Sardinops</i> <i>sagax</i>	Specified food category	Maximum levels fish peptide product	The designation of the novel food on the labelling of the foodstuffs	
	Foods based on yoghurt, yoghurt drinks, fermented milk products, and powdered milk	0,48 g/100 g (ready to eat/ drink)	containing it shall be 'Fish (<i>Sardinops</i> <i>sagax</i>) peptides'	
	Flavoured water, and vegetable- based drinks	0,3 g/100 g (ready to drink)		
	Breakfast cereals	2 g/100 g		
	Soups, stews and soup powders	0,3 g/100 g (ready to eat)		
Flavonoids from Glycyrrhiza glabra	Specified food category	Maximum levels of flavonoids from Glycyrrhiza glabra	1. The designat of the novel food on the	Beverages containing flavonoids shall be presented to the final consumer as
	Beverages based on milk	120 mg/day	labelling of the	single portions.
	Beverages based on yoghurt		foodstuf containin it shall	
	Beverages based on fruit or vegetables		be 'Flavono from	
	Food Supplements as defined in Directive 2002/46/EC	120 mg/day	Glycyrrh glabra L.' 2. The labelling of the	
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	120 mg/day	foods where the product was added	
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	120 mg/day	as a novel food ingredier shall bear a statemen that:	

		(a)	the
			product
			should
			not
			be
			consumed
			by
			pregnant
			and
			breast
			feeding
			women,
			children
			and
			young
			adolescents;
			and
		(b)	people
		(0)	taking
			prescription
			drugs
			should
			only
			consume the
			product
			under
			medical
		()	supervision;
		(c)	a .
			maximum
			of
			120 mg
			of
			flavonoids
			per
			day
			should
			be
			consumed.
	3.	The	
		amount	
		of	
		flavonoi	ds
		in the	
		final	
		food	
		shall be	
		indicated	ł
		on the	
		labelling	
		of the	
		food	
I			

Fucoidan extract from the seaweed <i>Fucus</i> <i>vesiculosus</i>	Specified food category Foods including food supplements as defined in Directive 2002/46/EC for the general population	Maximum levels 250 mg/day	containing it. The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Fucoidan extract from seaweed <i>Fucus</i> <i>vesiculosus</i> '.
Fucoidan extract from the seaweed <i>Undaria</i> <i>pinnatifida</i>	Specified food categoryFoods including food supplements as defined in Directive 2002/46/EC for the general population	Maximum levels 250 mg/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Fucoidan extract from seaweed Undaria pinnatifida'
2'- Fucosyllactose	Specified food category Unflavoured pasteurised and sterilised (including UHT) milk-based products Unflavoured fermented milk- based products	Maximum levels 1,2 g/l 1,2 g/l beverages 19,2 g/kg products other than beverages	1. The designation of the novel food on the labelling of the foodstuffs containing it shall be '2'- fucosyllactose'.
	Flavoured fermented milk- based products including heat- treated products Dairy analogues, including beverage whiteners	 1,2 g/l beverages 19,2 g/kg products other than beverages 1,2 g/l beverages 12 g/kg for products other than beverages 400 g/kg for whitener 	2. The labelling of food supplements containing 2'- fucosyllactose shall bear a statement that the supplements should not be
	Cereal bars	12 g/kg	used if other

Table-top sweeteners	200 g/kg	foods with added
Infant formula as defined in Regulation (EU) No 609/2013	1,2 g/l alone or in combination with up to 0,6 g/ l of lacto- <i>N</i> - neotetraose at a ratio of 2:1 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	2'- fucosyllactose are consumed the same day. 3. The labelling of food supplements containing
Follow-on formula as defined in Regulation (EU) No 609/2013	1,2 g/l alone or in combination with up to 0,6 g/ l of lacto- <i>N</i> - neotetraose at a ratio of 2:1 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	2'- fucosyllactose intended for young children shall bear a statement that the supplements should not be
Processed cereal-based food and baby	12 g/kg for products other than beverages	used if breast milk or
food for infants and young children as defined in Regulation (EU) No 609/2013	1,2 g/l for liquid food ready for use, marketed as such or reconstituted as instructed by the manufacturer	other foods with added 2'- fucosyllactose are consumed
Milk-based drinks and similar products intended for young children	1,2 g/l for milk- based drinks and similar products added alone or in combination with up to 0,6 g/l lacto- <i>N</i> - neotetraose, at a ratio of 2:1 in the final product ready for use, marketed as such or reconstituted	the same day.

Foods for special medical purposes as defined in Regulation (EU) No 609/2013	as instructed by the manufacturer In accordance with the particular nutritional requirements of the persons for whom the products are intended
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	4,8 g/l for drinks 40 g/kg for bars
Bread and pasta products bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014	60 g/kg
Flavoured drinks	1,2 g/l
Coffee, tea (excluding black tea), herbal and fruit infusions, chicory; tea, herbal and fruit infusions and chicory extracts; tea, plant, fruit and cereal preparations for infusions, as well as mixes and instant mixes of these products	9,6 g/l — the maximum level refers to the products ready to use
Food supplements as defined in Directive	3,0 g/day for general population

	2002/46/EC, excluding food supplements for infants	1,2 g/day for young children	
Galacto- oligosaccharide	Specified food category	Maximum levels (expressed as ratio kg galacto- oligosaccharide/ kg final food)	
	Food Supplements as defined in Directive 2002/46/EC	0,333	
	Milk	0,02	
	Milk drinks	0,03	
	Meal replacement for weight control (as drinks)	0,02	
	Dairy analogue drinks	0,02	
	Yoghurt	0,033	
	Dairy based deserts	0,043	
	Frozen dairy deserts	0,043	
	Fruit drinks and energy drinks	0,021	
	Infant meal replacement drinks	0,012	
	Baby juice	0,025	
	Baby yogurt drink	0,024	
	Baby desert	0,027	
	Baby snack	0,143	
	Baby cereals	0,027	
	Drinks intended to meet the expenditure of	0,013	

	intense muscular effort especially for sportsmen		
	Juice	0,021	
	Fruit pie fillings	0,059	
	Fruit preparations	0,125	
	Bars	0,125	
	Cereals	0,125	
	Infant formula and follow- on formula as defined in Regulation (EU) No 609/2013	0,008	
Glucosamine HCl	Specified food category	Maximum levels	
	Food Supplements as defined in Directive 2002/46/EC	In line with normal food use of glucosamine from shell fish	
	Foods covered by Regulation (EU) No 609/2013		
	Meal replacement for weight control		
	Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen		
	Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements		

Glucosamine sulphate KCl	Regulation (EU) No 828/2014 Specified food category Food Supplements as defined in Directive 2002/46/EC	<i>Maximum</i> <i>levels</i> In line with normal food use of glucosamine from shell fish			
Glucosamine sulphate NaCl	Specified food category	Maximum levels			
	Food Supplements as defined in Directive 2002/46/EC	In line with normal food use of glucosamine from shell fish			
Guar Gum	Specified food category	Maximum levels	1.	The designation	on
	Fresh dairy products such as yogurts, fermented milks, fresh cheeses and other dairy- based desserts.	1,5 g/100 g		of the novel food on the labelling of the foodstuffs containing it shall be 'Guar Guar Gum'.	s
	Fruit or vegetable- based liquid foodstuffs (of the 'smoothie' variety)	1,8 g/100 g	2.		
	Fruit or vegetable-based compotes	3,25 g/100 g	mention of the possible risks of digestive discomfo linked to the	of the possible	
	Cereals accompanied by a dairy product, in packaging containing two compartments	10 g/100 g in the cereals None in the accompanying dairy product 1 g/100 g in the product when ready to eat		digestive discomfort linked to the exposure of children aged under 8 to guar gum must be visible on the	rt

3.	of any foodstuffs containing it. For example, 'Excessive consumption of these products may cause digestive discomfort, especially for children under 8 years of age'. In the case of products with two compartments containing dairy and cereal products respectively, the instructions for use
2	
5.	
	two
	-
	the
	must clearly
	specify
	the
	need to
	mix the
	cereal and the
	dairy
	product
	before
	consumption,
	in order
	to take
	to take into
	to take into account
	to take into

Heat-treated milk products fermented with <i>Bacteroides</i> <i>xylanisolvens</i>	Specified food category Fermented milk products (in liquid, semi- liquid and spray- dried powder forms)	Maximum levels	risk of gastro- intestinal obstruction.
Hydroxytyrosol	Specified food category Fish and vegetable oils, (except olive oils and olive pomace oils as defined in Part VIII of Annex VII of Regulation (EU) No 1308/2013 ^f), placed as such on the market Spreadable fats as defined in Part VII of Annex VII of Regulation (EU) No 1308/2013, placed as such on the market	Maximum levels 0,215 g/kg 0,175 g/kg	The designation of the novel food on the labelling of the food products containing it shall be 'hydroxytyrosol'. The labelling of the food products containing hydroxytyrosol shall bear the following statements: (a) This food product should not be consumed by children under the age of three years, pregnant women, and lactating women; (b) This food product should not be used for

Ice Structuring Protein type III HPLC 12	<i>Specified food</i> <i>category</i> Edible ices	<i>Maximum</i> <i>levels</i> 0,01 %	cooking, baking or frying' The designation of the novel food on the labelling of the foodstuffs
			containing it shall be 'Ice Structuring Protein'
Aqueous extracts of	Specified food category	Maximum levels	The designation of the novel food
dried leaves of	Herbal infusions	In line with	on the labelling
Ilex guayusa	Food Supplements as defined in Directive 2002/46/EC	normal use in herbal infusions and food supplements of a similar aqueous extract of dried leaves of <i>Ilex</i> <i>paraguariensis</i>	of the foodstuffs containing it shall be 'Extracts of dried leaves of <i>Ilex guayusa</i> '
Isomalto- oligosaccharide	Specified food category	Maximum levels	1. The designation
	Energy-Reduced Soft Drinks	6,5 %	of the novel food
	Energy Drinks	5,0 %	on the
	Foods intended to meet the expenditure of intense muscular efforts, especially for sportsmen (including isotonic drinks)	6,5 %	labelling of the foodstuffs containing it shall be 'Isomaltooligosaccharide'. 2. Foods containing the
	Fruit Juices	5 %	novel
	Processed Vegetables and Vegetable Juices	5 %	ingredient must be labelled as 'a
	Other Soft Drinks	5 %	source of
	Cereals Bars	10 %	glucose'.
	Cookies, Biscuits	20 %	

			_	
	Breakfast Cereal Bars	25 %		
	Hard Candies	97 %		
	Soft Candies/ Chocolate Bars	25 %		
	Meal replacement for weight control (as bars or milk based)	20 %	-	
Isomaltulose	Not specified		 The designation of the novel food on the labelling of the foodstuff containing it shall be 'Isomalt The designation of the novel food on the labelling shall be accompany by indication that the 'Isomalt is a source of glucose and food food food food food food food fo	fs ng ulose'. ion anied an ulose
Lactitol	Specified food	Maximum	fructose The designation	•
	category	levels	of the novel food	
	Food	20 g/day	on the labelling of the food	
	Supplements as defined		supplements	
	in Directive		containing	
	2002/46/EC		it shall be	
	(capsules or		'Lactitol'	

Lacto-N- neotetraose	tablets) intended for the adult population Specified food <i>category</i> Unflavoured pasteurised and sterilised (including UHT) milk-based products	Maximum levels 0,6 g/l	1.	The designation of the novel food on the labelling of the foodstuffs
	Unflavoured fermented milk- based products	0,6 g/l for beverages 9,6 g/kg for products other than beverages		containing it shall be 'lacto-N- neotetraose'.
	Flavoured 0,6 g/l for fermented milk- based products 9,6 g/kg for including heat-	beverages 9,6 g/kg for	2.	The labelling of food supplements containing
	Dairy analogues, including beverage whiteners	0,6 g/l for beverages 6 g/kg for products other than beverages 200 g/kg for whitener		lacto- <i>N</i> - neotetraose shall bear a statement that the supplements should
	Cereal bars	6 g/kg		not be
	Table-top sweeteners	100 g/kg		used if other foods
	Infant formula as defined in Regulation (EU) No 609/2013	0,6 g/l in combination with up to 1,2 g/l of 2'- fucosyllactose at a ratio of 1:2 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	added lacto- <i>N</i> - neotetra are consum the same day. 3. The labelling of food supplen containi lacto- <i>N</i> - neotetra	lacto-N- neotetraose are consumed the same day. The labelling
	Follow-on formula as defined in Regulation (EU) No 609/2013	0,6 g/l in combination with up to 1,2 g/l of 2'- fucosyllactose at		containing lacto-N- neotetraose intended

Processed cereal-based food and baby food for infants and young children as defined in Regulation (EU) No 609/2013 Milk-based drinks and similar products intended for young children	a ratio of 1:2 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer 6 g/kg for products other than beverages 0,6 g/l for liquid food ready for use, marketed as such or reconstituted as instructed by the manufacturer 0,6 g/l for milk- based drinks and similar products added alone or in combination with 2'- fucosyllactose, at a ratio of 1:2 in the final product ready for use, marketed as such or reconstituted as instructed by the manufacturer	young children shall bear a statement that the supplements should not be used if breast milk or other foods with added lacto- <i>N</i> - neotetraose are consumed the same day.
special medical purposes as defined in Regulation (EU) No 609/2013	with the particular nutritional requirements of the persons for whom the products are intended	
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	2,4 g/l for drinks 20 g/kg for bars	
Bread and pasta products bearing statements on the absence or	30 g/kg	

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014			
	Flavoured drinks	0,6 g/l		
	Coffee, tea (excluding black tea), herbal and fruit infusions, chicory; tea, herbal and fruit infusions and chicory extracts; tea, plant, fruit and cereal preparations for infusions, as well as mixes and instant mixes of these products	4,8 g/l — the maximum level refers to the products ready to use		
	Food supplements as defined in Directive 2002/46/EC, excluding food supplements for infants	1,5 g/day for general population 0,6 g/day for young children		
Lucerne leaf extract from	Specified food category	Maximum levels	The designation of the novel food	
Medicago sativa	Food supplements as defined in Directive 2002/46/EC	10 g/day	on the labelling of the foodstuffs containing it shall be 'Lucerne (<i>Medicago</i> <i>sativa</i>) protein' or 'Alfalfa (<i>Medicago</i> <i>sativa</i>) protein'.	
Lycopene	Specified food category	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing	
	Fruit/vegetable juice-based drinks (including concentrates)	2,5 mg/100 g		

	Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen	2,5 mg/100 g	it shall be 'Lycopene'	
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	8 mg/meal		
	Breakfast cereals Fats and	5 mg/100 g 10 mg/100 g	-	
	dressings Soups other than tomato soups	1 mg/100 g	-	
	Bread (including crispy breads)	3 mg/100 g	-	
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Food supplements as defined in Directive 2002/46/EC	15 mg/day		
Lycopene from <i>Blakeslea</i> trispora	Specified food category	Maximum levels	The designation of the novel food	
	Fruit/vegetable juice-based drinks (including concentrates)	2,5 mg/100 g	on the labelling of the foodstuffs containing it shall be	
	Drinks intended to meet the expenditure of intense muscular	2,5 mg/100 g	- 'Lycopene'	

	effort especially for sportsmen Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	8 mg/meal		
	Breakfast cereals	5 mg/100 g		
	Fats and dressings	10 mg/100 g		
	Soups other than tomato soups	1 mg/100 g		
	Bread (including crispy breads)	3 mg/100 g		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Food supplements as defined in Directive 2002/46/EC	15 mg/day		
Lycopene from tomatoes	Specified food category	Maximum levels	The designation of the novel food	
	Fruit/vegetable juice-based drinks (including concentrates)	2,5 mg/100 g	on the labelling of the foodstuffs containing it shall be	
	Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen	2,5 mg/100 g	'Lycopene'	
	Total diet replacement for weight control	8 mg/meal		

	as defined in Regulation (EU) No 609/2013 and meal replacements for weight control Breakfast cereals Fats and dressings Soups other than tomato soups Bread (including	5 mg/100 g 10 mg/100 g 1 mg/100 g 3 mg/100 g		
	crispy breads) Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended		
	Food supplements as defined in Directive 2002/46/EC	15 mg/day		
Lycopene oleoresin from tomatoes	Specified food category Fruit/vegetable juice-based drinks (including concentrates)	Maximum levels of lycopene 2,5 mg/100 g	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Lycopene	
	Drinks intended to meet the expenditure of intense muscular effort especially for sportsmen	2,5 mg/100 g	oleoresin from tomatoes'	
	Total diet replacement for weight control covered by Regulation (EU) No 609/2013 and meal	8 mg/meal		

	replacements for weight control			
	Breakfast cereals	5 mg/100 g]	
	Fats and dressings	10 mg/100 g		
	Soups other than tomato soups	1 mg/100 g		
	Bread (including crispy breads)	3 mg/100 g		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended	-	
Magnesium citrate malate	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Magnesium citrate malate'	
Magnolia Bark Extract	Specified food category	Maximum levels	The designation of the novel food	
	Mints (confectionary products) Chewing gum	0,2 % for breath freshening purposes. Based on a 0,2 % maximum incorporation level and a maximum gum/ mint size of 1,5 g each, each gum or mint serving will contain no more than 3 mg of magnolia bark extract.	on the labelling of the foodstuffs containing it shall be 'Magnolia Bark Extract'	
Maize-germ oil high in	Specified food category	Maximum levels	The designation of the novel food	
unsaponifiable matter	Food Supplements	2 g/day	on the labelling of the foodstuffs	

Methylcellulose	as defined in Directive 2002/46/EC Chewing gum <i>Specified food</i> <i>category</i> Edible ices Flavoured drinks	2 % Maximum levels 2 %	containing it shall be 'Maize- germ oil extract'The designation of the novel food on the labelling of the foodstuffs	
	Flavoured or unflavoured fermented milk products		containing it shall be 'Methylcellulose'	prepared for young children
	Cold desserts (dairy, fat, fruit, cereal, egg-based products)			
	Fruit preparations (pulps, purees or compotes)			
	Soups and broths			
(6S)-5- methyltetrahydro acid, glucosamine salt		Maximum levels	The designation of the novel food on the labelling of the foodstuffs containing it shall be '(6S)-5- methyltetrahydrof acid, glucosamine salt' or '5MTHF- glucosamine'	olic
	Food Supplements as defined in Directive 2002/46/EC as a source of folate			
Monomethylsilan (Organic Silicon)Amerified food categoryMaximum levels of siliconFood Supplements as defined in Directive 2002/46/EC for10,40 mg/day	e Sabr ified food category	<i>levels of</i> of the nor	The designation of the novel food on the labelling	
	10,40 mg/day	of the food supplements containing it shall be		

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	adult population (in liquid form)		'Organic silicon (monomethylsilanetriol)'
Mycelial extract from Shiitake	Specified food category	Maximum levels	The designation of the novel food
mushroom	Bread products	2 ml/100 g	on the labelling
(Lentinula edodes)	Soft drinks	0,5 ml/100 ml	of the foodstuffs containing
	Ready prepared meals	2,5 ml per meal	it shall be 'extract from the mushroom
	Foods based on yoghurt	1,5 ml/100 ml	<i>Lentinula</i> <i>edodes</i> ' or
	Food supplements as defined in Directive 2002/46/EC	2,5 ml per day dose	'extract from Shiitake mushroom'
Noni fruit juice (<i>Morinda</i>	Specified food category	Maximum levels	The designation of the novel food
citrifolia)	<i>folia</i>) Pasteurised fruit and fruit nectar based drinks 100 % noni juice) of the hover it on the labellin of the foodstu- containing it	shall be 'Noni juice' or 'Juice of <i>Morinda</i>	
Noni fruit juice powder (<i>Morinda</i> <i>citrifolia</i>)	Food supplements as defined in Directive 2002/46/EC	6,6 g/day (equivalent to 30 ml of noni juice)	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Noni juice powder' or 'Juice powder of <i>Morinda</i> <i>citrifolia</i> '
Noni fruit puree and concentrate (<i>Morinda</i> <i>citrifolia</i>)	Specified food category	Maximum levels	The designation of the novel food
		Fruit puree	on the labelling
	Candy/ confectionery	45 g/100 g	- of the foodstuffs containing it shall be:
	Cereal bars	53 g/100 g	For fruit puree:
	Powdered nutritional drink mixes (dry weight)	53 g/100 g	<i>Citrifolia</i> fruit puree' or 'Noni fruit puree'

Carbonated beverages	11 g/100 g	For fruit concentrate: <i>Morinda</i>
Ice cream & sorbet	31 g/100 g	<i>citrifolia</i> fruit concentrate'
Yoghurt	12 g/100 g	or 'Noni fruit concentrate'
Biscuits	53 g/100 g	concentrate
Buns, cakes and pastries	53 g/100 g	
Breakfast cereals (wholegrain)	88 g/100 g	
Jams and jellies in accordance with Directive 2001/113/EC	133 g/100 g Based on pre- processing quantity to produce final 100 g product	
Sweet spreads, fillings and icings	31 g/100 g	-
Savoury sauces, pickles, gravies and condiments	88 g/100 g	-
Food Supplements as defined in Directive 2002/46/EC	26 g/day	
	Fruit concentrate	
Candy/ Confectionery	10 g/100 g	
Cereal bars	12 g/100 g	
Powdered nutritional drink mixes (dry weight)	12 g/100 g	
Carbonated beverages	3 g/100 g	
Ice cream & sorbet	7 g/100 g	
Yoghurt	3 g/100 g]
Biscuits	12 g/100 g]
Buns, cakes and pastries	12 g/100 g	

	Breakfast cereals (wholegrain)	20 g/100 g		
	Jams and jellies in accordance with Directive 2001/113/EC	30 g/100 g		
	Sweet spreads, fillings and icings	7 g/100 g		
	Savoury sauces, pickles, gravies and condiments	20 g/100 g		
	Food Supplements as defined in Directive 2002/46/EC	6 g/day	-	
Noni leaves (<i>Morinda</i>	Specified food category	Maximum levels	1.	The designation
(viorinaa citrifolia)	For the preparation of infusions	A cup of infusion to be consumed shall not be prepared with more than 1 g of dried and roasted leaves of <i>Morinda</i> <i>citrifolia</i>	2.	of the novel food on the labelling of the foodstuffs containing it shall be 'Noni leaves' or 'leaves of <i>Morinda</i> <i>citrifolia</i> '. Instructions shall be given to the consumer that a cup of infusion should not be prepared with more

			1 g of dried and roasted leaves of Morinda citrifolia	
Noni fruit	Specified food category	Maximum levels	The designation	
powder (<i>Morinda</i> <i>citrifolia</i>)	Food Supplements as defined in Directive 2002/46/EC	2,4 g per/day	of the novel food on the labelling of the foodstuffs containing it shall be 'Morinda citrifolia fruit powder' or 'Noni fruit powder'	
<i>Odontella aurita</i> microalgae	Specified food category	Maximum levels	The designation of the novel food	
linerbargae	Flavoured pasta	1,5 %	on the labelling	
	Fish soups	1 %	of the foodstuffs containing	
	Marine terrines	0,5 %	it shall be	
	Broth preparations	1 %	'Odontella aurita microalgae'	
	Crackers	1,5 %		
	Frozen breaded fish	1,5 %	-	
Oil enriched with phytosterols/ phytostanols	Specified food category	Maximum levels of phytosterols/ phytostanols	In accordance with Annex III.5 to Regulation (EU) No	
	Spreadable fats as defined in Annex VII, Part VII and Appendix II, points B and C of Regulation (EU) No 1308/2013, and excluding cooking and frying fats and spreads based on butter or other animal fat	1. The products containing the novel food ingredie shall be presente in such a manner that they can be	ng nt	

OH UAH AUUU	- <u>r</u> <u>j</u> <u>j</u> <u>j</u> <u>j</u> <u>j</u> <u>j</u> <u>j</u> -	levels of	-	ine designation	
Oil extracted	Specified food	Maximu	m	The designation	
			portions		
			single		
			as		
			packed		
			shall be		
			spicy		
			and spicy		
			mayonna	aise	
			dressing	-	
			Salad		
			3 g.		
			exceed		
			not		
			shall		
			beverage	2S	
			of	1	
	spicy sauces		to a containe	r	
	mayonnaise and		added		
	Salad dressings,		phytosta	nols	
	Soya drinks		phytoste		
			of		
	protein		amount		
	replaced by vegetable fat or		The		
	partly or fully		phytoste		
	protein has been		added phytoste	rols/	
	and the fat or		day) of		
	been reduced		per		
	the milk fat has		portions		
	where possibly		of three		
	per 100 g),		(in case		
	content ≤ 12 g		of 1 g		
	products (fat		maximu	m	
	cheese based		or a		
	milk such as yoghurt and		day)		
	on fermented		portion		
	products based		of one portion		
	and/or cereals,		(in case		
	addition of fruits		of 3 g		
	possibly with the		maximu	m	
	products,		either a		
	skimmed milk		contain		
	skimmed and		that		
	based on semi-		portions		
	products, such as products		divided into		
			uivided		

		and EPA combined	on the labelling of the foodstuffs		
	Dairy products except milk- based beverages	200 mg/100 g or for cheese products 600 mg/100 g	containing it shall be 'Squid oil'.	shall be 'Squid	
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g			
	Spreadable fat and dressings	600 mg/100 g	-		
	Breakfast cereals	500 mg/100 g			
	Bakery products (breads and bread rolls)	200 mg/100 g			
	Cereal bars	500 mg/100 g			
	Non-alcoholic beverages (including milk- based beverages)	60 mg/100 ml			
	Food Supplements as defined in Directive 2002/46/EC	3 000 mg/day for general population 450 mg/day for pregnant and lactating women			
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products intended			
	Total diet replacement for weight control defined in Regulation (EU) No 609/2013 and meal replacements for weight control	200 mg/meal			
Pasteurised fruit-based	Specified food category	Maximum levels	The wording 'pasteurised by		

preparations produced using high-pressure treatment	Types of fruit: apple, apricot, banana, blackberry, blueberry, cherry, coconut, fig, grape, grapefruit, mandarin, mango, melon, peach, pear, pineapple, prune, raspberry, rhubarb, strawberry		high-pressure treatment' shall be displayed next to the name of the fruit preparations as such and in any product in which it is used
Phosphated maize starch	Specified food category	Maximum levels	The designation of the novel food
maize starch	Baked bakery products Pasta Breakfast cereals	15 %	on the labelling of the foodstuffs containing it shall be 'Phosphated
	Cereal bars		maize starch'
Phosphatidylseri		Maximum	The designation
from fish	category	levels of	of the novel food
phospholipids		phosphatidylseri	
	Beverages based on yoghurt	50 mg/100 ml	of the foodstuffs containing it shall be 'Fish
	Powders based on milk powders	3 500 mg/100 g (equivalent to 40 mg/100 ml ready to drink)	phosphatidylserine'
	Foods based on yoghurt	80 mg/100 g	
	Cereal bars	350 mg/100 g	
	Chocolate based confectionary	200 mg/100 g	
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In compliance with Regulation (EU) No 609/2013	
	Food supplements as defined	300 mg/day	

	in Directive 2002/46/EC			
Phosphatidylseri from soya phospholipids	nSpecified food category	Maximum levels of phosphatidylseri	The designation of the novel food	
	Beverages based on yoghurt	50 mg/100 ml	of the foodstuffs containing it shall be 'Soya phosphatidylserine'	
	Powders based on milk powder	3,5 g/100 g (equivalent to 40 mg/100 ml ready to drink)		2'
	Foods based on yoghurt	80 mg/100 g		
	Cereal bars	350 mg/100 g		
	Chocolate based confectionary	200 mg/100 g		
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In compliance with Regulation (EU) No 609/2013		
Phospholipid product containing	Specified food category	Maximum levels of phosphatidylseri	The designation of the novel food	The product is not intended to be marketed
product containing equal	1 0 0		of the novel food of the labelling of the foodstuffs	not intended to be marketed to pregnant or
product containing equal amounts of	<i>category</i> Breakfast cereals	levels of phosphatidylseri	of the novel food 46 n the labelling of the foodstuffs containing	not intended to be marketed
product containing equal amounts of phosphatidylseri and phosphatidic	<i>category</i> Breakfast cereals	<i>levels of</i> <i>phosphatidylseri</i> 80 mg/100 g	of the novel food G n the labelling of the foodstuffs containing shall be 'Soy phosphatidylsering and phosphatidic	not intended to be marketed to pregnant or breast-feeding women
product containing equal amounts of phosphatidylseri and	<i>category</i> Breakfast cereals Cereal bars Foods based on	<i>levels of</i> <i>phosphatidylseri</i> 80 mg/100 g 350 mg/100 g	of the novel food of the labelling of the foodstuffs containing shall be 'Soy phosphatidylsering	not intended to be marketed to pregnant or breast-feeding women
product containing equal amounts of phosphatidylseri and phosphatidic	<i>category</i> Breakfast cereals ne ereal bars Foods based on yogurt Soy-based yogurt-like	<i>levels of</i> <i>phosphatidylseri</i> 80 mg/100 g 350 mg/100 g 80 mg/100 g	of the novel food G n the labelling of the foodstuffs containing shall be 'Soy phosphatidylsering and phosphatidic	not intended to be marketed to pregnant or breast-feeding women
product containing equal amounts of phosphatidylseri and phosphatidic	<i>category</i> Breakfast cereals Recereal bars Foods based on yogurt Soy-based yogurt-like products Yogurt based-	<i>levels of</i> <i>phosphatidylseri</i> 80 mg/100 g 350 mg/100 g 80 mg/100 g	of the novel food G n the labelling of the foodstuffs containing shall be 'Soy phosphatidylsering and phosphatidic	not intended to be marketed to pregnant or breast-feeding women
product containing equal amounts of phosphatidylseri and phosphatidic	<i>category</i> Breakfast cereals Cereal bars Foods based on yogurt Soy-based yogurt-like products Yogurt based- drinks Soy-based yogurt-like	<i>levels of</i> <i>phosphatidylseri</i> 80 mg/100 g 350 mg/100 g 80 mg/100 g 50 mg/100 g	of the novel food G n the labelling of the foodstuffs containing shall be 'Soy phosphatidylsering and phosphatidic	not intended to be marketed to pregnant or breast-feeding women

Phospholipides from egg yolk	in Directive 2002/46/EC Foods for special medical purposes as defined in Regulation (EU) No 609/2013 Specified food category	In compliance with Regulation (EU) No 609/2013 Maximum levels		
	Not specified			
Phytoglycogen	Specified food category	Maximum levels	The designation of the novel food	
	Processed foods	25 %	on the labelling of the foodstuffs containing it shall be 'Phytoglycogen'	
Phytosterols/ phytostanols	Specified food category	Maximum levels	In accordance with Annex III.5	
	Rice drinks	1. They shall be presente in such a manner that they can be easily divided into portions that contain either a maximum	of Regulation (EU) No 1169/2011	
	Rye bread with flour containing $\geq 50 \%$ rye (wholemeal rye flour, whole or cracked rye kernels and rye flakes) and $\leq 30 \%$ wheat; and with $\leq 4 \%$ added sugar but no fat added.			
	Salad dressings, mayonnaise and spicy sauces.		m	
	Soya drink	of 3 g		
	Milk type products, such as semi- skimmed and skimmed milk type products, possibly with the addition of fruits and/or cereals, where possibly the milk fat has	(in case of 1 portion/ day) or a maximum of 1 g (in case of 3 portions, day) of added		

	been reduced, or where milk fat and/or protein has been partly or fully replaced by vegetable fat and/or protein. Products based on fermented milk such as yoghurt and cheese type products (fat content < 12 % per 100 g), where possibly the milk fat has been reduced, or where milk fat and/or protein has been partly or fully replaced by vegetable fat and/or protein Spreadable fats as defined in Annex VII, Part VII and Appendix II, points B and C of Regulation (EU) No 1308/2013, and excluding cooking and frying fats and spreads based on butter or other animal fat. Food Supplements as defined in Directive 2002/46/EC	phytoste phytosta The amount of phytosterols/ phytostanols added to a container of beverages shall not exceed 3 g. Salad dressings, mayonnaise and spicy sauces shall be packed as single portions 3 g/day	
Plum kernel oil	Specified food category For frying and as seasoning	Maximum levels In line with normal food use of vegetable oils	

Potato proteins (coagulated) and hydrolysates thereof	Not specified		The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Potato protein'	
Prolyl oligopeptidase (enzyme preparation)	Specified food category Food Supplements as defined in Directive 2002/46/EC for general adult population	Maximum levels120 PPU/day $(2,7 g of enzymepreparation/day)(2 \times 10^6 PPI/day)PPU - ProlylPeptidase Unitsor ProlineProtease UnitsPPI - ProteasePicomoleInternational$	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Prolyl oligopeptidase'	
Protein extract from pig kidneys	Specified food category Food Supplements as defined in Directive 2002/46/EC Food for special medical purposes as defined in Regulation (EU) No 609/2013	Maximum levels 3 capsules/ day; equalizing 12,6 mg pig kidney extract a day Diamine oxidase (DAO) content: 0,9 mg/day (3 capsules with a content of DAO of 0,3 mg/ capsule)		
Rapeseed oil high in unsaponifiable matter	Specified food category Food Supplements as defined in Directive 2002/46/EC	Maximum levels 1,5 g per portion recommended for daily consumption	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Rapeseed oil extract'	
Rapeseed Protein	As a vegetable protein source in foods except in infant formula and follow-on formula		1. The designat of the novel food on the labelling	

		proximity to the list of ingredients.
	2.	be 'Rapeseed protein'. Any foodstuff containing 'rapeseed protein' shall bear a statement that this ingredient may cause allergic reaction to consumers who are allergic to mustard and products thereof. Where relevant, this statement shall appear in close

<i>Status:</i> Point in time view as at 31/01/2020.	
Changes to legislation: There are currently no known outstanding effects for the	
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)	

	(capsule or tablet form)		2.	supplements containing it shall be ' <i>Trans</i> - resveratrol'. The labelling of food supplements containing trans- resveratrol shall bear a statement that people using medicines should only consume the product under medical supervision.
Trans- resveratrol (microbial source)	Specified food category Food supplements as defined in Directive 2002/46/EC	Maximum levels In line with normal use in food supplements of resveratrol extracted from Japanese knotweed (Fallopia japonica)	1.	The designation of the novel food on the labelling of the food supplements containing it shall be ' <i>Trans</i> -resveratrol'. The labelling of food supplements containing trans-resveratrol shall bear a

			statement that people using medicine should only consume the product under medical supervise	es
Rooster comb extract	Specified food category Milk-based drinks	Maximum levels 40 mg/100 g or mg/100 ml	The designation of the novel food on the labelling of the foodstuffs	
	Milk based fermented drinks	80 mg/100 g or mg/100 ml	containing it shall be 'Rooster comb extract' or	
	Yoghurt-type products	65 mg/100 g or mg/100 ml	'Cockerel comb extract'	
	Fromage frais	110 mg/100 g or mg/100 ml		
0 1 1 1 1				
Sacha inchi oil from <i>Plukenetia</i> <i>volubilis</i>	Specified food category As for linseed oil	Maximum levels In line with normal food use of linseed oil	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Sacha inchi oil (Plukenetia volubilis)'	
from <i>Plukenetia</i>	category	<i>levels</i> In line with normal food use	of the novel food on the labelling of the foodstuffs containing it shall be 'Sacha inchi oil (Plukenetia	ion

			 2. There shall be a statement that excessive consumption may lead to gastro-intestinal disturbance. 3. There shall be a statement that the products are not intended for use by children.
<i>Schizochytrium</i> <i>sp</i> . oil rich in DHA and EPA	Specified food category	Maximum levels of DHA and EPA combined:	The designation of the novel food on the labelling of the foodstuffs
	Food Supplements as defined in Directive 2002/46/EC for adult population excluding pregnant and lactating women	3 000 mg/day	containing it shall be 'DHA and EPA-rich oil from the microalgae <i>Schizochytrium</i> sp.'
	Food Supplements as defined in Directive 2002/46/EC for pregnant and lactating women	450 mg/day	
	Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the	

	products are intended	
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal	
Milk-based drinks and similar products intended for young children Processed cereal based food and baby food for infants and young children as defined in Regulation (EU)	200 mg/100 g	
No 609/2013 Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen		
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014		
Bakery Products (Breads, Rolls and Sweet Biscuits)	200 mg/100 g	
Breakfast Cereals	500 mg/100 g	
Cooking Fats	360 mg/100 g	

	Dairy Analogues except drinks Dairy Products except milk- based drinks	600 mg/100 g for cheese; 200 mg/100 g for soy and imitation milk products (excluding drinks) 600 mg/100 g for cheese; 200 mg/100 g for milk products (including milk, fromage frais		
		and yoghurt products; excluding drinks)		
	Non-alcoholic Beverages (including dairy analogue and milk-based drinks)	80 mg/100 g		
	Cereal/Nutrition Bars	500 mg/100 g		
	Spreadable Fats and Dressings	600 mg/100 g		
<i>Schizochytrium</i> sp. (ATCC	Specified food category	Maximum levels of DHA	The designation of the novel food	
PTA-9695) oil	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g	on the labelling of the foodstuffs containing it shall be 'Oil from the	
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g		
	Spreadable fats and dressings	600 mg/100 g		
	Breakfast cereals	500 mg/100 g		
	Food Supplements as defined in Directive	250 mg DHA/ day for general population		
	2002/46/EC	450 mg DHA/ day for pregnant		

	and lactating women	
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal	
Milk-based drinks and similar products intended for young children	200 mg/100 g	
Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen		
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014		
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended	
Bakery products (breads,rolls, and, sweet biscuits)	200 mg/100 g	
Cereal bars	500 mg/100 g	
Cooking fats	360 mg/100 g	

	Non-alcoholic beverages (including dairy analogue and milk-based drinks)80 mg/100 mlInfant formula and follow- on formula as defined in Regulation (EU)In accordance with Regulation (EU) No 609/2013			
	No 609/2013 Processed cereal-based foods and baby foods for infants and young children as defined in Regulation (EU) No 609/2013	200 mg/100 g		
<i>Schizochytrium</i> sp. oil	Specified food category	Maximum levels of DHA	The designation of the novel food	
She ou	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g	on the labelling of the foodstuffs containing it shall be 'Oil from the microalgae to <i>Schizochytrium</i> ts sp.'	
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g		
	Spreadable fats and dressings	600 mg/100 g		
	Breakfast cereals	500 mg/100 g	/ al /	
	Food Supplements as defined	250 mg DHA/ day for general population		
	in Directive 2002/46/EC	450 mg DHA/ day for pregnant and lactating women		
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal	250 mg/meal		

replacements for weight control	
Milk-based drinks and similar products intended for young children	200 mg/100 g
Processed cereal-based foods and baby foods for infants and young children as defined in Regulation (EU) No 609/2013	
Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen	
Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014	
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended
Bakery products (breads, rolls and sweet biscuits)	200 mg/100 g
Cereal bars	500 mg/100 g
Cooking fats	360 mg/100 g

	[ſ	י ר	
	Non-alcoholic beverages (including dairy analogue and milk-based drinks)	80 mg/100 ml		
<i>Schizochytrium</i> sp. (T18) oil	Specified food category	Maximum levels of DHA	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Oil from the microalgae <i>Schizochytrium</i> sp.'	
1 ()	Dairy products except milk- based drinks	200 mg/100 g or for cheese products 600 mg/100 g		
	Dairy analogues except drinks	200 mg/100 g or for analogues to cheese products 600 mg/100 g		
	Spreadable fats and dressings	600 mg/100 g		
	Breakfast cereals	500 mg/100 g		
	Food Supplements as defined	250 mg DHA/ day for general population	ral	
	in Directive 2002/46/EC	450 mg DHA/ day for pregnant and lactating women		
	Total diet replacement for weight control as defined in Regulation (EU) No 609/2013 and meal replacements for weight control	250 mg/meal		
	Milk-based drinks and similar products intended for young children	200 mg/100 g		
	Foods intended to meet the expenditure of intense muscular effort, especially for sportsmen			

Foods bearing statements on the absence or reduced presence of gluten in accordance with the requirements of Commission Implementing Regulation (EU) No 828/2014	
Foods for special medical purposes as defined in Regulation (EU) No 609/2013	In accordance with the particular nutritional requirements of the persons for whom the products are intended
Bakery products (breads, rolls and, sweet biscuits)	200 mg/100 g
Cereal bars	500 mg/100 g
Cooking fats	360 mg/100 g
Non-alcoholic beverages (including dairy analogue and milk-based drinks)	80 mg/100 ml
Infant formula and follow- on formula as defined in Regulation (EU) No 609/2013	In accordance with Regulation (EU) No 609/2013
Processed cereal-based foods and baby foods for infants and young children as defined in Regulation (EU) No 609/2013	200 mg/100 g

Fermented	Specified food	Maximum levels	1. The
soybean extract	<i>category</i> Food Supplements as defined in Directive 2002/46/EC (capsules, tablets or powder form) intended for the adult population, excluding pregnant and lactating women	levels 100 mg/day	designation of the novel food on the labelling of the foodstuffs containing it shall be 'Fermented soybean extract'. 2. The labelling of food supplements containing fermented soybean extract shall bear a statement that persons taking medication should only consume the product under medical supervision.
Spermidine- rich wheat germ extract (<i>Triticum</i> <i>aestivum</i>)	Specified food category Food Supplements as defined in Directive 2002/46/EC intended for the adult population, excluding pregnant and lactating women	<i>Maximum</i> <i>levels</i> Equivalent of max. 6 mg/day spermidine	The designation of the novel food on the labelling of the food supplements containing it shall be 'spermidine- rich wheat germ extract'

Sucromalt	Specified food category	Maximum levels	1. The designation
	Not specified		of the novel food on the labelling of the foodstuffs containing it shall be 'Sucromalt'. 2. The designation of the novel food on the labelling shall be accompanied by indication that the product is a source of glucose and fructose.
Sugar cane fibre	Specified food category	Maximum levels	
	Bread	8 %	
	Bakery goods	5 %	
	Meat and muscle products	3 %	
	Seasonings and spices	3 %	
	Grated cheeses	2 %	
	Special diet foods	5 %	
	Sauces	2 %	
	Beverages	5 %	
Sunflower oil extract	Specified food category	Maximum levels	The designation of the novel food on the labelling

Dried <i>Tetraselmis</i> chuii microalgae	Food Supplements as defined in Directive 2002/46/EC <i>Specified food</i> <i>category</i> Sauces Special salts Condiment Food	1,1 g/day <i>Maximum</i> <i>levels</i> 20 % or 250mg/ day 1 % 250 mg/day 250 mg/day	of the foodstuffs containing it shall be 'Sunflower oil extract' The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Dried microalgae <i>Tetraselmis</i> <i>chuii</i> ' or 'Dried	
	Supplements as defined in Directive 2002/46/EC		microalgae <i>T.</i> <i>chuii</i> ' Food supplements containing dried microalgae <i>Tetraselmis</i> <i>chuii</i> shall bear the following statement: 'Contains negligible amounts of iodine'	
<i>Therapon</i> <i>barcoo</i> /Scortum	Intended use identical to that of the salmon, namely the preparation of culinary fish products and dishes, including cooked, raw, smoked and baked fish products			
D-Tagatose	Specified food category	Maximum levels	1. The designat	ion
	Not specified		of the novel food on the labelling of the foodstuf containin it shall be 'D- Tagatose 2. The labelling of any product	fs ng ?.

			where the level of D- Tagatose exceeds 15 g per serving and all beverage containin greater than 1 % D- Tagatose (as consume shall bear a statemen 'excessiv consump may produce laxative effects'.	es ng ed) tt
Taxifolin-rich extract	Specified food category Food Supplements as defined in Directive 2002/46/EC intended for the general population, excluding infants, young children, children and adolescents younger than 14	Maximum levels 100 mg/day	The designation of the novel food on the labelling of the foodstuffs containing it shall be 'taxifolin-rich extract'.	
Trehalose	years Specified food <i>category</i> Not specified	Maximum levels	1. The designat of the novel food on the labelling of the	

			2.	foodstuffs containing it shall be 'Trehalose' and shall be displayed on the labelling of the product as such or in the list of ingredients of foodstuffs containing it. The designation of the novel food on the labelling shall be accompanied by indication that the 'Trehalose is a source of glucose'.
UV-treated mushrooms (<i>Agaricus</i>	Specified food category	Maximum levels of vitamin D ₂		<u> </u>
bisporus)	Mushrooms (<i>Agaricus</i> <i>bisporus</i>)	10 μg of vitamin D ₂ /100 g fresh weight	1.	The designation on the label of the novel food as such or of the foodstuffs containing

			it shall
			be 'UV- treated mushrooms (<i>Agaricus</i>
			<i>bisporus</i>)'. 2. The designation on the label of the
			novel food as such or of the foodstuffs
			containing it shall be accompanied by indication
			that a 'controlled light treatment was
			used to increase vitamin D levels'
			or 'UV treatment was used to increase vitamin D ₂
UV-treated baker's yeast (<i>Saccharomyces</i>	Specified food category	Maximum levels of vitamin D ₂	Ievels'. The designation of the novel food on the labelling
cerevisiae)	Yeast-leavened breads and rolls	5 μ g of vitamin D ₂ /100 g	of the foodstuffs containing it shall be 'Vitamin
	Yeast-leavened fine bakery wares	$5 \ \mu g \ of vitamin D_2/100 \ g$	D yeast' or 'Vitamin D_2 yeast'
	Food Supplements	5 μg of vitamin D ₂ /day	

UV-treated bread	as defined in Directive 2002/46/EC Specified food <i>category</i> Yeast leavened bread and rolls (without toppings)	Maximum levels of vitamin D ₂ 3 μg vitamin D ₂ /100 g	The designation on the label of the novel food shall be accompanied by 'contains vitamin D produced by UV-treatment'	
UV-treated milk	Specified food category Pasteurised whole milk as defined in Regulation (EU) No 1308/2013 to be consumed as such	Maximum levels of vitamin D ₃ 5-32 μg/kg for general population excluding infants	1. The designat on the label of the novel food shall be 'UV- treated'.	ion
	Pasteurised semi-skimmed milk as defined in Regulation (EU) No 1308/2013 to be consumed as such	1-15 μg/kg for general population excluding infants	2. Where UV- treated milk contains an amount of vitamin D that is consider significa in accordar with Point 2 of Part A of Annex XIII to Regulati (EU) No 1169/20 of the Europea Parliame and of the	ed nt nce on 11

Vitamin K ₂ (menaquinone)	To be used in con Directive 2002/46 (EU) No 609/201 Regulation (EC) 1	5/EC, Regulation 3 and/or	Council, the designat for the labelling shall be accompa by 'contain vitamin D produced by UV- treatmen or 'milk containin vitamin D resulting from UV- treatmen The designation of the novel food on the labelling of the foodstuffs containing it shall be 'Menaquinone'	ion nied s d t'
Wheat bran	Specified food	Maximum	or 'Vitamin K ₂ ' The designation	The 'Wheat Bran
extract	category	levels	of the novel food	Extract' may not
	Beer and substitutes	0,4 g/100 g	on the labelling of the foodstuffs	be introduced onto the market as a food
	Ready to eat cereals	9 g/100 g	- containing it shall be 'Wheat bran extract'	supplement or food supplement
	Dairy products	2,4 g/100 g		ingredient.
	Fruit and vegetable juices	0,6 g/100 g	_	Nor may it be added to infant formula.
	Soft drinks	0,6 g/100 g		
	Meat preparations	2 g/100 g		
Yeast beta- glucans	Specified food category	Maximum levels of pure beta-glucans from yeast	The designation of the novel food on the labelling of the foodstuffs containing it	

	(Saccharomyces cervisiae)	(Saccharomyces
Food supplements as defined in Directive 2002/46/EC, excluding food supplements for infants and young children	1,275 g/day for children older than 12 years and general adult population 0,675 g/day for children younger than 12 years	<i>cerevisiae</i>) beta- glucans'
Total diet replacement for weight control as defined in Regulation (EU) No 609/2013	1,275 g/day	
Food for special medical purposes as defined in Regulation (EU) No 609/2013, excluding food for special medical purposes intended for infants and young children	1,275 g/day	
Beverages based on fruit and/ or vegetable juices including concentrate and dehydrated juices	1,3 g/kg	
Fruit-flavoured drinks	0,8 g/kg	
Cocoa beverages preparation powder	38,3 g/kg (powder)	
Other beverages	0,8 g/kg (ready to drink) 7 g/kg (powder)	
Cereal bars	6 g/kg	
Breakfast cereals	15,3 g/kg	

	Foods covered by Regulation (EU) No 609/2013	3 g/day	on the labelling of the foodstuffs containing it shall be 'Zinc L- pidolate'	
Zinc L-pidolate	Specified food category	Maximum levels	The designation of the novel food	
	Food Supplements as defined in Directive 2002/46/EC	2 mg/day	on the labelling of the foodstuffs containing it shall be 'synthetic zeaxanthin'	
Zeaxanthin	Specified food category	Maximum levels	The designation of the novel food	
	Jam, marmalade and other fruit spreads	11,3 g/kg		
	Protein bars and powders	19,1 g/kg		
	Chocolate and confectionery	4 g/kg		
		6,3 g/kg (powder)		
		1,8 g/kg (condensed)		
	Soups and soup mixes	0,9 g/kg (ready to eat)		
	Dried milk/milk powder	25,5 g/kg		
	Milk product analogues	3,8 g/kg		
	Fermented milk products	3,8 g/kg		
	Milk based beverages	3,8 g/kg		
	Cracker-type biscuits	6,7 g/kg		
	Cookie-type biscuits	6,7 g/kg		
	Wholegrain and high fibre instant hot breakfast cereals	1,5 g/kg		

- a Regulation (EU) No 609/2013 of the European Parliament and of the Council of 12 June 2013 on food intended for infants and young children, food for special medical purposes, and total diet replacement for weight control and repealing Council Directive 92/52/EEC, Commission Directives 96/8/EC, 1999/21/EC, 2006/125/EC and 2006/141/EC, Directive 2009/39/EC of the European Parliament and of the Council and Commission Regulations (EC) No 41/2009 and (EC) No 953/2009 (OJ L 181, 29.6.2013, p. 35).
- **b** Commission Implementing Regulation (EU) No 828/2014 of 30 July 2014 on the requirements for the provision of information to consumers on the absence or reduced presence of gluten in food (OJ L 228, 31.7.2014, p. 5).
- c Directive 2002/46/EC of the European Parliament and of the Council of 10 June 2002 on the approximation of the laws of the Member States relating to food supplements (OJ L 183, 12.7.2002, p. 51).
- **d** Regulation (EC) No 1925/2006 of the European Parliament and of the Council of 20 December 2006 on the addition of vitamins and minerals and of certain other substances to foods (OJ L 404, 30.12.2006, p. 26).
- e Council Directive 2001/113/EC of 20 December 2001 relating to fruit jams, jellies and marmalades and sweetened chestnut purée intended for human consumption (OJ L 10, 12.1.2002, p. 67).
- f Regulation (EU) No 1308/2013 of the European Parliament and of the Council of 17 December 2013 establishing a common organisation of the markets in agricultural products and repealing Council Regulation (EEC) No 922/72, (EEC) No 234/79, (EC) No 1037/2001 and (EC) 1234/2007 (OJ L 347, 20.12.2013, p. 671).

TABLE 2: SPECIFICATIONS

Authorised Novel Food	Specifications
N-Acetyl-D-	Description:
neuraminic acid	<i>N</i> -Acetyl-D-neuraminic acid is a white to off-white crystalline powder
	Definition:
	Chemical name:
	IUPAC names:
	<i>N</i> -Acetyl-D-neuraminic acid (dihydrate)
	5-Acetamido-3,5-dideoxy-D-glycero-D-galacto-non-2-ulopyranosonic
	acid (dihydrate)
	Synonyms:
	Sialic acid (dihydrate)
	Chemical formula:
	$C_{11}H_{19}NO_9$ (acid)
	$C_{11}H_{23}NO_{11}$ ($C_{11}H_{19}NO_9 * 2H_2O$) (dihydrate)
	Molecular mass:
	309,3 Da (acid)
	345,3(309,3+36,0) (dihydrate)
	CAS No.:
	131-48-6 (free acid)
	50795-27-2 (dihydrate)
	Specifications:
	Description: white to off-white crystalline powder
	pH (20 °C, 5 % solution): 1,7 – 2,5
	N-Acetyl-D-neuraminic acid (dihydrate): > 97,0 %
	Water (dihydrate calculates to 10,4 %): $\leq 12,5$ % (w/w)
	Ash, sulphated: $< 0.2 \%$ (w/w)
	Acetic acid (as free acid and/or sodium acetate): $< 0.5 \%$ (w/w)
	Heavy Metals:
	Iron: $< 20,0 \text{ mg/kg}$
	Lead: $< 0,1 \text{ mg/kg}$
	Residual proteins: $< 0.01 \%$ (w/w)
	Residual solvents:
	2-Propanol: $< 0,1 \% (w/w)$
	Acetone: $< 0,1 \% (w/w)$
	Ethyl acetate: $< 0,1 \% (w/w)$
	Microbiological criteria:
	Salmonella: Absence in 25 g
	Aerobic mesophilic total count:< 500 CFU/g
	Enterobacteriaceae: Absence in 10 g
	Cronobacter (Enterobacter) sakazakii: Absence in 10 g
	Listeria monocytogenes: Absence in 25 g
	<i>Bacillus cereus:</i> < 50 CFU/g
	Yeasts: < 10 CFU/g
	Moulds: < 10 CFU/g
a Commission Regul	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in

 a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	Residual endotoxins: < 10 EU/mg CFU: Colony Forming Units; EU: Endotoxin Units.
<i>Adansonia digitata</i> (Baobab) dried fruit pulp	Description/Definition: The Baobab (<i>Adansonia digitata</i>) fruits are harvested from trees. The hard shells are cracked open and the pulp is separated from the seeds and the shell. This is milled, separated into coarse and fine lots (particle size 3 to 600 μ) and then packaged. Typical nutritional components: Moisture (loss on drying) (g/100 g): 4,5-13,7 Protein (g/100 g): 1,8-9,3 Fat (g/100 g): 0-1,6 Total carbohydrate (g/100 g): 76,3-89,5 Total sugars (as glucose): 15,2-36,5 Sodium (mg/100 g): 0,1-25,2 Analytical specifications: Foreign matter: Not more than 0,2 % Moisture (loss on drying) (g/100 g): 4,5-13,7 Ash (g/100 g): 3,8-6,6
<i>Ajuga reptans</i> extract from cell cultures	Description/Definition: Hydroalcoholic extract from <i>Ajuga reptans</i> L. tissue cultures which is substantially equivalent to extracts from flowering aerial parts of <i>Ajuga reptans</i> obtained by traditional cultures.
L-Alanyl-L- Glutamine	Description/Definition: L-Alanyl-L-Glutamine is produced by fermentation with a genetically modified strain of <i>Escherichia coli</i> . During the fermentation process, the ingredient is secreted into the growth medium from which it is subsequently separated and purified to a concentration of > 98 %. Appearance: White crystalline powder Purity: > 98 % Infrared spectroscopy: Conformity with ref. standard Appearance of solution: Colourless and clear Assay (dry basis): 98-102 % Related substances (each): $\leq 0,2$ % Residue on ignition: $\leq 0,1$ % Loss on drying: $\leq 0,5$ % Optical rotation: +9,0 - +11,0° pH (1 %; H ₂ O): 5,0-6,0 Ammonium (NH ₄): $\leq 0,020$ % Sulphate (SO ₄): $\leq 0,020$ % Microbiological criteria: <i>Escherichia coli</i> : Absence/g
Algal oil from the microalgae <i>Ulkenia</i> sp.	Description/Definition: Oil from the micro-algae <i>Ulkenia</i> sp. Acid value: ≤ 0.5 mg KOH/g
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

Document Generated: 20	24-07-08 Status: Point in time view as at 31/01/2020.
	Changes to legislation: There are currently no known outstanding effects for the
Comn	nission Implementing Regulation (EU) 2018/1023. (See end of Document for details)
	Peroxide value (PV): \leq 5,0 meq/kg oil Moisture and volatiles: \leq 0,05 % Unsaponifiables: \leq 4,5 % Trans-fatty acids: \leq 1,0 % DHA content: \geq 32 %
411	
Allanblackia seed oil	Description/Definition: Allanblackia seed oil is obtained from the seeds of the allanblackia species: A. floribunda (synonymous with A. parviflora) and A. stuhlmannii. Composition of fatty acids: Lauric acid (C12:0): < 1,0 % Myristic acid (C14:0): < 1,0 % Palmitic acid (C16:1): < 1,0 % Stearic acid (C16:1): < 1,0 % Stearic acid (C18:1): < 45-58 % Oleic acid (C18:1): 40-51 % Linoleic acid (C18:2): < 1,0 % Y-Linolenic acid (C18:3): < 1,0 % Free fatty acids: max 0,1 % Characteristics: Trans fatty acids: max 0,5 % Peroxide value (PV): max 0,8 meq/kg Iodine value: < 46 g/100 g Unsaponifiable matter: max 1,0 % Saponification value: 185-198 mg KOH/g
<i>Aloe macroclada</i> Baker leaf extract	Description/Definition: Powdered gel extract derived from the leaves of <i>Aloe macroclada</i> Baker which is substantially equivalent to the same gel derived from <i>Aloe vera</i> (L.) Burm.f. leaves. Ash: 25 % Dietary fibres: 28,6 % Fat: 2,7 % Moisture: 4,7 % Polysaccharides: 9,5 % Protein: 1,63 % Glucose: 8,9 %
Antarctic Krill oil from <i>Euphausia</i> superba	Description/Definition: To produce lipid extract from Antarctic Krill (<i>Euphausia superba</i>) deep- frozen crushed krill or dried krill meal is subjected to lipid extraction with an approved extraction solvent (under Directive 2009/32/EC). Proteins and krill material are removed from the lipid extract by filtration. The extraction solvents and residual water are removed by evaporation. Saponification value: ≤ 230 mg KOH/g Peroxide value (PV): ≤ 3 meq O ₂ /kg oil
Annexes II and III p. 1).	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
b Commission Imple	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	Document Generated: 2024-07-0
	Status: Point in time view as at 31/01/2020. Changes to legislation: There are currently no known outstanding effects for the aission Implementing Regulation (EU) 2018/1023. (See end of Document for details)
	Oxidative stability: All food products containing Antarctic Krill oil from <i>Euphausia superba</i> should demonstrate oxidative stability by appropriate and recognised national/international test methodology (e.g. AOAC). Moisture and volatiles: $\leq 3 \%$ or 0,6 expressed as water activity at 25 °C Phospholipids: 35-50 % Trans-fatty acids: $\leq 1 \%$ EPA (eicosapentaenoic acid): $\geq 9 \%$ DHA (docosahexaenoic acid): $\geq 5 \%$
Antarctic Krill oil rich in phospholipids from <i>Euphausia</i> <i>superba</i>	Description/Definition: Oil rich in phospholipids is produced from Antarctic krill (<i>Euphausia superba</i>) by repeated solvent washings with an approved solvent (under Directive 2009/32/EC) to increase phospholipid content of the oil. Solvents are removed from the final product by evaporation. Saponification value: $\leq 230 \text{ mg KOH/g}$ Peroxide value (PV): $\leq 3 \text{ meq O}_2/\text{kg oil}$ Moisture and volatiles: $\leq 3 \%$ or 0,6 expressed as water activity at 25 °C Phospholipids: $\geq 60 \%$ Trans-fatty acids: $\leq 1 \%$ EPA (eicosapentaenoic acid): $\geq 9 \%$ DHA (docosahexaenoic acid): $\geq 5 \%$
Arachidonic acid-rich oil from the fungus <i>Mortierella</i> <i>alpina</i>	Description/Definition: The clear yellow arachidonic acid-rich oil is obtained by fermentation of the non-genetically modified strains IS-4, I49-N18, FJRK-MA01 and CBS 210.32 of the fungus <i>Mortierella alpina</i> using a suitable liquid. The oil is then extracted from the biomass and purified. Arachidonic acid: ≥ 40 % by weight of the total fatty acid content Free fatty acids: $\leq 0,45$ % of the total fatty acid content Trans fatty acids: $\leq 0,5$ % of the total fatty acid content Unsaponifiable matter: $\leq 1,5$ % Peroxide value (PV): ≤ 5 meq/kg Anisidin value: ≤ 20 Acid value: $\leq 1,0$ KOH/g Moisture: $\leq 0,5$ %
Argan oil from <i>Argania spinosa</i>	Description/Definition: Argan oil is the oil obtained by cold pressing of the almond like kernels of the fruits of <i>Argania spinosa</i> (L.) Skeels. Kernels may be roasted prior to pressing, but with no direct contact with a flame. Composition: Palmitic acid (C16:0): 12-15 % Stearic acid (C18:0): 5-7 % Oleic acid (C18:1): 43-50 % Linoleic acid (C18:2): 29-36 % Unsaponifiable matter: 0,3-2 % Total sterols: 100-500 mg/100 g Total tocopherols: 16-90 mg/100 g Oleic acidity: 0,2-1,5 %
	Oleic acidity: 0,2-1,5 % ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Peroxide value (PV): $< 10 \text{ meq O}_{2/2}$

Astaxanthin- rich oleoresin from <i>Haematococcus</i> <i>pluvialis</i> algae	Description/Definition:Astaxanthin is a carotenoid produced by Haematococcus pluvialis algae.Production methods for the growth of the algae are variable; using eitherclosed systems exposed to sunlight or strictly controlled illuminatedlight; alternatively open ponds may be used. The algal cells are harvestedand dried; the oleoresin is extracted using either super critical CO_2 or asolvent (ethyl acetate). The Astaxanthin is diluted and standardized to $2,5 \%, 5,0 \%, 7,0 \%, 10 \%, 15 \%$ or 20 % using olive oil, safflower oil,Sunflower oil or MCT (Medium Chain Triglycerides).Composition of the Oleoresin:Fat: $42,2-99 \%$ Protein: $0,3-4,4 \%$ Carbohydrate: $0-52,8 \%$ Fibre: < 1,0 %
Basil seeds (Ocimum basilicum)	Description/Definition: Basil (<i>Ocimum basilicum</i> L.) belongs to the family ' <i>Lamiaceae</i> ' within the order 'Lamiales'. Post-harvest the seeds are cleaned mechanically. Flowers, leaves and other parts of the plant are removed. Highest level of purity of Basil seeds has to be ensured by filtering (optical, mechanical). Production process of fruit juice and fruit/vegetable blend beverages containing Basil seeds (<i>Ocimum basilicum</i> L.) includes seed pre-hydration and pasteurisation steps. Microbiological controls and monitoring systems are in place. Dry Matter: 94,1 % Protein: 20,7 % Fat: 24,4 % Carbohydrate: 1,7 % Dietary Fibre: 40,5 % (Method: AOAC 958,29)
Annexes II and III p. 1).	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

Fermented black bean extract	Ash: 6,78 % Description/Definition: Fermented black bean extract (Touchi extract) is a fine light-brown protein-rich powder obtained by water extraction of small soybeans (<i>Glycine max (L.) Merr.</i>) fermented with <i>Aspergillus oryzae</i> . The extract contains an α -glucosidase inhibitor. Characteristics: Fat: $\leq 1,0 \%$ Protein: $\geq 55 \%$ Water: $\leq 7,0 \%$ Ash: $\leq 10 \%$ Carbohydrate: $\geq 20 \%$ α -glucosidase inhibitory activity: IC50 min 0,025 mg/ml Soy isoflavone: $\leq 0,3 \text{ g/100 g}$
Bovine lactoferrin	Description/Definition: Bovine lactoferrin is a protein that occurs naturally in cows' milk. It is an iron-binding glycoprotein of approximately 77 kDa and consists of a single polypeptide chain of 689 amino acids. Production process: Bovine lactoferrin is isolated from skimmed milk or cheese whey via ion exchange and subsequent ultra-filtration steps. Finally, it is dried by freeze drying or spraying and the large particles are sieved out. It is a virtually odourless, light pinkish powder. Physical-Chemical properties of Bovine lactoferrin: Moisture: < 4,5 % Ash: < 1,5 % Arsenic: < 2,0 mg/kg Iron: < 350 mg/kg Protein: > 93 % of which bovine lactoferrin: > 95 % of which other proteins: < 5,0 % pH (2 % solution, 20 °C): 5,2-7,2 Solubility (2 % solution, 20 °C): complete
<i>Buglossoides arvensis</i> seed oil	Description/Definition: Refined Buglossoides oil is extracted from the seeds of <i>Buglossoides</i> <i>arvensis</i> (L.) I.M.Johnst Alpha-linolenic acid: ≥ 35 % w/w of total fatty acids Stearidonic acid: ≥ 15 % w/w of total fatty acids Linoleic acid: $\geq 8,0$ % w/w of total fatty acids Trans fatty acids: $\leq 2,0$ % w/w of total fatty acids Acid value: $\leq 0,6$ mg KOH/g Peroxide value (PV): $\leq 5,0$ meq O ₂ /kg Unsaponifiable content: $\leq 2,0$ % Protein content (total nitrogen): ≤ 10 µg/ml Pyrrolizidine alkaloids: Not detectable with a detection limit of 4,0 µg/kg
a Commission Regul Annexes II and III p. 1).	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012)

Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Calanus finmarchicus oil	Description/Definition: The novel food is ruby coloured, slightly viscous oil with a slight shellfish odour extracted from the crustacean (marine zooplankton) <i>Calanus finmarchicus</i> . The ingredient consists primarily of wax esters (> 85 %) with minor amounts of triglycerides and other neutral lipids. Specifications: Water: < 1,0 % Wax esters: > 85 % Total fatty acids: > 46 % Eicosapentaenoic acid (EPA): > 3,0 % Docosahexaenoic acid (DHA): > 4,0 % Total fatty alcohols: > 28 % C20:1 n-9 fatty alcohol: > 9,0 % C22:1 n-11 fatty alcohol: > 12 % Trans fatty acids: < 1,0 % Astaxanthinesters: < 0,1 % Peroxide value (PV): < 3,0 meq. O ₂ /kg
Chewing gum base	Description/Definition: The novel food ingredient is a synthetic polymer (Patent
(monomethoxypo glycol)	hyenhyde We O2006016179). It consists of branched polymers of monomethoxypolyethylene glycol (MPEG) grafted onto polyisoprene-
giycolj	graft-maleic anhydride (PIP-g-MA), and unreacted MPEG (less than
	35 % by weight).
	White to off-white colour. CAS No.: 1246080-53-4
	Characteristics:
	Moisture: < 5,0 %
	Aluminium: < 3,0 mg/kg Lithium: < 0,5 mg/kg
	Nickel: $< 0.5 \text{ mg/kg}$
	Residual anhydride: $< 15 \mu mol/g$
	Polydispersity index: < 1,4
	Isoprene: < 0,05 mg/kg
	Ethylene oxide: < 0,2 mg/kg
	Free maleic anhydride: $< 0,1 \%$ Total oligomeres (less than 1 000 Dalton): $\le 50 \text{ mg/kg}$
	Ethylene glycol: < 200 mg/kg
	Diethylene glycol: < 30 mg/kg
	Monoethylene glycol methyl ether: $< 3,0 \text{ mg/kg}$
	Diethylene glycol methyl ether: < 4,0 mg/kg
	Triethylene glycol methyl ether: $< 7,0 \text{ mg/kg}$
	1,4-Dioxane: < 2,0 mg/kg Formaldehyde: < 10 mg/kg
Chewing	Description/Definition:
gum base (Methyl vinyl	Methyl vinyl ether-maleic anhydride copolymer is an anhydrous copolymer of methyl vinyl ether and maleic anhydride.
a Commission Regul	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

ether-maleic anhydride copolymer)	Free-flowing, white to white-off powder CAS No: 9011-16-9 Purity: Assay value: At least 99,5 % in dry matter Specific viscosity (1 % MEK): 2-10 Residual methyl vinyl ether: ≤ 150 ppm Residual maleic anhydride: ≤ 250 ppm Acetaldehyde: ≤ 500 ppm Methanol: ≤ 500 ppm Dilauroyl peroxide: ≤ 15 ppm Total heavy metals: ≤ 10 ppm Microbiological criteria: Total aerobic plate count: ≤ 500 CFU/g Mould/yeast: ≤ 500 CFU/g <i>Escherichia coli</i> : Negative to test <i>Salmonella</i> : Negative to test <i>Staphylococcus aureus:</i> Negative to test <i>Pseudomonas aeruginosa</i> : Negative to test
Chia oil from <i>Salvia hispanica</i>	Description/Definition: Chia oil is produced from Chia (<i>Salvia hispanica</i> L.) seeds (99,9 % pure) by cold pressing. No solvents are used and, once pressed, the oil is held in decantation tanks and a filtration process employed to remove impurities. It can also be produced by extraction with supercritical CO ₂ . Production process: Produced by cold pressing. No solvents are used and, once pressed, the oil is held in decantation tanks and a filtration process employed to remove impurities. Acidity expressed as oleic acid: $\leq 2,0$ % Peroxide value (PV): ≤ 10 meq/kg Insoluble impurities: $\leq 0,05$ % Alpha linolenic acid: ≥ 60 % Linoleic acid: 15-20 %
Chia seeds (Salvia hispanica)	Description/Definition: Chia (Salvia hispanica L.) is a summer annual herbaceous plant belonging to the Labiatae family. Post-harvest the seeds are cleaned mechanically. Flowers, leaves and other parts of the plant are removed. Dry matter: 90-97 % Protein: 15-26 % Fat: 18-39 % Carbohydrate (*): 18-43 % Crude Fibre(**): 18-43 % Ash: 3-7 % (*) Carbohydrates include the fibre value (**) Crude fibre is the part of fibre made mainly of indigestible cellulose, pentosans and lignin
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
-	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

Status: Point in time view as at 31/01/2020. Changes to legislation: There are currently no known outstanding effects for the

Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Production process: Production process of fruit juices and fruit juice blends beverages, containing Chia seeds, includes seed pre-hydration and pasteurisation steps. Microbiological controls and monitoring systems are in place.
Chitin- glucan from <i>Aspergillus</i> <i>niger</i>	Description/Definition: Chitin-glucan is obtained from the mycelium of Aspergillus niger; it is a slightly yellow, odourless, free-flowing powder. It has a dry matter content of 90 % or more. Chitin-glucan is composed largely of two polysaccharides: — chitin, composed of repeating units of N-acetyl-D-glucosamine (CAS No: 1398-61-4), — beta (1, 3)-glucan, composed of repeating units of D-glucose (CAS No: 9041-22-9). Loss on drying: ≤ 10 % Chitin-glucan: ≥ 90 % Ratio of chitin to glucan: 30:70 to 60:40 Ash: ≤ 3,0 % Lipids: ≤ 1,0 % Proteins: ≤ 6,0 %
Chitin-glucan complex from <i>Fomes</i> <i>fomentarius</i>	 Description/Definition: Chitin-glucan complex is obtained from the cell walls of the fruit bodies of the fungus <i>Fomes fomentarius</i>. It consists primarily of two polysaccharides: Chitin, composed of repeating units of N-acetyl-D-glucosamine (CAS No: 1398-61-4); Beta-(1,3)(1,6)-D-glucan, composed of repeating units of D-glucose (CAS No: 9041-22-9). The production process consists of several steps, including: cleaning, reduction in size and grinding, softening in water and heating in an alkaline solution, washing, drying. No hydrolysis is applied during the production process. Appearance: Powder, odourless, flavourless, brown Purity: Moisture: ≤ 15 % Ash: ≤ 3,0 % Chitin-glucan: ≥ 90 % Ratio of chitin to glucan: 70:20 Total carbohydrates, excluding glucans: ≤ 0,1 % Proteins: ≤ 2,0 % Lipids: ≤ 1,0 % Melanins: ≤ 8,3 % Additives: None pH: 6,7-7,5
	Lead (ppm): $\leq 1,00$

Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

Mercury (ppm): $\leq 0,03$ Arsenic (ppm): $\leq 0,20$ Microbiological criteria: Total mesophilic bacteria: $\leq 10^3/g$ Yeast and moulds: $\leq 10^3/g$ Coliforms at 30 °C: $\leq 10^3/g$ E. coli: $\leq 10/g$ Salmonella and other pathogenic bacteria: Absence/25 gChitosan extract from fungi (Agaricus) bisporus; Aspergillus niger)Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus miger. The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: ($C_6H_{11}NO_4$) Appearance: fine free-flowing powder Aspect: Off -white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus Ash (% w/w dry weight): ≤ 3.0
Microbiological criteria: Total mesophilic bacteria: $\leq 10^3/g$ Yeast and moulds: $\leq 10^3/g$ Coliforms at 30 °C: $\leq 10^3/g$ E. coli: $\leq 10/g$ Salmonella and other pathogenic bacteria: Absence/25 gChitosan extract from fungi (Agaricus bisporus; Aspergillus niger)Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus niger. The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan formula: (C ₆ H ₁₁ NO ₄) _n Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odour: Odouriess Purity: Chitosan content (% w/w dry weight): ≥ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
Total mesophilic bacteria: $\leq 10^3/g$ Yeast and moulds: $\leq 10^3/g$ Coliforms at $30 ^\circ$ C: $\leq 10^3/g$ E. coli: $\leq 10/g$ Salmonella and other pathogenic bacteria: Absence/25 gChitosan extract from fungi (Agaricus bisporus; Aspergillus niger)Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus niger. The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odour! So Purity: Chitosan content (% w/w dry weight): ≥ 15 Loss on drying (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
Yeast and moulds: $\leq 10^3/g$ Coliforms at 30 °C: $\leq 10^3/g$ <i>E. coli</i> : $\leq 10/g$ <i>Salmonella</i> and other pathogenic bacteria: Absence/25 gChitosan extract from fungi (<i>Agaricus</i> bisporus; <i>Aspergillus</i> niger)Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of <i>Agaricus bisporus</i> or from the mycelium of <i>Aspergillus niger</i> . The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: (C ₆ H ₁₁ NO ₄) _n Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Coliforms at 30 °C: $\leq 10^{3}/g$ E. coli: $\leq 10/g$ Salmonella and other pathogenic bacteria: Absence/25 gChitosanextract from fungi (Agaricus) bisporus; Aspergillus niger)Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus niger. The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off -white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
E. coli: $\leq 10/g$ Salmonella and other pathogenic bacteria: Absence/25 gChitosan extract from fungi (Agaricus bisporus; Aspergillus niger)Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus niger. The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO4)_n$ Appearance: fine free-flowing powder Aspect: Off -white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
Chitosan extract from fungi (Agaricus bisporus; Aspergillus miger)Description/Definition: The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus niger. The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
extract from fungi (Agaricus bisporus; Aspergillus niger)The chitosan extract (containing mainly poly(D-glucosamine)) is obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus niger.Aspergillus niger)The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
fungi (Agaricus bisporus; Aspergillus niger)obtained from stems of Agaricus bisporus or from the mycelium of Aspergillus niger.Aspergillus niger)The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
bisporus; Aspergillus niger)Aspergillus niger.The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
Aspergillus niger)The patented production process consists of several steps, including: extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
niger)extraction and deacetylation (hydrolysis) in alkaline medium, solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
solubilisation in acidic medium, precipitation in alkaline medium, washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
washing and drying. Synonym: Poly(D-glucosamine) Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Chitosan CAS number: 9012-76-4 Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Chitosan formula: $(C_6H_{11}NO_4)_n$ Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Appearance: fine free-flowing powder Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): \geq 85 Glucan content (% w/w dry weight): \leq 15 Loss on drying (% w/w dry weight): \leq 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Aspect: Off –white to slightly brownish Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Odour: Odourless Purity: Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Purity:Chitosan content (% w/w dry weight): \geq 85Glucan content (% w/w dry weight): \leq 15Loss on drying (% w/w dry weight): \leq 10Viscosity (1 % in 1 % acetic acid): 1-15Degree of acetylation (in % mol/wet weight): 0-30Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from Agaricus bisporus
Chitosan content (% w/w dry weight): ≥ 85 Glucan content (% w/w dry weight): ≤ 15 Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Loss on drying (% w/w dry weight): ≤ 10 Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Viscosity (1 % in 1 % acetic acid): 1-15 Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Degree of acetylation (in % mol/wet weight): 0-30 Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Viscosity (1 % in 1 % acetic acid) (mPa.s): 1-14 for chitosan from Aspergillus niger; 12-25 for chitin from <i>Agaricus bisporus</i>
Aspergillus niger; 12-25 for chitin from Agaricus bisporus
Proteins (% w/w dry weight): $\leq 2,0$
Particle size: > 100 nm
Tapped density (g/cm ³): 0,7-1,0
Fat binding capacity $800 \times (w/w \text{ wet weight})$: pass
Heavy metals: Moreovery $(nnm) \le 0.1$
Mercury (ppm): $\leq 0,1$ Lead (ppm): $\leq 1,0$
Arsenic (ppm): $\leq 1,0$
Cadmium (ppm): $\leq 0,5$
Microbiological criteria:
Aerobic count (CFU/g): $\leq 10^3$
Yeast and mould count (CFU/g): $\leq 10^3$
<i>Escherichia coli</i> (CFU/g): ≤ 10
Enterobacteriaceae (CFU/g): ≤ 10 a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in

 a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	Salmonella: Absence/25g Listeria monocytogenes: Absence/25g
Chondroitin sulphate	Description/Definition: Chondroitin sulphate (sodium salt) is a biosynthetic product. It is obtained by chemical sulphation of chondroitin derived from fermentation by the bacterium <i>Escherichia coli</i> O5:K4:H4 strain U1-41 (ATCC 23502). Chondroitin sulphate (sodium salt) (% dry basis): 95-105 MWw (weight avg.) (kDa): 5-12 MWn (number avg.) (kDa): 4-11 Dispersity ($w_h/w_{0,05}$): $\leq 0,7$ Sulphation pattern (Δ Di-6S) (%): ≤ 85 Loss on drying (%) (105 °C to constant weight): $\leq 10,0$ Residue on ignition (% dry basis): 20-30 Protein (% dry basis): $\leq 0,5$ Endotoxins (EU/mg): ≤ 100 Total organic impurities (mg/kg): ≤ 50
Chromium Picolinate	Description/Definition:Chromium picolinate is a reddish free-flowing powder, slightly soluble in water at pH 7. The salt is also soluble in polar organic solvents.Chemical name: tris(2pyridinecarboxylato-N,O)chromium(III) or 2- pyridinecarboxylic acid chromium(III) salt CAS No.: 14639-25-9Chemical formula: Cr(C ₆ H ₄ NO ₂) ₃ Chemical characteristics: Chromium Picolinate: \geq 95 % Chromium (III): 12-13 % Chromium (VI): not detected Water: \leq 4,0 %
<i>Cistus incanus</i> L. Pandalis herb	Description: <i>Cistus incanus</i> L. Pandalis herb; species belonging to the <i>Cistaceae</i> family and native to the Mediterranean region, Chalkidiki Peninsula. Composition: Moisture: 9–10 g/100 g herbs Protein: 6,1 g/100 g herbs Fat: 1,6 g/100 g herbs Carbohydrates: 50,1 g/100 g herbs Fiber: 27,1 g/100 g herbs Minerals: 4,4 g/100 g herbs Sodium: 0,18 g Potassium: 0,75 g Magnesium: 0,24 g Calcium: 1,0 g Iron: 65 mg Vitamin B ₁ : 3,0 μg
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,

Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	Vitamin B ₂ : $30 \ \mu g$ Vitamin B ₆ : $54 \ \mu g$ Vitamin C: $28 \ mg$ Vitamin A: less than 0,1 mg Vitamin E: $40-50 \ mg$ Alpha-Tocopherol: $20-50 \ mg$ Beta and Gamma-Tocopherols: $2-15 \ mg$ Delta-Tocopherol: $0,1-2 \ mg$
Citicoline	Description/Definition: Citicoline is produced by a microbial process. Citicoline is composed of cytosine, ribose, pyrophosphate and choline. White crystalline powder Chemical name: Choline cytidine 5'-pyrophosphate, Cytidine 5'- (trihydrogen diphosphate) P'-[2-(trimethylammonio)ethyl]ester inner salt Chemical formula: C ₁₄ H ₂₆ N ₄ O ₁₁ P ₂ Molecular weight: 488,32 g/mol CAS No.: 987-78-0 pH (sample solution of 1 %): 2,5-3,5 Purity: Assay value: ≥ 98 % of dry matter Loss on drying (100 °C for 4 hours): ≤ 5,0 % Armonium: ≤ 0,05 % Arsenic: Not more than 2 ppm Free phosphoric acids: ≤ 0,1 % 5'-Cytidylic acid: ≤ 1,0 % Microbiological criteria: Total plate count: ≤ 10 ³ CFU/g Yeast and moulds: ≤ 10 ² CFU/g <i>Escherichia coli</i> : Absence in 1 g
Clostridium butyricum	Description/Definition: <i>Clostridium butyricum</i> (CBM-588) is a Gram-positive, spore-forming, obligate anaerobic, non-pathogenic, non-genetically modified bacterium. Depository number FERM BP-2789 Microbiological criteria: Total viable aerobic count: $\leq 10^3$ CFU/g <i>Escherichia coli</i> : Not detected in 1 g <i>Staphylococcus aureus</i> : Not detected in 1 g <i>Pseudomonas aeruginosa</i> : Not detected in 1 g Yeast and moulds: $\leq 10^2$ CFU/g
Extract of defatted cocoa powder	Cocoa (<i>Theobroma cacao</i> L.) Extract Appearance: Dark brown powder free of visible impurities Physical and chemical properties: Polyphenol content: Min 55,0 % GAE Theobromine content: Max 10,0 %
	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

Low fat cocoa extract	Ash content: Max 5,0 % Moisture content: Max 8,0 % Bulk density: 0,40-0,55 g/cm ³ pH: 5,0-6,5 Residual solvent: Max 500 ppm Low fat Cocoa (<i>Theobroma cacao</i> L.) extract Appearance: Dark red to purple powder Cocoa extract, concentrate: Min 99 %
	Silicon dioxide (technological aid): Max 1,0 % Cocoa flavanols: Min. 300 mg/g — Epicatechin: Min. 45 mg/g Loss on drying: Max. 5,0 %
Coriander seed oil from <i>Coriandrum</i> <i>sativum</i>	Description/Definition: Coriander seed oil is an oil containing glycerides of fatty acids that is produced from the seeds of the coriander plant <i>Coriandrum sativum</i> L. Slight yellow colour, bland taste CAS No.: 8008-52-4 Composition of fatty acids: Palmitic acid (C16:0): 2-5 % Stearic acid (C18:0): < 1,5 % Petroselinic acid (cis-C18:1(n-12)): 60-75 % Oleic acid (cis-C18:1 (n-9)): 8-15 % Linoleic acid (C18:2): 12-19 % α -Linolenic acid (C18:3): < 1,0 % Purity: Refractive index (20 °C): 1,466-1,474 Acid value: $\leq 2,5$ mg KOH/g Peroxide value (PV): $\leq 5,0$ meq/kg Iodine value: 88-110 units Saponification value: 186-200 mg KOH/g Unsaponifiable matter: ≤ 15 g/kg
<i>Crataegus pinnatifida</i> dried fruit	Description/Definition: Dried fruits of <i>Crataegus pinnatifida</i> species belonging to the <i>Rosaceae</i> family and native to north China and Korea. Composition: Dry matter: 80 % Carbohydrates: 55 g/kg fresh weight Fructose: 26,5–29,3 g/100 g Glucose: 25,5–28,1 g/100 g Vitamin C: 29,1 mg/100 g fresh weight Sodium: 2,9 g/100 g fresh weight Compotes are products obtained by thermal processing of the edible part of one or several species of fruits, whole or in pieces, sieved or not,
Annexes II and III p. 1).	ulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in t to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020. Changes to legislation: There are currently no known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	without significant concentration. Sugars, water, cider, spices and lemon juice may be used.
a-cyclodextrin	juice may be used. Description/Definition: A non-reducing cyclic saccharide consisting of six α -1,4-linked D-glucopyranosyl units produced by the action of cyclodextrin glucosyltransferase (CGTase, EC 2.4.1.19) on hydrolyzed starch. Recovery and purification of α -cyclodextrin may be carried out using one of the following procedures: precipitation of a complex of α -cyclodextrin with 1-decanol, dissolution in water at elevated temperature and re- precipitation, steam-stripping of the complexant, and crystallisation of α -cyclodextrin from the solution; or chromatography with ion-exchange or gel filtration followed by crystallisation of α -cyclodextrin from the purified mother liquor; or membrane separation methods such as ultra- filtration and reverse osmosis: Description: Virtually odourless, white or almost white crystalline solid. Synonyms: α -cyclodextrin, α -dextrin, cyclohexaamylose, cyclomaltohexaose, α -cycloamylase Chemical name: Cyclohexaamylose CAS No.: 10016-20-3 Chemical formula: (C ₆ H ₁₀ O ₅) ₆ Formula weight: 972,85 Assay: \geq 98 % (dry basis) Identification: Melting range: Decomposes above 278 °C Solubility: Freely soluble in water; very slightly soluble in ethanol Specific rotation: [α] _D ²⁵ : Between +145° and +151° (1 % solution) Chromatogram of reference α -cyclodextrin (available from <i>Consortium</i> <i>für Elektrochemische Industrie GmbH, München, Germany or Wacker</i> <i>Biochem Group, Adrian, MI, USA</i>) using the conditions described in the METHOD OF ASSAY Purity: Water: \leq 11 % (Karl Fischer Method) Residual complexant: \leq 20 mg/kg (1-decanol) Reducing substances: \leq 0,5 % (as glucose) Sulphated ash: \leq 0,1 % Lead: \leq 0,5 mg/kg
	Method of assay: Determine by liquid chromatography using the following conditions: Sample solution: Weigh accurately about 100 mg of test sample into a 10 ml volumetric flask and add 8 ml of deionised water. Dissolve the sample completely using an ultra-sonification bath (10-15 min) and dilute to the work with purified deionised water. Filter through a 0.45
	dilute to the mark with purified deionised water. Filter through a 0,45- micrometer filter

Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, a p. 1).

Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable b to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	Reference solution: Weigh accurately about 100 mg of α -cyclodextrin into a 10 ml volumetric flask and add 8 ml of deionised water. Dissolve the sample completely using an ultra-sonification bath and dilute to the mark with purified deionised water. Chromatography: Liquid chromatograph equipped with a refractive index detector and an integrating recorder. Column and packing: Nucleosil-100-NH ₂ (10 µm) (<i>Macherey & Nagel</i> <i>Co. Düren</i> , Germany) or similar Length: 250 mm Diameter: 4 mm Temperature: 40 °C Mobile phase: acetonitrile/water (67/33, v/v) Flow rate: 2,0 ml/min Injection volume: 10 µl Procedure: Inject the sample solution into the chromatograph, record the chromatogram, and measure the area of the α -CD peak. Calculate the percentage of α -cyclodextrin in the test sample as follows: % α -cyclodextrin (dry basis) = 100 × (A _S /A _R) (W _R /W _S) where A _S and A _R are the areas of the peaks due to α -cyclodextrin for the sample solution and reference solution, respectively. W _S and W _R are the weights (mg) of the test sample and reference α - cyclodextrin, respectively, after correcting for water content.
γ-cyclodextrin	Description/Definition: A non-reducing cyclic saccharide consisting of eight α -1,4-linked D-glucopyranosyl units produced by the action of cyclodextrin glucosyltransferase (CGTase, EC 2.4.1.19) on hydrolysed starch. Recovery and purification of γ -cyclodextrin may be carried out by precipitation of a complex of γ -cyclodextrin with 8-cyclohexadecen-1- one, dissolution of the complex with water and n-decane, steam-stripping of the aqueous phase and recovery of gamma-CD from the solution by crystallisation. Virtually odourless, white or almost white crystalline solid Synonyms: γ -cyclodextrin, γ -dextrin, cyclooctaamylose, cyclomaltooctaose, γ -cycloamylase Chemical name: Cyclooctaamylose CAS number: 17465-86-0 Chemical formula: (C ₆ H ₁₀ O ₅) ₈ Assay: \geq 98 % (dry basis) Identification: Melting range: Decomposes above 285 °C Solubility: Freely soluble in water; very slightly soluble in ethanol Specific rotation: [α] _D ²⁵ : between + 174° and + 180° (1 % solution) Purity: Water: \leq 11 %
	Residual complexant (8-cyclohexadecen-1-one (CHDC)): $\leq 4 \text{ mg/kg}$ Ilation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed inI to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,

Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	Residual solvent (n-decane): $\leq 6mg/kg$ Reducing substances: $\leq 0.5 \%$ (as glucose) Sulphated ash: $\leq 0.1 \%$
Dextran preparation produced by <i>Leuconostoc</i> <i>mesenteroides</i>	 Powdered form: Carbohydrates: 60 % with: (Dextran: 50 %, Mannitol: 0,5 %, Fructose: 0,3 %, Leucrose: 9,2 %) Protein: 6,5 % Lipid: 0,5 % Lactic acid: 10 % Ethanol: traces Ash: 13 % Moisture: 10 % Liquid form: Carbohydrates: 12 % with: (Dextran: 6,9 %, Mannitol: 1,1 %, Fructose: 1,9 %, Leucrose: 2,2 %) Protein: 2,0 % Lipid: 0,1 % Lactic acid: 2,0 % Ethanol: 0,5 % Ash: 3,4 % Moisture: 80 %
Diacylglycerol oil of plant origin	Description/Definition: Manufactured from glycerol and fatty acids derived from edible vegetable oils, in particular from soybean oil (<i>Glycine max</i>) or rapeseed oil (<i>Brassica campestris, Brassica napus</i>) using a specific enzyme. Acylglycerol Distribution: Diacylglycerols (DAG): \geq 80 % 1,3-Diacylglycerols (1,3-DAG): \geq 50 % Triacylglycerols (TAG): \leq 20 % Monoacylglycerols (MAG): \leq 5,0 % Fatty Acid Composition (MAG, DAG, TAG): Oleic acid (C18:1): 20-65 % Linolenic acid (C18:2): 15-65 % Linolenic acid (C18:3): \leq 15 % Saturated fatty acids: \leq 10 % Others: Acid value: \leq 0,5 mg KOH/g Moisture and volatile: \leq 0,1 % Peroxide value (PV): \leq 1,0 meq/kg Unsaponifiables: \leq 2,0 % Trans fatty acids \leq 1,0 % MAG = monoacylglycerols, DAG = diacylglycerols, TAG = triacylglycerols
Dihydrogonsiato	Description/Definition.

Dihydrocapsiate Description/Definition: (DHC)

 a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020. ~~

Status: Point in time view as at 51/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Dihydrocapsiate is synthesised by enzyme-catalysed esterification of vanillyl alcohol and 8-methylnonanoic acid. Following the esterification dihydrocapsiate is extracted with n-hexane. Viscous to colourless to yellow liquid Chemical formula: C_{18} H ₂₈ O ₄ CAS No: 205687-03-2 Physical-chemical properties: Dihydrocapsiate: > 94 % 8-Methylnonanoic acid: < 6,0 % Vanillyl acohol: < 1,0 % Other synthesis related substances: < 2,0 %
Dried extract of <i>Lippia citriodora</i> from cell cultures	Description/Definition: Dried extract of <i>Lippia citriodora</i> (Palau) Kunth from cell cultures HTN [®] Vb.
<i>Echinacea</i> <i>angustifolia</i> extract from cell cultures	Description/Definition: Extract of the roots of <i>Echinacea angustifolia</i> obtained from plant tissue culture which is substantially equivalent to a root extract from <i>Echinacea angustifolia</i> obtained in ethanol-water titrated to 4 % echinacoside.
<i>Echinacea</i> <i>purpurea</i> extract from cell cultures	Description/Definition: Dried extract of <i>Echinacea purpurea</i> from cell cultures HTN [®] Vb
<i>Echium plantagineum</i> oil	Description/Definition: Echium oil is the pale yellow product obtained by refining oil extracted from the seeds of <i>Echium plantagineum</i> L. Stearidonic acid: ≥ 10 % w/w of total fatty acids Trans fatty acids: $\leq 2,0$ % (w/w of total fatty acids) Acid value: $\leq 0,6$ mg KOH/g Peroxide value (PV): $\leq 5,0$ meq O ₂ /kg Unsaponifiable content: $\leq 2,0$ % Protein content (total nitrogen): ≤ 20 µg/ml Pyrrolizidine alkaloids: Not detectable with a detection limit 4,0 µg/kg
Epigallocatechin gallate as a purified extract from green tea leaves (<i>Camellia</i> <i>sinensis</i>)	Description/Definition: A highly purified extract from the leaves of green tea (<i>Camellia sinensis</i> (<i>L.</i>) <i>Kuntze</i>) in the form of a fine, off-white to pale pink powder. It is composed of a minimum of 90 % epigallo-catechin gallate (EGCG), and has a melting point between approx. 210 and 215 °C Appearance: off-white to pale pink powder Chemical name: polyphenol (-) epigallocatechin-3-gallate Synonyms: epigallocatechin gallate (EGCG) CAS No.: 989-51-5 INCI name: epigallocatechin gallate Molecular mass: 458,4 g/mol
	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and .2.2015, p. 10).

	Loss on drying: m Heavy metals: Arsenic: max 3,0 Lead: max 5,0 pp Assay: Min. 94 % EGCG max. 0,1 % caffei Solubility: EGCG acetone	ppm m (on dry material) ne	water, ethanol, methanol and
L-ergothioneine	Definition Chemical name (IUPAC): (2S)-3-(2-thioxo-2,3-dihydro-1 <i>H</i> -imidazol-4- yl)-2-(trimethylammonio)-Propanoate Chemical formula: C ₉ H ₁₅ N ₃ O ₂ S Molecular mass: 229,3 Da CAS No.: 497-30-3		
	Parameter	Specification	Method
	Appearance	White powder	Visual
	Optical rotation	$\begin{split} & [\alpha]_D \ge (+) \ 122^{\circ} \\ & (c = 1, H_2O)^{a)} \end{split}$	Polarimetry
	Chemical purity	≥ 99,5 % ≥ 99,0 %	HPLC [Eur. Ph. 2,2.29] 1H-NMR
	Identification	Compliant with the structure C: $47,14 \pm 0,4 \%$ H: $6,59 \pm 0,4 \%$ N: $18,32 \pm 0,4 \%$	1H-NMR Elemental analysis
	Total residual solvents (methanol, ethyl acetate, isopropanol, ethanol)	[Eur. Ph. 01/2008:50400] < 1 000 ppm	Gas chromatography [Eur. Ph. 01/2008:20424]
	Loss on drying	Internal standard < 0,5 %	[Eur. Ph. 01/2008:20232]
	Impurities	< 0,8 %	HPLC/GPC or 1H-NMR
	Heavy metals ^{b) c)}		
	Lead	< 3,0 ppm	ICP/AES
	Cadmium	< 1,0 ppm	(Pb, Cd)
	Mercury	< 0,1 ppm	Atomic fluorescence (Hg)

 a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.

Changes to legislation: There are currently no known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Microbiological s	specifications ^{b)}	
	Total viable aerobic count (TVAC)	$\leq 1 \times 10^3 $ CFU/g	[Eur. Ph. 01/2011:50104]
	Total yeast and mould count (TYMC)	$\leq 1 \ge 10^2 \text{ CFU/g}$	
	Escherichia coli	Absence in 1 g	
	resonance; HPLC:permeation chromatomic emission sCFU: colony-forma)Lit. $[\alpha]_D$ b)	high-performance atography; ICP/Al pectroscopy; ning units. = (+) 126,6° (c = $\frac{1}{2}$ s conducted on eac m levels in accorda	/
erric Sodium	Description/Defin		
EDTA	Ferric Sodium EDTA (ethylenediaminetetraacetic acid) is an odourless free-flowing, yellow to brown powder with a chemical purity of more than 99 % (w/w). It is freely soluble in water. Chemical formula: $C_{10}H_{12}FeN_2NaO_8 * 3H_2O$ Chemical characteristics: pH of 1 % solution: 3,5-5,5 Iron: 12,5-13,5 % Sodium: 5,5 % Water: 12,8 % Organic matter (CHNO): 68,4 % EDTA: 65,5-70,5 % Water insoluble matter: $\leq 0,1$ % Nitrilo-triacetic acid: $\leq 0,1$ %		
Ferrous ammonium ohosphate	insoluble in water CAS No.: 10101-0 Chemical formula Chemical character	m phosphate is a g and soluble in dilu 60-7 : FeNH ₄ PO ₄ eristics: sion in water: 6,8-7 % (w/w) (w/w)	
			wn specifications for food additives listed in Parliament and of the Council (OJ L 83, 22.3.2012

to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Fish peptides from <i>Sardinops</i> <i>sagax</i>	Description/Definition:The novel food ingredient is a peptide mixture, which is obtained by an alkaline protease-catalysed hydrolysis of fish (Sardinops sagax) muscle, subsequent isolation of the peptide fraction by column chromatography, concentration under vacuum and spray drying.Yellowish white powderPeptides (¹) (short chain peptides, dipeptides and tripeptides with a molecular weight of less than 2 kDa): ≥ 85 g/100 g Val-Tyr (dipeptide): 0,1-0,16 g/100 g Moisture: ≤ 8 g/100 g(¹)Kjeldahl method
Flavonoids from Glycyrrhiza glabra	Description/Definition: Flavonoids derived from the roots or rootstock of <i>Glycyrrhiza glabra</i> L. are extracted with ethanol followed by further extraction of this ethanolic extract with medium-chain triglycerides. It is a dark-brown coloured liquid, containing 2,5 % to 3,5 % of glabridin. Moisture: < 0,5 % Ash: < 0,1 % Peroxide value (PV): < 0,5 meq/kg Glabridin: 2,5-3,5 % of fat Glycyrrhizinic acid: < 0,005 % Fat including polyphenol-type substances: \geq 99 % Protein: < 0,1 % Carbohydrates: not detectable
Fucoidan extract from the seaweed <i>Fucus</i> <i>vesiculosus</i>	Description/Definition: Fucoidan from the seaweed <i>Fucus vesiculosus</i> is extracted using aqueous extraction in acidic solution and filtration processes without the use of organic solvents. The resulting extract is concentrated and dried to yield the fucoidan extract with the following specifications: Off-white to brown powder Odour and Taste: Bland odour and taste Moisture: < 10 % (105 °C for 2 hours) pH value: 4,0-7,0 (1 % suspension at 25 °C) Heavy metals: Arsenic (inorganic): < 1,0 ppm Cadmium: < 3,0 ppm Lead: < 2,0 ppm Mercury: < 1,0 ppm Microbiological criteria: Total aerobic microbial count: < 10 000 CFU/g Yeast and mould count: < 100 CFU/g Total enterobacteria count: Absence/g <i>Escherichia coli</i> : Absence/g <i>Salmonella</i> : Absence/10 g

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.

Changes to legislation: There are	currently no known outstanding effects for the
Commission Implementing Regulation	(EU) 2018/1023. (See end of Document for details)

	Staphylococcus aureus: Absence/gComposition of the two permitted types of extracts, based on the level offucoidan:Extract 1:Fucoidan: 75-95 %Alginate: 2,0-5,5 %Polyphloroglucinol: 0,5-15 %Mannitol: 1-5 %Natural salts/Free Minerals: 0,5-2,5 %Other carbohydrates: 0,5-1,0 %Protein: 2,0-2,5 %Extract 2:Fucoidan: 60-65 %Alginate: 3,0-6,0 %Polyphloroglucinol: 20-30 %Mannitol: < 1,0 %Natural salts/Free Minerals: 0,5-2,0 %Other carbohydrates: 0,5-2,0 %Protein: 2,0-2,5 %
Fucoidan extract from the seaweed Undaria pinnatifida	Description/Definition:Fucoidan from seaweed Undaria pinnatifida is extracted using aqueous extraction in acidic solution and filtration processes without the use of organic solvents. The resulting extract is concentrated and dried to yield the fucoidan extract with the following specifications: Off-white to brown powder Odour and Taste: Bland odour and taste Moisture: < 10 % (105 °C for 2 hours) pH value: 4,0-7,0 (1 % suspension at 25 °C) Heavy metals: Arsenic (inorganic): < 1,0 ppm Cadmium: < 3,0 ppm Lead: < 2,0 ppm Mercury: < 1,0 ppm
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

	Document Generated: 2024-07-6 Status: Point in time view as at 31/01/2020.
	Changes to legislation: There are currently no known outstanding effects for the
Com	mission Implementing Regulation (EU) 2018/1023. (See end of Document for details)
	Natural salts/Free Minerals: 0,5-1,0 % Other carbohydrates: 0,5-2,0 % Protein: 2,0-2,5 % <i>Extract 2:</i> Fucoidan: 50-55 % Alginate: 2,0-4,0 % Polyphloroglucinol: 1,0-3,0 % Mannitol: 25-35 % Natural salts/Free Minerals: 8-10 % Other carbohydrates: 0,5-2,0 % Protein: 1,0-1,5 %
2'- Fucosyllactose (synthetic)	Definition: Chemical name: α-L-Fucopyranosyl-(1→2)-β-D-galactopyranosyl- (1→4)- D-glucopyranose Chemical formula: C18H32O15 CAS No: 41263-94-9 Molecular weight: 488,44 g/mol Description: 2'-fucosyllactose is a white to off-white powder that is produced by a chemical synthesis process. Purity: 2'-Fucosyllactose: ≥ 95 % D-Lactose: ≤ 1,0 w/w % Difucosyl-D-lactulose isomers: ≤ 1,0 w/w % Difucosyl-D-lactulose: ≤ 0,6 w/w % pH (20 °C, 5 % solution): 3,2-7,0 Water (%): ≤ 9,0 % Ash, sulphated: ≤ 0,2 % Acetic acid: ≤ 0,3 % Residual solvents (methanol, 2-propanol, methyl acetate, acetone): ≤ 50,0 mg/kg singly, ≤ 200,0 mg/kg in combination Residual proteins: ≤ 0,01 % Heavy Metals: Palladium: ≤ 0,1 mg/kg Nickel: ≤ 3,0 mg/kg Microbiological criteria: Aerobic mesophilic bacteria total count: ≤ 500 CFU/g Yeasts and Moulds: ≤ 10 CFU/g Residual endotoxins: ≤ 10 EU/mg
2'- Fucosyllactose (microbial source)	Definition: Chemical name: α -L-Fucopyranosyl- $(1\rightarrow 2)$ - β -D-galactopyranosyl- $(1\rightarrow 4)$ -D-glucopyranose Chemical formula: C ₁₈ H ₃₂ O ₁₅ CAS No: 41263-94-9 Molecular weight: 488,44 g/mol
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable aar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.

Changes to legislation: There are currently no known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

Source: Genetically modified strain of	Source: Genetically modified strain of
Escherichia coli K-12	Escherichia coli BL21
Description: 2'-Fucosyllactose is a white to off- white powder that is produced by a microbial process. Purity: 2'-Fucosyllactose: $\geq 90 \%$ D-Lactose: $\leq 3,0 \%$ L-Fucose: $\leq 2,0$ Difucosyl-D-lactose: $\leq 2,0 \%$ 2'-Fucosyl-D-lactulose: $\leq 1,0 \%$ pH (20 °C, 5 % solution): 3,0-7,5 Water: $\leq 9,0 \%$ Sulphated ash: $\leq 2,0 \%$ Acetic acid: $\leq 1,0 \%$ Residual proteins: $\leq 0,01 \%$ Microbiological criteria: Aerobic mesophilic bacteria total count: $\leq 3 000 \text{ CFU/g}$ Yeasts: $\leq 100 \text{ CFU/g}$ Moulds: $\leq 100 \text{ CFU/g}$ Endotoxins: $\leq 10 \text{ EU/mg}$	Description: 2'-Fucosyllactose is a white to off white powder and the liquid concentrate ($45 \% \pm 5 \%$ w/v) aqueous solution is a colourless to slight yellow clear aqueous solution. 2'-Fucosyllactose is produced by a microbiological process. Purity: 2'-Fucosyllactose: $\geq 90 \%$ Lactose: $\leq 5,0 \%$ Fucose: $\leq 3,0 \%$ 3-Fucosyllactose: $\leq 5,0 \%$ Fucosylgalactose: $\leq 5,0 \%$ Glucose: $\leq 3,0 \%$ Glucose: $\leq 3,0 \%$ Glactose: $\leq 3,0 \%$ Galactose: $\leq 3,0 \%$ Water: $\leq 9,0 \%$ (powder) Ash, sulphated: $\leq 0,5 \%$ (powder and liquid) Residual proteins: $\leq 0,01 \%$ (powder and liquid) Heavy Metals: Lead: $\leq 0,02 \text{ mg/kg}$ (powder and liquid); Arsenic: $\leq 0,2 \text{ mg/kg}$ (powder and liquid) Cadmium: $\leq 0,1 \text{ mg/kg}$ (powder and liquid) Mercury: $\leq 0,5 \text{ mg/kg}$ (powder and liquid) Mercury: $\leq 0,5 \text{ mg/kg}$ (powder and liquid) Mercury: $\leq 5 000 \text{ CFU/g}$ (liquid) Yeasts and Moulds: $\leq 100 \text{ CFU/g}$ (powder), $\leq 5 000 \text{ CFU/g}$ (liquid) Enterobacteriaceae/Coliforms: absence in 11g (powder and liquid) <i>Salmonella</i> : negative/100 g (powder), negative/200 ml (liquid) <i>Cronobacter</i> : negative/100 g (powder), negative/200 ml (liquid)

a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Galacto- oligosaccharide	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	
Glucosamine HCl from Aspergillus niger and genetically modified strain of <i>E. coli</i> K-12	White crystalline odourless powder Molecular formula: $C_6H_{13}NO_5 \cdot HCl$ Relative molecular mass: 215,63 g/mol D-Glucosamine HCl 98,0-102,0 % of reference standard (HPLC) Specific rotation + 70,0° - + 73,0°	
Glucosamine sulphate KCl from <i>Aspergillus</i> <i>niger</i> and genetically modified strain of <i>E. coli</i> K-12	White crystalline odourless powder Molecular formula: $(C_6H_{14}NO_5)_2SO_4 \cdot 2KCl$ Relative molecular mass: 605,52 g/mol D-Glucosamine Sulphate 2KCl 98,0-102,0 % of reference standard (HPLC) Specific Rotation +50,0° to +52,0°	
Glucosamine sulphate NaCl from <i>Aspergillus</i> <i>niger</i> and genetically modified strain of <i>E. coli</i> K-12	White crystalline odourless powder Molecular formula: $(C_6H_{14}NO_5)_2SO_4 \cdot 2NaCl$ Relative molecular mass: 573,31 g/mol D-Glucosamine HCl: 98-102 % of reference standard (HPLC) Specific Optical Rotation: $+52^\circ - +54^\circ$	
Guar Gum	Description/Definition: Native guar gum is the ground endosperm of seeds from natural strains of guar <i>Cyamopsis tetragonolobus</i> L. Taub. (<i>Leguminosae</i> family). It consists of a high molecular weight polysaccharide, primarily composed of galactopyranose and mannopyranose units combined	
Annexes II and III p. 1). b Commission Imple	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and	
dioxins (OJ L 30, 6		

Status: Point in time view as at 31/01/2020.

Changes to legislation: There are currently no known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	 through glycosidic linkages, which may be described chemically as a galactomannan (galactomannan content not less than 75 %). Appearance: White to yellowish powder Molecular weight: Between 50 000 – 8 000 000 Daltons CAS number: 9000-30-0 Einecs Number: 232-536-8 Purity: As specified by Commission Regulation (EU) No 231/2012 layin down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council^a & by Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins^b. Physico-chemical properties:
	Powder Shelf-life: 2 years Colour: White Odour: Light
	Average diameter of particles: 60-70µm Moisture: Max 15 % Viscosity * at 1 hour — Viscosity * at 2 hours: Min 3 600 mPa.s Viscosity * at 24 hours: Min 4 000 mPa.s Solubility: Soluble in hot and cold water
	pH for 10g/L, at 25 °C - 6-7,5 Flakes Useful life: 1 year Colour: White/off white with absence or minimal presence of black spots Odour: Light Average diameter of particles: 1-10 mm
	Moisture: Max 15 % Viscosity * at 1 hour: Min 3 000 mPa.s Viscosity * at 2 hours — Viscosity * at 24 hours — Solubility — Soluble in hot and cold water pH for 10g/L, at 25 °C - 5-7,5
	(*) The measurements of viscosity are carried out under the following conditions: 1 %, 25 °C, 20 rpm
Heat-treated milk products fermented with <i>Bacteroides</i> <i>xylanisolvens</i>	Description/Definition: Heat-treated fermented milk products are produced with <i>Bacteroides xylanisolvens</i> (DSM 23964) as starter culture. Semi-skimmed milk (between 1,5 % and 1,8 % fat) or skimmed milk (0,5 % fat or less) is pasteurised or ultra-heat-treated before starting the fermentation with <i>Bacteroides xylanisolvens</i> (DSM 23964). The resulting fermented milk product is homogenised and then heat-treated to inactivate <i>Bacteroides xylanisolvens</i> (DSM 23964). The final product
	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012
	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable argum originating in or consigned from India due to contamination risks by pentachlorophenol and

	Status: Point in time view as at 31/01/2020. Changes to legislation: There are currently no known outstanding effects for the nission Implementing Regulation (EU) 2018/1023. (See end of Document for details)
	does not contain viable cells of <i>Bacteroides xylanisolvens</i> (DSM 23964) (¹).
	$(^{1})$ Modified DIN EN ISO 21528-2.
Hydroxytyrosol	Description/Definition: Hydroxytyrosol is a pale yellow viscous liquid obtained by chemical synthesis Molecular formula: $C_8H_{10}O_3$ Molecular weight: 154,6 g/mol CAS No: 10597-60-1 Moisture ≤ 0,4 % Odour: Characteristic Taste: Slightly bitter Solubility (water): Miscible with water pH: 3,5-4,5 Refractive Index: 1,571-1,575 Purity: Hydroxytyrosol: ≥ 99 % Acetic acid: ≤ 0,4 % Hydroxytyrosol acetate: ≤ 0,3 % Sum of homovanillic acid, iso-homovanilic acid, and 3- methoxy-4hydroxyphenylglycol: ≤ 0,3 % Heavy Metals Lead: ≤ 0,03 mg/kg Cadmium: ≤ 0,01 mg/kg Mercury: ≤ 0,01 mg/kg Residual Solvents Ethyl acetate: ≤ 25,0 mg/kg Isopropanol: ≤ 2,50 mg/kg
Ice Structuring Protein type III HPLC 12	Tetrahydrofuran: $\leq 0,01 \text{ mg/kg}$ Description/Definition: The Ice Structuring Protein (ISP) preparation is a light-brown liquid produced by submerged fermentation of a genetically-modified strain of food-grade baker's yeast (<i>Saccharomyces cerevisiae</i>) in which a synthetic gene for the ISP has been inserted into the yeast's genome. The protein is expressed and secreted into the growth medium where it is separated from the yeast cells by micro-filtration and concentrated by ultra-filtration. As a result, the yeast cells are not transferred into the ISP preparation as such or under an altered form. The ISP preparation consists of native ISP, glycosylated ISP and proteins and peptides from the yeast and sugars as well as acids and salts commonly found in food. The concentrate is stabilised with 10 mM citric acid buffer. Assay: ≥ 5 g/l active ISP pH: 2,5-3,5 Ash: $\leq 2,0 \%$
	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10). Status: Point in time view as at 31/01/2020.

Changes to legislation: There are currently no known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

DNA: Not detectable

Aqueous extract of dried leaves of <i>Ilex guayusa</i>	Description/Definition: Dark brown liquid. Aqueous extracts of dried leaves of <i>Ilex guayusa</i> . Composition: Protein: < 0,1 g/100 ml Fat: < 0,1 g/100 ml Carbohydrate: 0,2–0,3 g/100 ml Total sugars: < 0,2 g/100 ml Caffeine: 19,8–57,7 mg/100 ml Theobromine: 0,14–2,0 mg/100 ml Chlorogenic acids: 9,9–72,4 mg/100ml
Isomalto- oligosaccharide	Powder: Solubility (water) (%): > 99 Glucose (% dry basis): \leq 5,0 Isomaltose + DP3 to DP9 (% dry basis): \geq 90 Moisture (%): \leq 4,0 Sulphated ash(g/100 g): \leq 0,3 Heavy metals: Lead (mg/kg): \leq 0,5 Arsenic (mg/kg): \leq 0,5 Syrup: Dried solids (g/100 g): > 75 Glucose (% dry basis): \leq 5,0 Isomaltose + DP3 to DP9 (% dry basis): \geq 90 pH: 4 - 6 Sulphated ash(g/100 g): \leq 0,3 Heavy metals: Lead (mg/kg): \leq 0,5 Arsenic (mg/kg): \leq 0,5
Isomaltulose	Description/Definition: A reducing disaccharide that consists of one glucose and one fructose moiety linked by an alpha-1,6-glycosidic bond. It is obtained from sucrose by an enzymatic process. The commercial product is the monohydrate. Appearance: Virtually odourless, white or almost white crystals with a sweet taste Chemical name: $6-O-\alpha$ -D-glucopyranosyl-D-fructofuranose, monohydrate CAS No.: 13718-94-0 Chemical formula: $C_{12}H_{22}O_{11} \cdot H_2O$ Structural formula
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012
b Commission Imple	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).

dioxins (OJ L 30, 6.2.2015, p. 10).

	HOW HO HO HO HO
	Formula weight: 360,3 (monohydrate) Purity: Assay: \geq 98 % on the dry basis Loss on drying: \leq 6,5 % (60 °C, 5 hours) Heavy metals: Lead: \leq 0,1 mg/kg Determine using an atomic absorption technique appropriate to the specified level. The selection of sample size and method of sample preparation may be based on the principles of the method described in FNP 5(¹), 'Instrumental methods'
	(¹) Food and Nutrition Paper 5 Rev. 2 — Guide to specifications for general notices, general analytical techniques, identification tests, test solutions and other reference materials (JECFA), 1991, 322 pp., English, ISBN 92-5-102991-1.
Lactitol	Description/Definition: Crystalline powder or colourless solution manufactured via catalytic hydrogenation of lactose. Crystalline products occur in anhydrous, monohydrate and dihydrate forms. Nickel is used as a catalyst. Chemical name: 4-O- β -D-Galactopyranosyl-D-glucitol Chemical formula: C ₁₂ H ₂₄ O ₁₁ Molecular weight: 344,31 g/mol CAS No: 585-86-4 Purity: Solubility (in water): Very soluble in water Specific rotation [α] _D ²⁰ = + 13° to + 16° Assay: \geq 95 % d.b (d.b — expressed on the dry weight basis) Water: \leq 10,5 % Other polyols: \leq 2,5 % d.b Reducing sugars: \leq 0,2 % d.b Chlorides: \leq 100 mg/kg d.b Sulphates: \leq 200 mg/kg d.b
	Sulphated ash: $\leq 0,1$ % d.b Nickel: $\leq 2,0$ mg/kg d.b lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	Arsenic: $\leq 3,0 \text{ mg/kg d.b}$ Lead: $\leq 1,0 \text{ mg/kg d.b}$
Lacto-N- neotetraose (synthetic)	Definition: Chemical name: β -D-Galactopyranosyl- $(1 \rightarrow 4)$ -2-acetamido-2- deoxy- β -D-glucopyranosyl- $(1 \rightarrow 3)$ - β -D-galactopyranosyl- $(1 \rightarrow 4)$ - D- glucopyranose Chemical formula: C ₂₆ H ₄₅ NO ₂₁ CAS No: 13007-32-4 Molecular weight: 707,63 g/mol Description: Lacto-N-neotetraose is a white to off-white powder. Produced by a chemical synthesis process and is isolated by crystallisation. Purity: Assay (water free): ≥ 96 % D-Lactose: $\leq 1,0$ % Lacto-N-neotetraose fructose isomer: $\leq 0,6$ % pH (20 °C, 5 % solution): 5,0-7,0 Water: $\leq 9,0$ % Ash, sulphated: $\leq 0,4$ % Acetic acid: $\leq 0,3$ % Residual solvents (methanol, 2-propanol, methyl acetate, acetone): ≤ 50 mg/kg singly, ≤ 200 mg/kg in combination Residual proteins: $\leq 0,01$ % Palladium: $\leq 0,1$ mg/kg Nickel: $\leq 3,0$ mg/kg Microbiological criteria: Aerobic mesophilic bacteria total count: ≤ 500 CFU/g Yeasts: ≤ 10 CFU/g Moulds: ≤ 10 CFU/g Residual endotoxins: ≤ 10 EU/mg
Lacto- <i>N</i> - neotetraose (microbial source)	Definition: Chemical name: β -D-Galactopyranosyl- $(1 \rightarrow 4)$ -2-acetamido-2-deoxy- β - D-glucopyranosyl- $(1 \rightarrow 3)$ - β -D-galactopyranosyl- $(1 \rightarrow 4)$ -D-glucopyranose Chemical formula: $C_{26}H_{45}NO_{21}$ CAS No: 13007-32-4 Molecular weight: 707,63 g/mol Source: Genetically modified strain of <i>Escherichia coli</i> K-12 Description: Lacto- <i>N</i> -neotetraose is a white to off-white powder that is produced by a microbiological process. Lacto- <i>N</i> -neotetraose is isolated by crystallisation. Purity: Assay (water free): ≥ 92 % D-Lactose: $\leq 3,0$ %
	ilation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable are gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Lacto-N-triose II: $\leq 3,0 \%$ <i>para</i> -Lacto-N-neohexaose: $\leq 3,0 \%$ Lacto-N-neotetraose fructose isomer: $\leq 1,0 \%$ pH (20 °C, 5 % solution): 4,0-7,0 Water: $\leq 9,0 \%$ Ash, sulphated: $\leq 0,4 \%$ Residual solvents (methanol): $\leq 100 \text{ mg/kg}$ Residual proteins: $\leq 0,01 \%$ Microbiological criteria: Aerobic mesophilic bacteria total count: $\leq 500 \text{ CFU/g}$ Yeasts: $\leq 10 \text{ CFU/g}$ Moulds: $\leq 10 \text{ CFU/g}$ Residual endotoxins: $\leq 10 \text{ EU/mg}$
Lucerne leaf extract from <i>Medicago sativa</i>	Description/Definition: The Lucerne (<i>Medicago sativa</i> L.) is processed within 2 hours after harvest. It is chopped and crushed. By passing through an oleaginous- type press, the Lucerne provides a fibrous residue and press juice (10 % of dry matter). The dry matter of this juice contains about 35 % of crude protein. The press juice (pH 5,8-6,2) is neutralised. Preheating and vapour injection allows coagulation of proteins associated with carotenoid and chlorophyll pigments. The protein precipitate is separated by centrifugation and thereafter dried. After adding ascorbic acid the Lucerne protein concentrate is granulated and stored in inert gas or in cold storage. Composition: Protein: 45-60 % Fat: 9-11 % Free carbohydrates (soluble fibre): 1-2 % Polysaccharides (insoluble fibre): 11-15 % including cellulose: 2-3 % Minerals: 8-13 % Saponins: $\leq 1,4$ % Isoflavones: ≤ 350 mg/kg Coumestrol: ≤ 100 mg/kg Phytates: ≤ 200 mg/kg L-canavanine: $\leq 4,5$ mg/kg
Lycopene	Description/Definition: Synthetic lycopene is produced by the Wittig condensation of synthetic intermediates commonly used in the production of other carotenoids used in food. Synthetic lycopene consists of \geq 96 % lycopene and minor quantities of other related carotenoid components. Lycopene is presented either as a powder in a suitable matrix or an oily dispersion. The colour is dark red or red-violet. Antioxidative protection has to be assured. Chemical name: Lycopene CAS No.: 502-65-8 (<i>all</i> -trans lycopene) Chemical formula: C ₄₀ H ₅₆
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

Formula weight: 536,85 Da				
Lycopene from BlakesleaDescription/Definition: The purified lycopene from Blakeslea trispora consists of ≥ 95 % lycopene and ≤ 5 % other carotenoids. It is presented either as a point a suitable matrix or an oily dispersion. The colour is dark red on violet. Anti-oxidative protection has to be assured. Chemical name: Lycopene CAS No.: 502-65-8 (all trans lycopene) Chemical formula: C40H56 Formula weight: 536,85 Da				
Lycopene from tomatoes	Description/Definition: The purified lycopene from tomatoes (<i>Lycopersicon esculantum</i> L.) consists of \geq 95 % lycopene and \leq 5 % other carotenoids. It is presented either as a powder in a suitable matrix or an oily dispersion. The colour i dark red or red-violet. Anti-oxidative protection has to be assured. Chemical name: Lycopene CAS No.: 502-65-8 (all trans lycopene) Chemical formula: C ₄₀ H ₅₆ Formula weight: 536,85 Da			
Lycopene oleoresin from tomatoes	Description/Definition: Lycopene oleoresin from tomatoes is obtained by solvent extraction of ripe tomatoes (<i>Lycopersicon esculentum Mill.</i>) with subsequent removal of the solvent. It is a red to dark brown viscous, clear liquid. Total lycopene: 5-15 % Thereof trans-lycopene: 90-95 % Total carotenoids (calculated as lycopene): 6,5-16,5 % Other carotenoids: 1,75 % (Phytoene/phytofluene/ β -carotene): (0,5-0,75/0,4-0,65/0,2-0,35 %) Total tocopherols: 1,5-3,0 % Unsaponifiable matter: 13-20 % Total fatty acids: 60-75 % Water (Karl Fischer): \leq 0,5 %			
Magnesium citrate malate	Description/Definition:Magnesium citrate malate is a white to yellowish-white, amorphous powder.Chemical formula: Mg ₅ (C ₆ H ₅ O ₇) ₂ (C ₄ H ₄ O ₅) ₂ Chemical name: Pentamagnesium di-(2-hydroxybutanedioate)-di-(2- hydroxypropane-1,2,3-tricarboxylate)CAS No.: 1259381-40-2Molecular weight: 763,99 Daltons (anhydrous)Solubility: Freely soluble in water (about 20 g in 100 ml)Description of the physical state: Amorphous powderAssay magnesium: 12,0-15,0 %Loss on drying (120 °C/4 hours): ≤ 15 %Colour (solid): White to yellowish-white			
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,			
to the import of gu	 b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10). 			

Status: Point in time view as al 310/1200. Clanges to legislation: There are currently in known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details) Colour (20 % aqueous solution): Colourless to yellowish Appearance (20 % aqueous solution): Clear solution pH (20 % aqueous solution): Approx. 6,0 Impurities: Chloride: \$ 0,05 % Sulphate: \$ 0,05 % Sulphate: \$ 0,05 % Sulphate: \$ 0,05 % Sulphate: \$ 0,05 % Magnolia Bark Description/Definition: Magnolia bark extract is obtained from the bark of the plant Magnolia of dictinals 1. and produced with supercritical carbon dioxide. The bark is washed and oven dried to reduce moisture content before being crushed and extracted with supercritical carbon dioxide. The bark is washed and oven dried to reduce moisture content before being crushed and extract e thanol and re-crystallised to yield magnolia bark extract. Magnolia bark extract is mainly composed of two phenolic compounds, magnoloi ald honokiol. Appearance: Light brownish powder Purity: Magnoliol & Honokiol: \$ 2 % Magnoliol & Honokiol: \$ 2 % Methyl eugenol (ppm): \$ 0,5 Methyl eugenol (ppm): \$ 0,0 Total Alkaloid (ppm): \$ 0,0 Total Alkaloid (ppm): \$ 0,0 Maize-germ oil high in unsaponifiable matter is produced by vacuum unsaponifiable matter: \$ 0,0 g/100 g Total Alkaloid (ppm): \$ 0,0 for tal Alkaloid (ppm): \$ 0,0 g T		Document Generated: 2024-07-0	
Appearance (20 % aqueous solution): Clear solution H (20 % aqueous solution): Approx. 6,0 Impurities: Chloride: ≤ 0,05 % Sulphate: ≤ 0,05 % Sulphate: ≤ 0,05 % Arsenic:: ≤ 3,0 ppm Lead: ≤ 2,0 ppm Cadmium: ≤ 1 ppm Mercury: ≤ 0,1 ppm Magnolia Bark Extract Øficinalis L. and produced with supercritical carbon dioxide. The bark is washed and oven dried to reduce moisture content before being crushed and extracted with supercritical carbon dioxide. The extract is dissolved in medical-grade ethanol and re-crystallised to yield magnolia bark extract. Magnolia bark extract is mainly composed of two phenolic compounds, magnolol and honokiol. Appearance: Light brownish powder Purity: Magnolik ≥ 0,5 % Matze-germ oil high in unsaponifiable matter is produced by vacuum distilition and it is different from refined maize-germ oil in the concentration of the unsaponifiable matter		Changes to legislation: There are currently no known outstanding effects for the	
Extract Magnolia bark extract is obtained from the bark of the plant Magnolia officinalis L. and produced with supercritical carbon dioxide. The bark is washed and oven dried to reduce moisture content before being crushed and extracted with supercritical carbon dioxide. The extract is dissolved in medical-grade ethanol and re-crystallised to yield magnolia bark extract. Magnolia bark extract is mainly composed of two phenolic compounds, magnolol and honokiol. Appearance: Light brownish powder Purity: Magnolia E85,2 % Honokiol: ≥ 0,5 % Magnolol & Honokiol: ≥ 94 % Total Eudesmol: ≤ 2 % Moisture: 0,50 % Magnoli (ppm): ≤ 0,5 Leady metals: Arsenic (ppm): ≤ 0,5 Lead (ppm): ≤ 10 Tubocurarine (ppm): ≤ 2,0 Total Alkaloid (ppm): ≤ 100 Maize-germ oil high in unsaponifiable matter is produced by vacuum distillation and it is different from refined maize-germ oil in the concentration of the unsaponifiable matter :> 9,0 g/100 g matter Description/Definition: Misue-germ oil high in untsaponifiable matter: > 9,0 g/100 g Tocopherol (%): 63,89 % %-tocopherol (%): 61,025 % Bytopherol (%): 64,89 % %-tocopherol (%): 61,025 % Stocopherol (%): 63,0 % %-tocopherol (%): 64,89 % Stocopherol (%): 67,0 % %-tocopherol (%): 63,0 % Storobyl condes: > 6,5 g/100 g Totals in triglycerides: 4		Appearance (20% aqueous solution): Clear solution pH (20% aqueous solution): Approx. 6,0 Impurities: Chloride: $\leq 0,05\%$ Sulphate: $\leq 0,05\%$ Arsenic: $\leq 3,0$ ppm Lead: $\leq 2,0$ ppm Cadmium: ≤ 1 ppm	
Maize-germ oil high in unsaponifiable matterDescription/Definition: Maize-germ oil high in unsaponifiable matter is produced by vacuum distillation and it is different from refined maize-germ oil in the concentration of the unsaponifiable fraction (1,2 g in refined maize-germ oil and 10 g in 'maize-germ oil high in unsaponifiable matter'). Purity: Unsaponifiable matter: > 9,0 g/100 g Tocopherols: ≥ 1,3 g/100 g α-tocopherol (%): 10-25 % β-tocopherol (%): < 3,0 % γ-tocopherol (%): < 53,0 % Sterols, triterpenic alcohols, methylsterols: > 6,5 g/100 g Fatty acids in triglycerides:aCommission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in	-	Description/Definition: Magnolia bark extract is obtained from the bark of the plant <i>Magnolia</i> <i>officinalis</i> L. and produced with supercritical carbon dioxide. The bark is washed and oven dried to reduce moisture content before being crushed and extracted with supercritical carbon dioxide. The extract is dissolved in medical-grade ethanol and re-crystallised to yield magnolia bark extract. Magnolia bark extract is mainly composed of two phenolic compounds, magnolol and honokiol. Appearance: Light brownish powder Purity: Magnolol: $\geq 85,2 \%$ Honokiol: $\geq 0,5 \%$ Magnolol & Honokiol: $\geq 94 \%$ Total Eudesmol: $\leq 2 \%$ Moisture: 0,50 % Heavy metals: Arsenic (ppm): $\leq 0,5$ Lead (ppm): $\leq 0,5$ Methyl eugenol (ppm): ≤ 10 Tubocurarine (ppm): $\leq 2,0$	
	oil high in unsaponifiable	Description/Definition: Maize-germ oil high in unsaponifiable matter is produced by vacuum distillation and it is different from refined maize-germ oil in the concentration of the unsaponifiable fraction (1,2 g in refined maize-germ oil and 10 g in 'maize-germ oil high in unsaponifiable matter'). Purity: Unsaponifiable matter: > 9,0 g/100 g Tocopherols: $\geq 1,3$ g/100 g α -tocopherol (%): 10-25 % β -tocopherol (%): < 3,0 % γ -tocopherol (%): < 3,0 % Sterols, triterpenic alcohols, methylsterols: > 6,5 g/100 g	
p. 1).	Annexes II and III	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in	

b Commission implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

	palmitic acid: 10,0-20,0 % stearic acid: $< 3,3 \%$ oleic acid: 20,0-42,2 % linoleic acid: 34,0-65,6 % linolenic acid: $< 2,0 \%$ Acid value: $\le 6,0 \text{ mg KOH/g}$ Peroxide value (PV): $\le 10 \text{ mEq O}_2/\text{kg}$ Heavy metals: Iron (Fe): $< 1500 \mu\text{g/kg}$ Copper (Cu): $< 100 \mu\text{g/kg}$ Impurities:
	Polycyclic aromatic hydrocarbons (PAH) Benzo(a)pyrene: $< 2 \mu g/kg$ Treatment with active carbon is required to ensure that polycyclic aromatic hydrocarbons (PAH) are not enriched in the production of 'maize-germ oil high in unsaponifiable matter'
Methylcellulose	Description/Definition: Methyl cellulose is cellulose obtained directly from natural strains of fibrous plant material and partially etherified with methyl groups. Chemical name: Methyl ether of cellulose Chemical formula: The polymers contain substituted anhydroglucose units with the following general formula: C6H7O2(OR1)(OR2)(OR3) where R1, R2, R3 each may be one of the following: — H — CH ₃ or — CH ₂ CH ₃ Molecular weight: Macromolecules: from about 20 000 (n about 100) up to about 380 000 g/mol (n about 2 000) Assay: Content not less than 25 % and not more than 33 % of methoxyl groups (-OCH ₃) and not more than 5 % of hydroxyethoxyl groups (- OCH ₂ CH ₂ OH) Slightly hygroscopic white or slightly yellowish or greyish odourless and tasteless, granular or fibrous powder. Solubility: Swelling in water, producing a clear to opalescent, viscous, colloidal solution. Insoluble in ethanol, ether and chloroform. Soluble in glacial acetic acid. Purity: Loss on drying: ≤ 10 % (105 °C, 3 hours) Sulphated Ash: ≤ 1,5 % determined at 800 ± 25 °C pH: ≥ 5,0 and ≤ 8,0 (1 % colloidal solution) Heavy metals: Arsenic: ≤ 3,0 mg/kg Lead: ≤ 2,0 mg/kg Mercury: ≤ 1,0 mg/kg Tation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in

a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

(6S)-5-	Description/Definition:		
methyltetrahydr	fölia mical name: N-[4-[[[(6S)-2-amino-1,4,5,6,7,8-hexahydro-5-methyl-4-		
acid, oxo-6-pteridinyl]methyl]amino]benzoyl]-L-glutamic acid, glucosa			
glucosamine salt			
salt	Chemical formula: C ₃₂ H ₅₁ N ₉ O ₁₆		
	Molecular weight: 817,80 g/mol (anhydrous)		
	CAS No.: 1181972-37-1		
	Appearance: Creamy to light-brown powder		
	Purity:		
	Diastereoisomeric purity: At least 99 % of (6S)-5-methyltetrahydrofolic acid		
	Glucosamine assay: 34-46 % in dry basis		
	5-Methyltetrahydrofolic acid assay: 54-59 % in dry basis		
	Water: $\leq 8,0\%$		
	Heavy metals:		
	Lead: $\leq 2,0$ ppm		
	Cadmium: $\leq 1,0$ ppm		
	Mercury: $\leq 0,1$ ppm		
	Arsenic: $\leq 2,0$ ppm		
	Boron: $\leq 10 \text{ ppm}$		
	Microbiological criteria:		
	Total aerobic microbial count: $\leq 100 \text{ CFU/g}$		
	Yeasts and moulds: $\leq 100 \text{ CFU/g}$		
	Escherichia coli: Absence in 10g		
(Organic Silicon)	Chemical name: Silanetriol, 1-methyl- Chemical formula: CH_6O_3Si Molecular weight: 94,14 g/mol CAS No: 2445-53-6 Purity: Organic Silicon (monomethylsilanetriol) preparation (aqueous solution): Acidity (pH): 6,4-6,8 Silicon: 100-150 mg Si/l Heavy metals: Lead: $\leq 1,0 \mu g/l$ Mercury: $\leq 1,0 \mu g/l$ Cadmium: $\leq 1,0 \mu g/l$ Arsenic: $\leq 3,0 \mu g/l$ Solvents: Methanol: $\leq 5,0 mg/kg$ (residual presence)		
Musslial autroat			
Mycelial extract from Shiitake mushroom (<i>Lentinula</i> <i>edodes</i>)	Description/Definition: The novel food ingredient is a sterile aqueous extract obtained from the mycelium of <i>Lentinula edodes</i> cultivated in a submerged fermentation. It is a light brown, slightly turbid liquid.		
a Commission Regul	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,		
to the import of gu			
-			

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Lentinan is a β -(1-3) β -(1-6)-D-glucan which has a molecular weight of approximately 5 × 10 ⁵ Daltons, a degree of branching of 2/5 and a triple helical tertiary structure. Purity/Composition of the mycelial extract from <i>Lentinula edodes</i> : Moisture: 98 % Dry matter: 2 % Free glucose: < 20 mg/ml Total protein(¹): < 0,1 mg/ml N-containing constituents(²): < 10 mg/ml Lentinan: 0,8 – 1,2 mg/ml\ (¹) Bradford method (²) Kjeldahl method
Noni fruit juice (<i>Morinda</i> <i>citrifolia</i>)	Description/Definition: Noni fruits (fruits of <i>Morinda citrifolia</i> L.) are pressed. The obtained juice is pasteurised. An optional fermentation step before or after the pressing may occur. Rubiadin: $\leq 10 \ \mu g/kg$ Lucidin: $\leq 10 \ \mu g/kg$
Noni fruit juice powder (<i>Morinda</i> <i>citrifolia</i>)	Description/Definition: Seeds and skin of the sun-dried fruits of <i>Morinda citrifolia</i> are separated. The obtained pulp is filtered to separate juice from the flesh. Desiccation of the produced juice occurs in one or two ways: Either by atomisation using maize maltodextrins, this mixture is obtained by keeping the rates of inflow of the juice and maltodextrins constant Or by zeodratation or drying and then mixing with an excipient, this process allows the juice to be dried initially and then mixed with maltodextrins (same amount as used in atomisation).
Noni fruit puree and concentrate (<i>Morinda</i> <i>citrifolia</i>)	Description/Definition: The fruits of <i>Morinda citrifolia</i> are harvested by hand. Seeds and skin may be separated mechanically from the pureed fruits. After pasteurisation, the puree is packaged in aseptic containers and stored under cold conditions. <i>Morinda citrifolia</i> concentrate is prepared from <i>M. citrifolia</i> puree by treatment with pectinolytic enzymes ($50-60 \degree C$ for 1-2 h). Then the puree is heated to inactivate the pectinases and then immediately cooled. The juice is separated in a decanter centrifuge. Afterwards the juice is collected and pasteurised, prior to being concentrated in a vacuum evaporator from a brix of 6 to 8 to a brix of 49 to 51 in the final concentrate. Composition: Puree: Moisture: $89-93 \%$ Protein: $< 0,6 \text{ g}/100 \text{ g}$

p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

		Fat: $\leq 0,4 \text{ g}/100 \text{ g}$ Ash: $< 1,0 \text{ g}/100 \text{ g}$ Total carbohydrates: 5-10 g/100 g Fructose: 0,5-3,82 g/100 g Glucose: 0,5-3,14 g/100 g Dietary fibre: $< 0,5-3 \text{ g}/100 \text{ g}$ 5,15-dimethylmorindol (1): $\leq 0,254 \mu\text{g/ml}$ Lucidin (1): Not detectable Alizarin (1): Not detectable Rubiadin (1): Not detectable Rubiadin (1): Not detectable Concentrate: Moisture: 48-53 % Protein: 3-3,5 g/100 g Fat: $< 0,04 \text{ g}/100 \text{ g}$ Ash: 4,5-5,0 g/100 g Total carbohydrates: 37-45 g/100 g Fructose: 9-11 g/100 g Glucose: 9-11 g/100 g Dietary fibre: 1,5-5,0 g/100 g 5,15-dimethylmorindol (¹): $\leq 0,254 \mu\text{g/ml}$ (¹) By an HPLC-UV method developed and validated for the analysis of anthraquinones in Morinda citrifolia puree and concentrate. Limits of detection: 2,5 ng/ml (5,15 dimethylmorindol); 50,0 ng/ml (lucidin); 6,3 ng/ml (alizarin) and 62 S ng/ml (mukiadin)	
(Ma	ni leaves orinda ifolia)	and 62,5 ng/ml (rubiadin). Description/Definition: After cutting, the leaves of <i>Morinda citrifolia</i> are subject to drying and roasting steps. The product has a particle size ranging from broken leaves to coarse powder with fines. It is of greenish brown to brown colour. Purity/Composition: Moisture: $< 5,2 \%$ Protein: 17- 20 % Carbohydrate: 55-65 % Ash: 10-13 % Fat: 4-9 % Oxalic acid: $< 0,14 \%$ Tannic acid: $< 2,7 \%$ 5,15-dimethylmorindol: $< 47 \text{ mg/kg}$ Rubiadin: non detectable, $\le 10 \mu\text{g/kg}$	
pow (Ma citri	ni fruit vder o <i>rinda</i> ifolia)	Description/Definition: Noni fruit powder is made from pulped noni (<i>Morinda citrifolia L.</i>) fruits by freeze-drying. Fruits are pulped and seeds are removed. After freeze- drying during which water is removed from noni fruits, the remaining noni pulp is milled to a powder and encapsulated. ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in	
b	Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).		

	Purity/CompositionMoisture: $5,3-9 \%$ Protein: $3,8-4,8 g/100 g$ Fat: $1-2 g/100 g$ Ash: $4,6-5,7 g/100 g$ Total carbohydrates: $80-85 g/100 g$ Fructose: $20,4-22,5 g/100 g$ Glucose: $22-25 g/100 g$ Dietary fibre: $15,4-24,5 g/100 g$ $5,15$ -dimethylmorindol (1): $\leq 2,0 \mu g/ml$ (1)By an HPLC-UV method developed and validated for the analysis of anthraquinones in Morinda citrifolia fruit powder. Limits of detection: $2,5 ng/ml$ ($5,15$ dimethylmorindol)
<i>Odontella aurita</i> microalgae	Silicon: 3,3 % Crystalline silica: max 0,1-0,3 % as impurity
Oil extracted from squidsAcid value: ≤ 0.5 KOH/g oil Peroxide value (PV): ≤ 5 meq O2/kg oil p-Anisidine value: ≤ 20 Cold test at 0 °C: ≤ 3 hours Moisture: ≤ 0.1 % (w/w)	
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Unsaponifiable ma Trans fatty acids: Docosahexaeonic Eicosapentaenoic	$\leq 1,0\%$ acid: $\geq 20\%$	
Pasteurised	Parameter	Target	Comments
fruit-based preparations produced using high-pressure treatment	Fruit storage before high- pressure treatment	Minimum 15 days at – 20 °C	Fruit harvested and stored in conjunction with good/hygienic agricultural and manufacturing practices
treatment	Fruit added	40 % to 60 % of thawed fruit	Fruit homogenised and added to other ingredients
	pН	3,2 to 4,2	
	° Brix	7 to 42	Assured by added sugars
	a _w	< 0,95	Assured by added sugars
	Final storage	60 days maximum	EquisalEntraxistomage regimen for conventionally processed product
maize starch	Description/Definition: Phosphated maize starch (phosphated distarch phosphate) is a chemically modified resistant starch derived from high amylose starch by combining chemical treatments to create phosphate cross-links between carbohydrate residues and esterified hydroxyl groups. The novel food ingredient is a white or nearly white powder. CAS No: 11120-02-8 Chemical formula: $(C_6H_{10}O_5)_n [(C_6H_9O_5)_2PO_2H]x [(C_6H_9O_5)PO_3H_2]y$ n = number of glucose units; x, y = degrees of substitution The chemical characteristics of phosphated distarch phosphate: Loss on drying: 10-14 % pH: 4,5-7,5 Dietary fibre: ≥ 70 % Starch: 7-14 % Protein: $\leq 0,8$ % Lipids: $\leq 0,8$ % Residual bound phosphorus: $\leq 0,4$ % (as phosphorus) 'high amylose maize' as source		
	nDescription/Defin		
from fish phospholipids	The novel food ingredient is yellow to brown powder. Phosphatidylserine is obtained from fish phospholipids by an enzymatic transphosphorylation with the amino acid L-serine. Specification of the phosphatidylserine product manufactured from fish phospholipids: Moisture: $< 5,0 \%$ Phospholipids: $\geq 75 \%$		
	Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012		
 b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10). 			

Phosphatidylser from soya phospholipids	Phosphatidylserine: $\geq 35 \%$ Glycerides: $< 4,0 \%$ Free L-serine: $< 1,0 \%$ Tocopherols: $< 0,5 \%$ (¹) Peroxide value (PV): $< 5,0 \text{ meq } O_2/kg$ (¹) Tocopherols may be added as antioxidants according to Commission Regulation (EU) No 1129/2011 nDescription/Definition: The novel food ingredient is off-white to light yellow powder. It is also available in liquid form with a clear brown to orange colour. The liquid form contains medium chain triacylglycerides (MCT) as a carrier. It contains lower levels of Phosphatidylserine due to the fact that it includes significant amounts of oil (MCT). Phosphatidylserine from soya phospholipids is obtained through enzymatic transphosphatidylation of high-phosphatidylcholine soybean lecithin with the amino acid L-serine. Phosphatidylserine consists of a glycerophosphate skeleton conjugated with two fatty acids and L-serine via a phosphodiester linkage. Characteristics of Phosphatidylserine from soya phospholipids: Powder form: Moisture: $< 2,0 \%$			
	Phospholipids: $\geq 85 \%$ Phosphatidylserine: $\geq 61 \%$			
	Glycerides: < 2,0 % free L-serine: < 1,0 %			
	Tocopherols: < 0,3 % Phytosterols: < 0,2 %			
	Liquid form:			
	Moisture: $< 2,0 \%$			
	Phospholipids: $\geq 25 \%$ Phosphatidylserine: $\geq 20 \%$			
	Glycerides: not applicable			
	free L-serine: < 1,0 %			
	Tocopherols: $< 0,3 \%$			
	Phytosterols: < 0,2 %			
Phospholipid Description/Definition:				
product	The product is manufactured through enzymatic conversion of soy			
containing	lecithin. The phospholipid product is a highly concentrated, yellow-			
equal	brown powder form of phosphatidylserine and phosphatidic acid at an			
amounts of equal level. phosphatidylserinspecification of the product:				
and Moisture: $\leq 2,0\%$				
phosphatidic	Total phospholipids: $\geq 70 \%$			
acid	Phosphatidylserine: $\geq 20 \%$			
	Phosphatidic acid: ≥ 20 %			
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,			
to the import of g	b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).			

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Glycerides: $\leq 1,0 \%$ Free L-serine: $\leq 1,0 \%$ Tocopherols: $\leq 0,3 \%$ Phytosterols: $\leq 2,0 \%$ Silicon dioxide is used with a maximum content of 1,0 %
Phospholipides from egg yolk	85 % and 100 % pure Phospholipides from egg yolk
Phytoglycogen	Description: White to off-white powder which is an odourless, colourless, flavourless polysaccharide derived from non-GM sweet corn using conventional food processing techniques Definition: Glucose polymer ($C_6H_{12}O_6$)n with linear linkages of $\alpha(1 - 4)$ glycosidic bonds branched every 8 to 12 glucose units by $\alpha(1 - 6)$ glycosidic bonds Specifications: Carbohydrates: 97 % Sugars: 0,5 % Fibre: 0,8 % Fat: 0,2 % Protein: 0,6 %
Phytosterols/ phytostanols	Description/Definition: Phytosterols and phytostanols are sterols and stanols that are extracted from plants and may be presented as free sterols and stanols or esterified with food grade fatty acids. Composition (with GC-FID or equivalent method): β -sitosterol: < 81 % β -sitostanol: < 35 % campesterol: < 40 % campestanol: < 15 % stigmasterol: < 30 % brassicasterol: < 3,0 % other sterols/stanols: < 3,0 % Contamination/Purity (GC-FID or equivalent method): Phytosterols and phytostanols extracted from sources other than vegetable oil suitable for food have to be free of contaminants, best ensured by a purity of more than 99 % of the phytosterol/phytostanol ingredient.
Plum kernel oil	Description/Definition: Plum kernel oil is a vegetable oil obtained by cold pressing of plum (<i>Prunus domestica</i>) kernels. Composition: Oleic acid (C18:1): 68 % Linoleic acid (C18:2): 23 % γ -Tocopherol:80 % of total tocopherols β -Sitosterol: 80-90 % of total sterols Triolein: 40-55 % of triglycerides
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).

Cyanhydric acid: maximum 5 mg/kg oil

	substance: $\geq 800 \text{ mg/g}$
and Ash: hydrolysates Glyc thereof Lysi	ein (N * 6,25): $\geq 600 \text{ mg/g}$ (dry substance) $\leq 400 \text{ mg/g}$ (dry substance) coalkaloid (total): $\leq 150 \text{ mg/kg}$ noalanine (total): $\leq 500 \text{ mg/kg}$ noalanine (free): $\leq 10 \text{ mg/kg}$
oligopeptidase (enzyme preparation)Syste Synd ended Molde Enzy CASS Sour Desc conta Spect Activ App Colde batel Dry Glut Hea 	Effication of the enzyme: ematic name: Prolyl oligopeptidase onyms: Prolyl endopeptidase, proline-specific endopeptidase, prolylpeptidase ecular weight: 66 kDa /me Commission number: EC 3.4.21.26 6 number: 72162-84-6 ce: A genetically modified strain of <i>Aspergillus niger</i> (GEP-44) cription: Prolyl oligopeptidase is available as an enzyme preparation aining approximately 30 % maltodextrin. crifications of the enzyme preparation of prolyl oligopeptidase: vity: > 580 000 PPI(¹ /g (> 34,8 PPU(²)/g) earance: Microgranulate vur: Off-white to orange yellowish. The colour may change from h to batch Matter: > 94 % en: < 20 ppm vy metals: L: $\leq 1,0$ mg/kg mic: $\leq 1,0$ mg/kg mic: $\leq 1,0$ mg/kg cury: $\leq 0,1$ mg/kg robiological criteria: I aerobic plate count: $\leq 10^3$ CFU/g I yeasts and moulds: $\leq 10^2$ CFU/g hite reducing anaerobes: ≤ 30 CFU/g <i>robacteriaceae</i> : < 10 CFU/g <i>notella</i> : Absence in 25 g <i>terichia coli</i> : Absence in 25 g <i>terichia coli</i> : Absence in 10 g <i>domonas aeruginosa</i> : Absence in 10 g <i>dria monocytogenes</i> : Absence in 10 g <i>dria monocytogenes</i> : Absence in 25 g microbial activity: Absent otoxins: Below limits of detection: Aflatoxin B1, B2, G1, G2 25 µg/kg), total Aflatoxins (< 2,0 µg/kg), Ochratoxin A (< 0,20 µg/ T-2 Toxin (< 5 µg/kg), Zearalenone (< 2,5 µg/kg), Fumonisin B1 and < 2,5 µg/kg) PPI – Protease Picomole International

a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

$(^{2})$	PPU – Prolyl Peptidase Units or Proline Protease Units
----------	--

Protein extract from pig kidneys	Description/Definition: The protein extract is obtained from homogenised pig kidneys through a combination of salt precipitation and high speed centrifugation. The obtained precipitate contains essentially proteins with 7 % of the enzyme diamine oxidase (enzyme nomenclature E.C. 1.4.3.22) and is resuspended in a physiologic buffer system. The obtained pig kidney extract is formulated as encapsulated enteric coated pellets to reach the active sites of digestion. Basic Product: Specification: pig kidney protein excerpt with natural content of Diamin oxidase (DAO): Physical condition: liquid Colour: brownish Appearance: slightly turbid solution pH value: 6,4-6,8 Enzymatic activity: > 2 677 kHDU DAO/ml (DAO REA (DAO Radioextractionassay)) Microbiological criteria: Brachyspira spp.: negative (Real Time PCR) <i>Listeria monocytogenes</i> : negative (Real Time PCR) <i>Listeria monocytogenes</i> : negative (Real Time PCR) <i>Staphylococcus aureus</i> : < 100 CFU/g Influenza A: negative (Reverse Transcription Real Time PCR) <i>Escherichia coli</i> : < 10 CFU/g Total aerobic microbiological count: < 10 ⁵ CFU/g Yeasts/moulds count: < 10 ⁵ CFU/g Salmonella: Absence/10g Bile salt resistant enterobacteriaceae: < 10 ⁴ CFU/g Final product: Specification pig kidney protein excerpt with natural content of DAO (E.C. 1.4.3.22) in an enteric coated formulation: Physical condition: solid Colour: yellow gray Appearance: micropellets Enzymatic activity: 110-220 kHDU DAO/g pellet (DAO REA (DAO Radioextractionassay)) Acid stability 15 min 0,1M HCl followed by 60 min Borat pH = 9,0: > 68 KHDU DAO'g pellet (DAO REA (DAO Radioextractionassay)) Humidity: < 10 % Staphylococcus aureus: < 100 CFU/g
	<i>Staphylococcus aureus</i> : < 100 CFU/g
	<i>Escherichia coli</i> : < 10 CFU/g
	Total aerobic microbiological count: $< 10^4$ CFU/g
	Total combined yeasts/moulds count: < 10 ³ CFU/g Salmonella: Absence/10g
	Bile salt resistant enterobacteriaceae: $< 10^2$ CFU/g

Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Rapeseed	Description/Definition:					
oil high in	Rapeseed oil high in unsaponifiable matter' is produced by vacuum					
oil high in unsaponifiable matter	distillation and it is different from refined rapeseed oil in the concentration of the unsaponifiable fraction (1 g in refined rapeseed oil and 9 g in 'rapeseed oil high in unsaponifiable matter'). There is a minor reduction of triglycerides containing monounsaturated and polyunsaturated fatty acids. Purity: Unsaponifiable matter: > 7,0 g/100 g Tocopherols: > 0,8 g/100 g α -tocopherol (%): 30-50 % γ -tocopherol (%): 50-70 % δ -tocopherol (%): 50-70 % δ -tocopherol (%): < 6,0 % Sterols, triterpenic alcohols, methylsterols: > 5,0 g/100 g Fatty acids in triglycerides: palmitic acid: 3-8 % stearic acid: 0,8-2,5 % oleic acid: 50-70 % linolenic acid: 6-14 % erucic acid: < 2,0 % Acid value: \leq 6,0 mg KOH/g Peroxide value (PV): \leq 10 mEq O ₂ /kg Heavy metals:					
	Iron (Fe): < 1 000 μg/kg Copper (Cu): < 100 μg/kg Impurities: Polycyclic aromatic hydrocarbons (PAH) Benzo(a)pyrene: < 2 μg/kg Treatment with active carbon is required to ensure that polycyclic aromatic hydrocarbons (PAH) are not enriched in the production of 'rapeseed oil high in unsaponifiable matter.					
Rapeseed Protein	Definition: Rapeseed protein is an aqueous protein-rich extract from rapeseed press cake originating from non-genetically modified <i>Brassica napus</i> L. and					
	Brassica rapa L. Description: White to off-white, spray dried powder Total protein: $\geq 90 \%$ Soluble protein: $\geq 85 \%$ Moisture: $\leq 7,0 \%$ Carbohydrates: $\leq 7,0 \%$ Fat: $\leq 2,0 \%$ Ash: $\leq 4,0 \%$ Fibre: $\leq 0,5 \%$ Total glucosinolates: $\leq 1 \text{ mmol/kg}$ Purity:					

Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Total phytate: $\leq 1,5 \%$ Lead: $\leq 0,5 \text{ mg/kg}$ Microbiological criteria: Yeast and mould count: $\leq 100 \text{ CFU/g}$ Aerobic bacteria count: $\leq 10 000 \text{ CFU/g}$ Total coliform count: $\leq 10 \text{ CFU/g}$ <i>Escherichia coli:</i> Absence in 10 g <i>Salmonella</i> : Absence in 25 g				
Trans- resveratrol	Description/Definition:Synthetic Trans-resveratrol is off-white to beige crystals.Chemical name: 5-[(E)-2-(4-hydroxyphenyl)ethenyl]benzene-1,3-diolChemical formula: $C_{14}H_{12}O_3$ Molecular weight: 228,25 DaCAS No: 501-36-0Purity:Trans-resveratrol: ≥ 98 %-99 %Total by-products (related substances): $\leq 0,5$ %Any single related substance: $\leq 0,1$ %Sulphated ash: $\leq 0,1$ %Loss on drying: $\leq 0,5$ %Heavy metals:Lead: $\leq 1,0$ ppmMercury: $\leq 0,1$ ppmArsenic: $\leq 1,0$ ppmImpurities:Diisopropylamine: ≤ 50 mg/kgMicrobial source: A genetically modified strain of SaccharomycescerevisiaeAppearance: Off-white to slight yellow powderParticle size: 100 % less than 62,23 µmTrans-resveratrol content: Min. 98 % w/w (dry weight basis)Ash: Max. 0,5 % w/wMoisture: Max. 3 % w/w				
Rooster comb extract	Description/Definition:Rooster comb extract is obtained from Gallus gallus by enzymatichydrolysis of rooster comb and by subsequent filtration, concentrationand precipitation steps. The principal constituents of rooster comb extractare the glycosaminoglycans hyaluronic acid, chondroitin sulphate Aand dermatan sulphate (chondroitin sulphate B). White or almost whitehygroscopic powder.Hyaluronic acid: 60-80 %Chondroitin sulphate A: $\leq 5,0$ %Dermatan sulphate (chondroitin sulphate B): ≤ 25 %pH: 5,0-8,5Purity:Chlorides: $\leq 1,0$ %				
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,				
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).				

	Nitrogen: $\leq 8,0 \%$ Loss on drying: (105 °C for 6 hours): $\leq 10 \%$ Heavy metals: Mercury: $\leq 0,1 \text{ mg/kg}$ Arsenic: $\leq 1,0 \text{ mg/kg}$ Cadmium: $\leq 1,0 \text{ mg/kg}$ Chromium: $\leq 10 \text{ mg/kg}$ Lead: $\leq 0,5 \text{ mg/kg}$ Microbiological criteria: Total viable aerobic count: $\leq 10^2 \text{ CFU/g}$ <i>Escherichia coli</i> : Absence in 1 g <i>Salmonella</i> : Absence in 1 g <i>Staphylococcus aureus</i> : Absence in 1 g <i>Pseudomonas aeruginosa</i> : Absence in 1g
Sacha Inchi oil from <i>Plukenetia</i> <i>volubilis</i>	Description/Definition: Sacha inchi oil is a 100 % cold pressed vegetable oil obtained from the seeds of <i>Plukenetia volubiis</i> L. It is a transparent, fluid (liquid) and shiny oil at room temperature. It has a fruity, light, green vegetable taste without undesirable flavours. Aspect, limpidity, shine, colour: Fluid at room temperature, clean, shiny yellow gold Odour and taste: Fruity, vegetable without non acceptable taste or odour Purity: Water and Volatiles: < 0,2 g/100 g Impurities insoluble in hexane: < 0,05 g/100 g Oleic acidity: < 2,0 g/100 g Peroxide value (PV): < 15 meq O ₂ /kg Trans fatty acids: < 1,0 g/100 g Total unsaturated fatty acids: > 90 % Omega 3 alpha linolenic acid (ALA): > 45 % Saturated fatty acids: < 10 % No trans fatty acids (< 0,5 %) No erucic acid (< 0,2 %) More than 50 % of tri-linolenin and di-linolenin-triglycerides Phytosterols composition and level No cholesterol (< 5,0 mg/100 g)
Salatrims	Description/Definition: Salatrim is the internationally recognised acronym for (short and long chain acyl triglyceride molecules). Salatrim is prepared by non-enzymatic inter-esterification of triacetin, tripropionin, tributyrin, or their mixtures with hydrogenated canola, soybean, cottonseed, or sunflower oil. Description: Clear, slightly amber liquid to a light coloured waxy solid at room temperature. Free of particulate matter and of foreign or rancid odour. Glycerol ester disribution: Triacylglycerols: > 87 %
Annexes II and III p. 1). b Commission Imple	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, menting Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Diacylglycerols: $\leq 10 \%$ Monoacylglycerols: $\leq 2,0 \%$ Fatty acid composition: MOLE % LCFA (long chain fatty acids): 33-70 % MOLE % SCFA (short chain fatty acids): 30-67 % Saturated long chain fatty acids: $< 70 \%$ by weight Trans fatty acids: $\leq 1,0 \%$ Free fatty acids as oleic acid: $\leq 0,5 \%$ Triacylglycerol profile: Triesters (short/long of 0,5 to 2,0): $\geq 90 \%$ Triesters (short/long = 0): $\leq 10 \%$ Unsaponifiable material: $\leq 1,0 \%$ Moisture: $\leq 0,3 \%$ Ash: $\leq 0,1 \%$ Colour: $\leq 3,5 \text{ Red (Lovibond)}$ Peroxide value (PV): $\leq 2,0 \text{ Meq/Kg}$
<i>Schizochytrium</i> <i>sp</i> . oil rich in DHA and EPA	Acid value: $\leq 0,5$ mg KOH/g Peroxide value (PV): $\leq 5,0$ meq/kg oil Oxidative stability: All food products containing <i>Schizochytrium sp.</i> oil rich in DHA and EPA should demonstrate oxidative stability by appropriate and recognised national/international test methodology (e.g. AOAC) Moisture and volatiles: $\leq 0,05$ % Unsaponifiables: $\leq 4,5$ % Trans-fatty acids: ≤ 1 % DHA content: $\geq 22,5$ % EPA content: ≥ 10 %
<i>Schizochytrium</i> sp. (ATCC PTA-9695) oil	Peroxide value (PV): \leq 5,0 meq/kg oil Unsaponifiables: \leq 3,5 % Trans-fatty acids: \leq 2,0 % Free fatty acids: \leq 0,4 % Docosapentaenoic acid (DPA) n-6: \leq 7,5 % DHA content: \geq 35 %
<i>Schizochytrium</i> sp. oil	Acid value: $\leq 0.5 \text{ mg KOH/g}$ Peroxide value (PV): $\leq 5.0 \text{ meq/kg oil}$ Moisture and volatiles: $\leq 0.05 \%$ Unsaponifiables: $\leq 4.5 \%$ Trans-fatty acids: $\leq 1.0 \%$ DHA content: $\geq 32.0 \%$
<i>Schizochytrium</i> sp. (T18) oil	Acid value: $\leq 0.5 \text{ mg KOH/g}$ Peroxide value (PV): $\leq 5.0 \text{ meq/kg oil}$ Moisture and volatiles: $\leq 0.05 \%$ Unsaponifiables: $\leq 3.5 \%$ Trans-fatty acids: $\leq 2.0 \%$ Free fatty acids: $\leq 0.4 \%$
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

DHA content:	\geq	35	%	
--------------	--------	----	---	--

Fermented	Description/Definition:
Fermented soybean extract	Description/Definition: Fermented soybean extract is an odourless milk-white coloured powder. It is comprised of 30 % fermented soybean extract powder and 70 % resistant dextrin (as carrier) from corn-starch, which is added during the processing. Vitamin K ₂ is removed during the manufacturing process. Fermented soybean extract contains nattokinase isolated from natto, a foodstuff produced by the fermentation of non-genetically modified soybeans (<i>Glycine max</i> (L.)) with a selected strain of <i>Bacillus subtilis</i> var. natto. Nattokinase activity: 20 000 -28 000 Fibrin degradation unit/g(¹) Identity: Confirmable Condition: No offensive taste or smell Loss on drying: $\leq 10 \%$
	Vitamin K_2 : $\leq 0,1 \text{ mg/kg}$
	Heavy metals: Lead: $\leq 5,0$ mg/kg Arsenic: $\leq 3,0$ mg/kg
	Microbiological criteria: Total viable aerobic count: $\leq 10^3$ CFU(³)/g
	Yeast and mould: $\leq 10^2$ CFU/g Coliforms: ≤ 30 CFU/g Spore-forming bacteria: ≤ 10 CFU/g <i>Escherichia coli</i> : Absence/25 g <i>Salmonella</i> : Absence/25 g <i>Listeria</i> : Absence/25 g
	$(^1)$ Assay method as described by Takaoka et al. (2010).
Spermidine- rich wheat germ extract (<i>Triticum</i> <i>aestivum</i>)	Description/Definition: Spermidine-rich wheat germ extract is obtained from non-fermented, non-sprouting wheat germs (<i>Triticum aestivum</i>) by the process of solid- liquid extraction targeting specifically, but not exclusively polyamines. Spermidine: 0,8-2,4 mg/g Spermidine trichloride < 0,1 μ g/g Putrescine: < 0,3 mg/g Cadaverine: < 0,3 mg/g Cadaverine: < 0,1 μ g/g Mycotoxins: Aflatoxins (total): < 0,4 μ g/kg Microbiological criteria: Total aerobic bacteria: < 10 000 CFU/g Yeast and moulds: < 100 CFU/g <i>Escherichia coli:</i> < 10 CFU/g

a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Sucromalt	Description/Definition: Sucromalt is a complex mixture of saccharides which is produced from sucrose and a starch hydrolysate by means of an enzymatic reaction. In this process, glucose units are attached to saccharides from the starch hydrolysate by means of an enzyme produced by the bacterium <i>Leuconostoc citreum</i> or by means of a recombinant strain of the production organism <i>Bacillus licheniformis</i> . The resulting oligosaccharides are characterised by the presence of α -(1 \rightarrow 6) and α - (1 \rightarrow 3) glycosidic compounds. The overall product is syrup, in addition to these oligosaccharides, contains mainly fructose but also the disaccharide leucrose and other disaccharides. Total solids: 75-80 % Moisture: 20-25 % Sulphatase: Max 0,05 % pH: 3,5-6,0 Conductivity < 200 (30 %)
	Nitrogen < 10 ppm Fructose: 35-45 % d.w. Leucrose: 7-15 % d.w. Other disaccharides: Max 3 % Higher saccharides: 40-60 % d.w
Sugar cane fibre	Description/Definition:Sugar Cane Fibre is derived from the dry cell wall or fibrous residueremaining after expression or extraction of sugar juice from sugarcane, of the Saccharum genotype. It consists primarily of cellulose andhemicellulose.The production process consists of several steps, including:chipping, alkaline digestion, removal of lignins and other non-cellulosic components, bleaching of purified fibres, acid washing andneutralization.Moisture: $\leq 7,0 \%$ Ash: $\leq 0,3 \%$ Total Dietary Fibre (AOAC) dry basis (all insoluble): $\geq 95 \%$ of which: Hemicellulose (20-25 %) and cellulose (70-75 %)Silica (ppm): ≤ 200 Protein: $0,0 \%$ Fat: TracepH: 4-7Heavy metals:Mercury (ppm): $\leq 0,1$ Lead (ppm): $\leq 1,0$ Cadmium (ppm): $\leq 0,1$ Microbiological criteria:Yeast and moulds (CFU/g): $\leq 1 000$ Salmonella: Absence

Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Listeria monocytogenes: Absence

Sunflower oil extract	Description/Definition: The sunflower extract is obtained by a concentration factor of 10 of the unsaponifiable fraction of refined sunflower oil extracted from the seeds of the sunflower, <i>Helianthus Annuus</i> L. Composition: Oleic acid (C18:1): 20 % Linoleic acid (C18:2): 70 % Unsaponifiable matter: 8,0 % Phytosterols: 5,5 % Tocopherols: 1,1 %
Dried <i>Tetraselmis</i> <i>chuii</i> microalgae	Description/Definition: The dried product is obtained from the marine microalgae <i>Tetraselmis</i> <i>chuii</i> , belonging to the <i>Chlorodendraceae</i> family, cultivated in sterile sea water in closed photobioreactors insulated from the outside air. Purity/Composition: Identified by means of nuclear marker rDNA 18 S (sequence analysed no less than 1 600 base pairs) in the National Centre for Biotechnology information (NCBI) database: Not less than 99,9 % Humidity: $\leq 7,0 \%$ Proteins: 35-40 % Ashes: 14-16 % Carbohydrates: 30-32 % Fibre: 2-3 % Fat: 5-8 % Saturated fatty acids: 29-31 % of total fatty acids Monounsaturated fatty acids: 21-24 % of total fatty acids Polyunsaturated fatty acids: 44-49 % of total fatty acids Iodine: $\leq 15 \text{ mg/kg}$
<i>Therapon</i> <i>barcoo</i> /Scortum	Description/Definition: Scortum/ <i>Therapon barcoo</i> is a species of fish in the family Terapontidae It is an endemic fresh water species from Australia. It is now reared in fish farms. Taxonomic Identification: Class: Actinopterygii > order: Perciformes > family: Terapontidae > genus: <i>Therapon</i> or <i>Scortum barcoo</i> Composition of fish flesh: Protein (%): 18-25 Moisture (%): 65-75 Ash (%): 0,5-2,0 Energy (KJ/Kg): 6000-11500 Carbohydrates (%): 0,0 Fat (%): 5-15 Fatty acids (mg FA/g fillet): Σ PUFA n-3: 1,2-20,0 Σ PUFA n-6: 0,3-2,0 PUFA n-3/n-6: 1,5-15,0

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

D-Tagatose	Total omega 6 acids: 2,6-10,0 Description/Definition: Tagatose is produced by isomerization of galactose by means of chemica or enzymatic conversion, or by epimerization of fructose by means of enzymatic conversion. These are single-step conversions. Appearance: White or almost white crystals Chemical name: D-tagatose Synonym: D- <i>lyxo</i> -Hexulose CAS number: 87-81-0 Chemical formula: C ₆ H ₁₂ O ₆ Formula weight: 180,16 (g/mol) Purity: Assay: \geq 98 % on a dry weight basis Loss on drying: \leq 0,5 % (102 °C, 2 hours) Specific Rotation: $[\alpha]_D^{20}$: - 4 to - 5,6° (1 % aqueous solution)(¹)
	Melting range: 133–137 °C Heavy metals: Lead: ≤ 1,0 mg/kg(*)
	(*) Determine using an atomic absorption technique appropriate to the specified level. The selection of sample size and method of sample preparation may be based on the principles of the method described in FNP 5. 'Instrumental methods'(¹).
	(¹) Food and nutrition paper 5 Rev 2 – Guide to specifications for general notices, general analytical techniques, identification tests, test solutions and other reference materials (JECFA) 1991 307 p.; English – ISBN 92-5-102991-1
Taxifolin-rich extract	Description:Taxifolin-rich extract from the wood of Dahurian Larch (<i>Larix gmelinii</i> (Rupr.) Rupr) is a white to pale-yellow powder that crystallizes from hot aqueous solutions.Definition:Chemical name: [(2R,3R)-2-(3,4 dihydroxyphenyl)-3,5,7-trihydroxy-2,3- dihydrochromen-4-one, also called (+) trans (2R,3R)- dihydroquercetin] Chemical formula: $C_{15}H_{12}O_7$ Molecular mass: 304,25 Da CAS No: 480-18-2 Specifications: Physical parameter Moisture: $\leq 10 \%$ Compound analysis Taxifolin (m/m): $\geq 90,0 \%$ of the dry weight Heavy Metals, Pesticide
	Lead: $\leq 0.5 \text{ mg/kg}$ ulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed inII to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012)

commission implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10). Status: Point in time view as at 31/01/2020.

Changes to legislation: There are currently no known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Residual solventEthanol: < 5 000MicrobiologicalTotal Plate CounEnterobacteria: ≤Yeast and MouldEscherichia coli:Salmonella: AbsStaphylococcus aPseudomonas: A	mg/kg ng/kg eltrichloroethane (DDT): $\leq 0,05$ mg/kg is mg/kg criteria t (TPC): $\leq 10^4$ CFU/g $\leq 100/g$ $l: \leq 100$ CFU/g \approx Absence/1 g ence/10 g <i>ureus</i> : Absence/1 g
	Extract component	Content, usual observed range (%)
	Taxifolin	90 - 93
	Aromadendrin	2,5-3,5
	Eriodictyol	0,1-0,3
	Quercetin	0,3-0,5
	Naringenin	0,2-0,3
	Kaempferol	0,01 - 0,1
	Pinocembrin	0,05 - 0,12
	Unidentified flavonoids	1 – 3
	Water(*)	1,5
	crystal.	in in its hydrated form and during the drying process is a This results on the inclusion of water of crystallisation antity of 1,5 %.
Trehalose	by an α-1,1-gluc sucrose by a mul dihydrate. Virtua taste Synonyms: α,α-t	disaccharide that consists of two glucose moieties linkes osidic bond. It is obtained from liquefied starch or from tistep enzymatic process. The commercial product is the lly odourless, white or almost white crystals with a sweet rehalose α -D-glucopyranosyl- α -D-glucopyranoside, dihydrate
	gulation (EU) No 231/2012	of 9 March 2012 laying down specifications for food additives listed in 333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
to the import of) 2015/175 of 5 February 2015 laying down special conditions applicable consigned from India due to contamination risks by pentachlorophenol and

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	Chamical formula: C. H. O 2H. O. (dihydrata)
	Chemical formula: $C_{12}H_{22}O_{11} \cdot 2H_2O$ (dihydrate) Formula weight: 378,33 (dihydrate)
	Assay: ≥ 98 % on the dry basis
	Determine using an atomic absorption technique appropriate to the
	specified level. The selection of sample size and method of sample
	preparation may be based on the principles of the method described in
	FNP 5 (1), 'Instrumental methods'
	Method of assay:
	Principle: trehalose is identified by liquid chromatography and quantified
	by comparison to a reference standard containing standard trehalose
	Preparation of sample solution: weigh accurately about 3 g of dry sample
	into a 100 ml volumetric flask and add about 80 ml of purified, deionised
	water. Bring sample to complete dissolution and dilute to mark with
	purified deionised water. Filter through a 0,45 micron filter
	Preparation of standard solution: dissolve accurately weighed quantities
	of dry standard reference trehalose in water to obtain a solution having known concentration of about 30 mg of trehalose per ml.
	Apparatus: liquid chromatography equipped with a refractive index
	detector and integrating recorder
	Conditions:
	Column: Shodex Ionpack KS-801 (Showa Denko Co.) or equivalent
	length: 300 mm
	— diameter: 10 mm
	— temperature: 50 °C
	Mobile phase: water
	flow rate: 0,4 ml/min
	Injection volume: $8 \mu l$
	Procedure: inject separately equal volumes of the sample solution and the
	standard solution into the chromatograph. Record the chromatograms and measure the size of response of the
	trehalose peak
	Calculate the quantity, in mg, of trehalose in 1 ml of the sample solution
	by the following formula:
	$\%$ trehalose = $100 \times (R_U/R_S) (W_S/W_U)$
	where
	$R_{\rm S}$ = peak area of trehalose in the standard preparation
	$R_{\rm U}$ = peak area of trehalose in the standard preparation $R_{\rm U}$ = peak area of trehalose in the sample preparation
	W_S = yeak area of itenaiose in the sample preparation W_S = weight in mg of trehalose in the standard preparation
	W_U = weight of dry sample in mg
	Characteristics:
	Identification:
	Solubility: Freely soluble in water, very slightly soluble in ethanol
	Specific rotation: $[\alpha]_D^{20} = +179^\circ$ (5 % aqueous solution, dihydrate),
	$+199^{\circ}$ (5 % aqueous solution, anhydrous substance)
	Melting point: 97 °C (dihydrate) Purity:
Commission Decul	ation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in
	to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
p. 1).	

Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of t p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

a

<i>Status:</i> Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)


	Loss on drying: $\leq 1,5 \%$ (60 °C, 5h) Total ash: $\leq 0,05 \%$ Heavy metals: Lead: $\leq 1,0 \text{ mg/kg}$
UV treated mushrooms (<i>Agaricus</i> <i>bisporus</i>)	Description/Definition: Commercially grown <i>Agaricus bisporus</i> to which UV light treatment is applied to harvested mushrooms. UV radiation: a process of radiation in ultraviolet light within the wavelength of 200-800 nm. Vitamin D₂: Chemical name: $(3\beta,5Z,7E,22E)-9,10$ -secoergosta-5,7,10(19),22-tetraen-3-ol Synonym: Ergocalciferol CAS No: 50-14-6 Molecular weight: 396,65 g/mol Contents: Vitamin D ₂ in the final product: 5-10 µg/100 g fresh weight at the expiration of shelf life
UV-treated baker's yeast (<i>Saccharomyces</i> <i>cerevisiae</i>)	Description/Definition: Baker's yeast (<i>Saccharomyces cerevisiae</i>) is treated with ultraviolet light to induce the conversion of ergosterol to vitamin D ₂ (ergocalciferol). Vitamin D ₂ content in the yeast concentrate varies between 1 800 000-3 500 000 IU vitamin D/100 g (450-875 μ g/g). Tan-coloured, free-flowing granules Vitamin D₂: Chemical name: (5Z,7E,22E)-3S-9,10-secoergosta-5,7,10(19),22- tetraen-3-ol Synonym: Ergocalciferol CAS No.: 50-14-6 Molecular weight: 396,65 g/mol Microbiological criteria for the yeast concentrate: Coliforms: $\leq 10^3/$ g <i>Escherichia coli</i> : $\leq 10/$ g <i>Salmonella</i> : Absence in 25g
UV-treated bread	 Description/Definition: UV-treated bread is yeast leavened bread and rolls (without toppings) to which a treatment with ultraviolet radiation is applied after baking in order to convert ergosterol to vitamin D₂ (ergocalciferol). UV radiation: A process of radiation in ultraviolet light within the wavelength of 240-315 nm for maximum of 5 seconds with energy input of 10-50 mJ/cm². Vitamin D₂: Chemical name: (5Z,7E,22E)-3S-9,10-secoergosta-5,7,10(19),22-tetraen-3-ol Synonym: Ergocalciferol
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 5.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

	CAS No: 50-14-6 Molecular weight: 396,65 g/mol Contents:Vitamin D_2 (ergocalciferol) in the final product: 0,75-3 μ g/100 g(¹) Yeast in dough: 1-5 g/100 g (²)(¹)EN 12821, 2009, European Standard.(²)Recipe calculation.
UV-treated milk	Description/Definition: UV-treated milk is cow's milk (whole and semi-skimmed) to which a treatment with ultraviolet (UV) radiation via turbulent flow is applied after pasteurisation. The treatment of the pasteurised milk with UV radiation results in an increase in the vitamin D ₃ (cholecalciferol) concentrations by conversion of 7-dehydrocholesterol to vitamin D ₃ . UV radiation: A process of radiation in ultraviolet light within the wavelength of 200-310 nm with energy input of 1 045 J/l. Vitamin D₃: Chemical name: (1S,3Z)-3-[(2E)-2-[(1R,3aS,7aR)-7a-methyl-1- [(2R)-6-methylheptan-2-yl]-2,3,3a,5,6,7-hexahydro-1H-inden-4- ylidene]ethylidene]-4-methylidenecyclohexan-1-ol Synonym: Cholecalciferol CAS No: 67-97-0 Molecular weight: 384,6377 g/mol Contents: Vitamin D ₃ in the final product: Whole milk(¹)0,5-3,2 µg/100 g(²) Semi-skimmed milk(1): 0,1–1,5 µg/100 g(²) (¹) As defined by Regulation (EU) No 1308/2013 of the Expersent Darliement and a false General of 17 December
	 European Parliament and of the Council of 17 December 2013 establishing a common organisation of the markets in agricultural products and repealing Council Regulations (EEC) No 922/72, (EEC) No 234/79, (EC) No 1037/2001 and (EC) No 1234/2007 (OJ L 347, 20.12.2013, p. 671). (²) HPLC
Vitamin K ₂ (menaquinone)	This novel food is produced by a synthetic or microbiological process. Vitamin K_2 (2-methyl-3-all-trans-polyprenyl-1,4-naphthoquinones), or the menaquinone series, is a group of prenylated naphthoquinone derivatives. The number of isoprene residues, where 1 isoprene unit consists of 5 carbons comprising the side chain, is used to characterise the menaquinone homologues containing primarily MK-7 and to a smaller extent MK-6.
	lation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012,
	ementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable ar gum originating in or consigned from India due to contamination risks by pentachlorophenol and 6.2.2015, p. 10).

Status: Point in time view as at 51/01/2020.
Changes to legislation: There are currently no known outstanding effects for the
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)

21/01/2020



Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Status: Point in time view as at 31/01/2020.	
Changes to legislation: There are currently no known outstanding effects for the	
Commission Implementing Regulation (EU) 2018/1023. (See end of Document for details)	

	Fungi: Max 100/g Salmonella: Absence in 25g Bacillus cereus: Max 1000/g			
	Clostridium perfringens: Max 1000/g			
Yeast beta- glucans	 Clostridium perfringens: Max 1000/g Description/Definition: Beta-glucans are complex, high molecular mass (100–200 kDa) polysaccharides, found in the cell wall of many yeasts and cereals. The chemical name for 'yeast beta-glucans' is (1-3),(1-6)-β-D-glucans. Beta-glucans consist of a backbone of β-1-3-linked glucose residues that are branched by β-1-6-linkages, to which chitin and mannoproteins are linked by β-1-4-bonds. Beta-glucans are isolated from yeast <i>Saccharomyces cerevisiae</i>. The tertiary structure of the glucan cell wall of <i>Saccharomyces cerevisiae</i> consists of chains of β-1,3-linked glucose residues, branched by β-1,6- linkages, forming a backbone to which are linked chitin via β-1,4- bonds, β-1,6-glucans and some mannoproteins. This novel food is available in three different forms: soluble, insoluble and insoluble in water, but dispersible in many liquid matrices. Chemical characteristics yeast (<i>Saccharomyces cerevisiae</i>) beta- glucans: Soluble form: 			
	Total carbohydrates: > 75 %			
	Beta-glucans $(1,3/1,6)$: > 75 %			
	Ash: < 4,0 %			
	Moisture: < 8,0 %			
	Protein: < 3,5 %			
	Fat: < 10 %			
	Insoluble form:			
	Total carbohydrates: $> 70 \%$			
	Beta-glucans $(1,3/1,6)$: > 70 %			
	Ash: $\leq 12 \%$			
	Moisture: < 8,0 %			
	Protein: < 10 %			
	Fat: < 20 %			
	Insoluble in water, but dispersible in many liquid matrices:			
	(1,3)-(1,6)-B-D-Glucans: > 80%			
	Ash: < 2,0 %			
	Moisture: < 6,0 %			
	Protein: < 4,0 %			
	Total fat: < 3,0 %			
	Microbiological data for insoluble in water, but dispersible in many			
	liquid matrices:			
	Total plate count: < 1 000 CFU/g			
	Enterobacteriaceae: < 100 CFU/g			
	Total coliforms: < 10 CFU/g			
	Yeast: < 25 CFU/g			
	Mould: < 25 CFU/g			

 a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Changes to legislation: There are	currently no know	wn outstanding effects for the
Commission Implementing Regulation	(EU) 2018/1023.	(See end of Document for details)

	Salmonella: Absence in 25 g Escherichia coli: Absence in 1 g Bacillus cereus: < 100 CFU/g Staphylococcus aureus: Absence in 1 g Heavy metals for insoluble in water, but dispersible in many liquid matrices: Lead: < 0,2 mg/g Arsenic: < 0,2 mg/g Mercury: < 0,1 mg/g Cadmium: < 0,1 mg/g
Zeaxanthin	Description/Definition: Zeaxanthin is a naturally occurring xanthophyll pigment, it is an oxygenated carotenoid. The synthetic zeaxanthin is presented either as a spray-dried powder of gelatin or starch base ('beadlets') with added α-tocopherol and ascorbyl palmitate or as a corn oil suspension with added α-tocopherol. Synthetic zeaxanthin is produced by a multi-step chemical synthesis from smaller molecules. Orange-red crystalline powder with little or no odour. Chemical formula: $C_{40}H_{56}O_2$ CAS No: 144-68-3 Molecular weight: 568,9 daltons Physical-chemical properties: Loss on drying: < 0,2 % <i>All</i> -trans zeaxanthin: > 96 % Cis-zeaxanthin: < 2,0 % Other carotenoids: < 1,5 % Triphenylphosphine oxid (CAS No 791-28-6): < 50 mg/kg
Zinc L-pidolate	Description/Definition: Zinc L-pidolate is a white to off-white powder, with characteristic odour. International non-proprietary name (INN): L-pyroglutamic acid, Zinc salt Synonyms: Zinc 5-oxoproline, Zinc pyroglutamate, Zinc pyrrolidone carboxylate, Zinc PCA, L-Zinc pidolate CAS No.: 15454-75-8 Molecular formula: $(C_5 H_6 NO_3)_2 Zn$ Relative anhydrous molecular mass: 321,4 Appearance: White to slightly white powder Purity: Zinc L-pidolate (purity): $\geq 98 \%$ pH (10 % aqueous sol.): 5,0-6,0 Specific rotation: 19,6°- 22,8° Water: $\leq 10,0 \%$ Glutamic acid: $< 2,0 \%$ Heavy metals: Lead: $\leq 3,0$ ppm Arsenic: $\leq 2,0$ ppm

 a Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

Cadmium: $\leq 1,0$ ppm Mercury: $\leq 0,1$ ppm **Microbiological criteria:** Total viable mesophilic count: $\leq 1\ 000\ CFU/g$ Yeasts and moulds: $\leq 100\ CFU/g$ Pathogen: Absence

Commission Regulation (EU) No 231/2012 of 9 March 2012 laying down specifications for food additives listed in Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council (OJ L 83, 22.3.2012, p. 1).

b Commission Implementing Regulation (EU) 2015/175 of 5 February 2015 laying down special conditions applicable to the import of guar gum originating in or consigned from India due to contamination risks by pentachlorophenol and dioxins (OJ L 30, 6.2.2015, p. 10).

(**1**) OJ L 327, 11.12.2015, p. 1.

- (2) Regulation (EC) No 258/97 of the European Parliament and of the Council of 27 January 1997 concerning novel foods and novel food ingredients (OJ L 43, 14.2.1997, p. 1).
- (3) Commission Implementing Regulation (EU) 2017/2470 of 20 December 2017 establishing the Union list of novel foods in accordance with Regulation (EU) 2015/2283 of the European Parliament and of the Council on novel foods (OJ L 351, 30.12.2017, p. 72).
- (4) Commission Implementing Decision (EU) 2016/375 of 11 March 2016 authorising the placing on the market of lacto-N-neotetraose as a novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council (OJ L 70, 16.3.2016, p. 22).
- (5) Letter of 8 May 2015 (https://ec.europa.eu/food/sites/food/files/safety/docs/novelfood authorisation 2015 auth-letter krill-oil en.pdf)
- (6) Commission Decision 2009/827/EC of 13 October 2009 authorising the placing on the market of Chia seed (*Salvia hispanica*) as novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council (OJ L 294, 11.11.2009, p. 14).
- (7) Commission Implementing Decision 2014/423/EU of 1 July 2014 authorising the placing on the market of citicoline as a novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council (OJ L 196, 3.7.2014, p. 24).
- (8) Commission Decision 2009/345/EC of 22 April 2009 authorising the placing on the market of Vitamin K₂ (menaquinone) from *Bacillus subtilis* natto as a novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council (OJ L 105, 25.4.2009, p. 16).
- (9) Commission Implementing Decision 2011/762/EU of 24 November 2011 authorising the placing on the market of yeast beta-glucans as a novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council (OJ L 313, 26.11.2011, p. 41).
- (10) Commission Decision 2004/333/EC of 31 March 2004 authorising the placing on the market of yellow fat spreads, salad dressings, milk type products, fermented milk type products, soya drinks and cheese type products with added phytosterols/phytostanols as novel foods or novel food ingredients under Regulation (EC) No 258/97 of the European Parliament and of the Council (OJ L 105, 14.4.2004, p. 40).
- (11) Commission Decision 2008/968/EC of 12 December 2008 authorising the placing on the market of arachidonic acid-rich oil from *Mortierella alpina* as a novel food ingredient under Regulation (EC) No 258/97 of the European Parliament and of the Council (OJ L 344, 20.12.2008, p. 123).

Status:

Point in time view as at 31/01/2020.

Changes to legislation:

There are currently no known outstanding effects for the Commission Implementing Regulation (EU) 2018/1023.