
Status: This is the original version (as it was originally adopted).

		<p>produced by fermentation with <i>Komagataella phaffii</i> CGMCC 12056</p> <p><i>Analytical method^b</i> For the quantification of phytase activity in the feed additive:</p> <p>— colorimetric method based on the enzymatic reaction of phytase on the phytate</p> <p>— VDLUFA 27.1.4</p> <p>For the quantification of phytase activity in premixtures:</p> <p>— colorimetric method based on the enzymatic reaction of</p>				<p>operational procedures and organisational measures to address potential risks resulting from its use. Where those risks cannot be eliminated or reduced to a minimum by such procedures and measures, the additive and premixtures shall be used with personal protective equipment, including breathing protection.</p>
--	--	---	--	--	--	---

a One unit is the amount of enzyme which releases one micromole of inorganic phosphate from phytate per minute at pH 5,5 and 37 °C.

b Details of the analytical methods are available at the following address of the Reference Laboratory: <https://ec.europa.eu/jrc/en/eurl/feed-additives/evaluation-reports>

			<p>phytase on the phytate - VDLUFA 27.1.3</p> <p>For the quantification of phytase activity in feed materials and compound feed: —</p> <p>colorimetric method based on the enzymatic reaction of phytase on the phytate - EN ISO 30024</p>			
--	--	--	--	--	--	--

a One unit is the amount of enzyme which releases one micromole of inorganic phosphate from phytate per minute at pH 5,5 and 37 °C.

b Details of the analytical methods are available at the following address of the Reference Laboratory: <https://ec.europa.eu/jrc/en/eurl/feed-additives/evaluation-reports>