ANNEX I

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

ANNEX I

CLINICAL EXAMINATIONS, SAMPLING PROCEDURES, DIAGNOSTIC METHODS OF CATEGORY A DISEASES AND TRANSPORT OF SAMPLES(as referred to in Article 3 of this Regulation)

A. Sampling procedures

- A.1 SAMPLING OF ANIMALS FOR CLINICAL EXAMINATIONS
- 1. Clinical examinations must include, if possible:
- (a) animals showing clinical signs of category A diseases;
- (b) animals likely to have recently died from the suspected/confirmed disease;
- (c) animals with epidemiological link to a suspected or confirmed case; and
- (d) animals that obtained positive or non-conclusive results in previous laboratory examinations.
- 2. Animals to examine must be selected at random, in a number large enough to allow the detection of the disease, if present, where there are no obvious signs of disease or post-mortem lesions suggesting category A diseases.
- 3. The animals to examine and the sampling method must be chosen in accordance with the instructions of the competent authority and with the relevant contingency plan as referred to in Article 43 of Regulation (EU) 2016/429. The animals to examine and the sampling method must take into account the disease profile and:
- (a) the purpose of the sampling;
- (b) the listed species kept in the establishment;
- (c) the number of animals of listed species kept in the establishment;
- (d) the category of the kept animals;
- (e) the available production, health and traceability records of the kept animals relevant for the investigation;
- (f) the type of establishment and the husbandry practices;
- (g) the level of exposure risk:
 - (i) likelihood of exposure to the disease agent or to the vector;
 - (ii) absence of immunisation of the animals due to vaccination or maternal immunity; and
 - (iii) history of residence in the establishment;
- (h) other relevant epidemiological factors.
- 4. The minimum number of animals to examine must be in accordance with the instructions of the competent authority and with the relevant contingency plan as referred to in Article 43 of Regulation (EU) 2016/429. The minimum number of animals to examine must take into account the disease profile and in particular:
- (a) the expected prevalence in the establishment;

- (b) the level of confidence desired of the survey results, which in any case must not be lower than 95 %; and
- (c) international standards and available scientific evidence.

A.2 SAMPLING OF ANIMALS FOR LABORATORY EXAMINATIONS

- 1. Sampling for laboratory examinations must take into account the outcome of the clinical examinations referred to in point A.1 and, if possible, must include animals referred to in paragraph 1 of point A.1.
- 2. If there are no obvious signs of disease or post-mortem lesions suggesting category A diseases, samples must be collected at random in each epidemiological unit of the establishment and must allow the detection of the disease, if present.
- 3. The animals to sample, the nature of the samples to collect and the sampling method must be in accordance with the instructions of the competent authority and with the relevant contingency plan as referred to in Article 43 of the Regulation (EU) 2016/429. The animals to sample, the nature of the samples to collect and the sampling method must take into account the disease profile and the criteria set out in paragraph 3 of point A.1.
- 4. The minimum number of animals to sample must be in accordance with the instructions of the competent authority and the relevant contingency plan as referred to in Article 43 of the Regulation (EU) 2016/429. The minimum number of animals to sample must take into account the criteria set out in paragraph 4 of point A.1 and the performance of the tests used.
- 5. In the case of wild animals, samples must be collected from animals shot, found dead or purposely trapped or must be obtained on the basis of non-invasive methods such as salt licks and chewing ropes or baits. The minimum number and the nature of the samples must take into account the estimated size of the wild population and the relevant criteria set out in paragraph 3 and 4 of point A.1.

A.3 SAMPLING OF ESTABLISHMENTS FOR VISITS

- 1. The choice of establishments to sample and the sampling method must be in accordance with the instructions of the competent authority and with the relevant contingency plan as referred to in Article 43 of the Regulation (EU) 2016/429. The choice of establishments to sample and the sampling method must take into account the disease profile and the criteria set out in paragraph 3 of point A1.
- 2. The minimum number of establishments to visit must be in accordance with the instructions of the competent authority and with the relevant contingency plan, as referred to in Article 43 of the Regulation (EU) 2016/429.

B. **Diagnostic methods**

The techniques, reference materials, their standardisation and the interpretation of the results of tests carried out using the relevant diagnostic methods for category A diseases must comply with Article 6 and Part III of Annex VI to Delegated Regulation (EU) 2020/689.

The diagnostic methodology must aim to maximise the sensitivity of the surveillance. In certain circumstances this surveillance may include the use of laboratory examinations in order to assess previous exposure to disease.

C. Transport of samples

ANNEX II

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

- 1. All samples taken to confirm or rule out the presence of a category A disease must be sent, with a proper labelling and identification, to an official laboratory which has been informed of their arrival. These samples must be accompanied by the appropriate forms, in accordance with the requirements established by the competent authority and the laboratory receiving the samples. These forms must include at least:
- (a) the establishment of origin of the sampled animals;
- (b) information on the species, age and category of the sampled animals;
- (c) the clinical history of the animals, if available and relevant;
- (d) the clinical signs and post-mortem findings; and
- (e) any other relevant information.
- 2. All samples must be:
- (a) stored in watertight and unbreakable containers and packages and in accordance with applicable international standards;
- (b) kept at the most appropriate temperature and other conditions during transport taking into account the factors that may affect the sample quality.
- 3. The exterior of the package must be labelled with the address of the recipient laboratory and the following message must be prominently displayed:
- 'Animal pathological material; perishable; fragile; do not open outside the laboratory of destination.'
- 4. The person responsible in the official laboratory receiving the samples must be informed in due time of the arrival of the samples.

ANNEX II

MONITORING PERIOD(as referred to in Articles 8, 17, 27, 32, 48, 57 and 59 of this Regulation)

Category A diseases	Monitoring period
Foot and mouth disease (FMD)	21 days
Infection with rinderpest virus (RP)	21 days
Infection with Rift Valley fever virus (RVFV)	30 days
Infection with lumpy skin disease virus (LSD)	28 days
Infection with <i>Mycoplasma mycoides</i> subsp. mycoides SC (Contagious bovine pleuropneumonia) (CBPP)	45 days
Sheep pox and goat pox (SPGP)	21 days
Infection with peste des petits ruminants virus (PPR)	21 days

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

Contagious caprine pleuropneumonia (CCPP)	45 days
African horse sickness (AHS)	14 days
Infection with <i>Burkholderia mallei</i> (Glanders)	6 months
Classical swine fever (CSF)	15 days
African swine fever (ASF)	15 days
Highly pathogenic avian influenza (HPAI)	21 days
Infection with Newcastle disease virus (NCD)	21 days

ANNEX III

CONDITIONS FOR CERTAIN DEROGATIONS FROM ARTICLE 12(1)(a) IN EQUINE ANIMALS(as referred to in Article 13(4))

- 1. In the event of an outbreak of African horse sickness the competent authority may derogate from Article 12(1)(a) the affected and the unaffected animals, provided that:
- (a) the affected animals subject to the derogation are isolated in vector-protected premises which avoid any transmission of the disease agent from the animals to the relevant vectors until 40 days, corresponding to the infective period established in the relevant Chapter of the Terrestrial Animal Health Code of the World Organisation for Animal Health (OIE), have elapsed after the entry of the animals into the vector protected premises; and
- (b) surveillance, including if needed laboratory examinations, carried out by the competent authority, indicates that none of the animals in the vector protected premises poses a risk of virus transmission.
- 2. In the event of an outbreak of infection with *Burkholderia mallei* (Glanders) the competent authority may derogate from Article 12(1)(a) the unaffected animals, provided that the animals subject to the derogation are quarantined until:
- (a) the affected animals have been killed and destroyed;
- (b) after the killing, the cleaning and disinfection of the establishment has been completed as provided for in Article 15; and
- (c) the remaining animals have been subjected to a complement fixation test carried with negative result at a serum dilution of 1 in 5 on samples taken at least 6 months after the cleaning and disinfection referred to in point (b).

ANŃEX IV

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

ANNEX IV

PROCEDURES FOR CLEANING, DISINFECTION AND WHEN NECESSARY CONTROL OF INSECTS AND RODENTS (as referred to in Articles 12, 15, 16, 39, 45 and 57 of this Regulation)

A. General requirements

- 1. The choice of biocidal products and procedures for cleaning and disinfection operations must take into account:
- (a) the causal agent of infection;
- (b) the nature of the establishments, vehicles, objects and materials which are to be treated; and
- (c) the applicable legislation.
- 2. The conditions under which biocidal products are used must ensure that their efficacy is not impaired. In particular technical parameters provided by the manufacturer, such as pressure, temperature, required contact time or storage must be observed. The activity of the disinfectant must not be compromised by interaction with other substances.
- 3. Re-contamination of the previously cleaned parts must be avoided, in particular where washing is carried out with liquids applied under pressure.
- 4. The water used for cleaning operations must be contained and disposed of in a way that avoids any risk of spreading category A disease agents.
- 5. Biocidal products must be used in a way that reduces as much as possible any adverse impact on the environment and on public health that may arise from their use.

B. Preliminary cleaning and disinfection

For preliminary cleaning and disinfection under Article 15, to avoid spreading the category A disease:

- (a) entire bodies or parts of dead kept animals of listed species must be sprayed with disinfectant and removed from the establishment, in closed and leak-proof vehicles or containers for processing and disposal;
- (b) any tissue or blood which may have been spilled during killing, slaughter or postmortem examination must be carefully collected and disposed of;
- (c) as soon as the entire bodies or parts of dead kept animals of listed species have been removed for processing or disposal, the parts of the establishment in which these animals were kept and any parts of other buildings, surfaces or equipment contaminated during killing or post-mortem examination must be sprayed with disinfectant;
- (d) manure, including litter and used bedding, must be thoroughly soaked with disinfectant:
- (e) the disinfectant must remain on the treated surface for at least 24 hours;
- (f) equipment, containers, consumption utensils, surfaces or any material likely to be contaminated after the washing and disinfecting must be destroyed.

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

C. Final cleaning and disinfection:

For final cleaning and disinfection for the purpose of Article 57:

- 1. Manure, including litter and used bedding, must be removed and treated as follows:
 - (a) the solid phase of manure, including litter and used bedding, must either:
 - (i) undergo a steam treatment at a temperature of at least 70 °C;
 - (ii) be destroyed by burning;
 - (iii) be buried deep enough to prevent access by animals; or
 - (iv) be stacked to heat, sprayed with disinfectant and left for at least 42 days, during which the stack must be either covered or re-stacked to ensure thermic treatment of all layers;
 - (b) the liquid phase of manure must be stored for at least 42 days, and in the case of highly pathogenic avian influenza 60 days, after the last addition of infective material.
- 2. Buildings, surfaces and equipment must be thoroughly washed and cleaned by removing the remaining grease and dirt and sprayed with disinfectants.
- 3. After 7 days the establishments must be cleaned and disinfected again.

ANNEX V

MINIMUM RADIUS OF PROTECTION AND SURVEILLANCE ZONES(as referred to in Article 21 of this Regulation)

Indicated as radius of a circle centred on the establishment

Category A diseases	Protection Zone	Surveillance Zone		
Foot and mouth disease	3 km	10 km		
Infection with rinderpest virus	3 km	10 km		
Infection with Rift Valley fever virus	20 km	50 km		
Infection with lumpy skin disease virus	20 km	50 km		
Infection with Mycoplasma mycoides subsp. mycoides SC (Contagious bovine pleuropneumonia)	Establishment	3 km		
Sheep pox and goat pox	3 km	10 km		
Infection with peste des petits ruminants virus	3 km	10 km		

ANNEX VI

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

Contagious caprine pleuropneumonia	Establishment	3 km		
African horse sickness	100 km	150 km		
Infection with <i>Burkholderia</i> mallei (Glanders)	Establishment	Establishment		
Classical swine fever	3 km	10 km		
African swine fever	3 km	10 km		
Highly pathogenic avian influenza	3 km	10 km		
Infection with Newcastle disease virus	3 km	10 km		

ANNEX VI

PROHIBITIONS IN THE RESTRICTED ZONE(as referred to in Article 27 of this Regulation)

Table: Prohibitions of activities concerning animals of listed species and products from those animals

PRO	HIRA	TERN	SRVE	M.SD	CRP	PSPG	PPPR	CCF	PCSE	ASE	AHS	GLA	NIPA	INCD
OF	- 1-1414	J 101 (VLSD	CDI	151 0	1111		1031	7101	71110	GL	1111111	псь
-	IVIT	ES												
		NING												
	MALS													
AND														
PRO	DUC '	ΓS												
Move	nXents	X	X	X	X	X	X	X	X	X	X	NA	X	X
of														
kept														
anima	ıls													
of														
listed														
specie	es													
from														
establ	ishme	nts												
in														
the														
restric	eted													
zone														

a	only oocytes and embryo
---	-------------------------

b only oocytes and embryo.

c Disease abbreviations in accordance with Annex II.

NA = Not applicable. X = prohibition. NP = Not prohibited.

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

MoverXents of kept animals of listed species to establishme in the restricted		X	X	X	X	X	X	X	X	X	NA	X	X
Restocking of game animals of listed species	X	X	X	X	X	X	X	X	X	X	NA	X	X
Fairs, X markets, shows and other gatherings of kept animals of listed species including collection and dispersion of those species	X	X	X	X	X	X	X	X	X	X	NA	X	X
MoverMents of semen, oocytes and	X	X	X ^b	X	X	X	X	X	X	X	NA	NA	NA

a	only	oocytes	and	amhr	70
а	OHITY	oocytes	and	CHIDI	yO.

only oocytes and embryo.

Disease abbreviations in accordance with Annex II.

NA X NP

Not applicable. prohibition. Not prohibited.

=

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

embryos obtained from kept animals of listed species from establishme in the restricted zone	ents												
Collection of semen, oocytes and embryo from kept animals of listed species	X	X	X	X	X	X	X	X	X	NP	NA	NA	NA
Itinerakt artificial inseminatio of kept animals of listed species	X n	X	X	X	X	X	X	X	X	X	NA	NA	NA
ItineraXt natural service of kept animals of listed species	X	X	X	X	X	X	X	X	X	X	NA	NA	NA

a	only	oocytes	and	embr	70
а	OHIY	oocytes	anu	CIIIOI	yυ.

only oocytes and embryo.

Disease abbreviations in accordance with Annex II.

NA X NP

Not applicable. prohibition. Not prohibited. =

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

Movements of hatching eggs from establishme in the restricted zone		NA	X	X									
MoverXents of fresh meat excluding offal from kept and wild animals of listed species from slaughterho or game handling establishme in the restricted zone	uses	X	NP	NP	X	X	NP	X	X	NP	NA	X	X
MoverMents of offal from kept and wild animals of listed species	X	X	X	X	X	X	X	X	X	NP	NA	X	X

a or	ılv oocyt	ec and er	nhrvo

b only oocytes and embryo.

Disease abbreviations in accordance with Annex II.

NA	=	Not applicable.
X	=	Not applicable. prohibition.
NP	=	Not prohibited.

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

from slaughterho or game handling establishme in the restricted zone													
MoverMents of meat products obtained from fresh meat of listed species from establishme in the restricted zone		X	NP	NP	NP	X	NP	X	X	NP	NA	X	X
MoverXent of raw milk and colostrum obtained from kept animals of listed species from establishme in the	X	X	X	NP	X	X	NP	NA	NA	NP	NA	NA	NA

a	only	oocytes	and	embry	VO.

Disease abbreviations in accordance with Annex II.

Not applicable. prohibition. Not prohibited.

only oocytes and embryo.

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

restricted zone													
MoverMent of dairy products and colostrum based products from establishme in the restricted zone	X	X	X	NP	X	X	NP	NA	NA	NP	NA	NA	NA
Moverheat of eggs for human consumptio from establishme in the restricted zone		NA	X	X									
MoverXent of manure, including litter and used bedding from kept animals of listed species from establishme in	X	X	X	NP	X	X	NP	X	X	NP	NA	X	X

a	only	oocytes	and	embryo
---	------	---------	-----	--------

only oocytes and embryo.

Disease abbreviations in accordance with Annex II.

=

Not applicable. prohibition. Not prohibited. =

ANNEX VII

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

the restricted zone													
MoverXent of hides, skins, wool, bristles and feathers from kept animals of listed species from establishme in the restricted zone	ents	X	X	NP	X	X	NP	X	X	NP	NA	X	X
MoverXent of feed material of plant origin and straw obtained in the protection zone ^a	X	NP	NA	NP	NP								

	a	only	oocytes	and	embryo.
--	---	------	---------	-----	---------

NA = Not applicable. X = prohibition. NP = Not prohibited.

ANNEX VII

RISK-MITIGATING TREATMENTS FOR PRODUCTS OF ANIMAL ORIGIN FROM THE RESTRICTED ZONE (as referred to in Articles 27, 33 and 49 of this Regulation)

b only oocytes and embryo.

Disease abbreviations in accordance with Annex II.

ANNEX VII

Document Generated: 2023-11-13

Treatment N RP	RVFVLSD	CBPP SPGP PPR	CCPPCSF	ASF	AHS	HPA	I NCD
MEAT							
Heat x treatment in an hermetically sealed container, to achieve a minimum F_0^g value of 3		X	X	X		X	X
Heat X treatment to achieve a core temperature of 80 °C		X	X	X		X	X
Heat X treatment to achieve a core temperature of 70 °C		X	X			X	X
Heat X treatment (to meat previously		X	X				

- Safe commodity.
- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- ${f g}$ F₀ is the calculated killing effect on bacterial spores. An F₀ value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

ANŃEX VII

Document Generated: 2023-11-13

de-boned and defatted) to achieve a core temperature of 70 °C for a minimum of 30 minutes								
In an X hermetically sealed container, applying 60 °C for a minimum of 4 hours				X	X	X		
Core X temperature of 73,9 °C for a minimum of 0,51 seconds ^f							X	X
Core temperature of 70,0 °C for a	odity.						X	X

- Safe commodity.
- b Not for bovine, ovine, caprine and porcine casings.
- Not for bovine, ovine, caprine and porcine casings. c
- d Safe commodity.
- e Only for porcine animals.
- Only for poultry meat.
- F_0 is the calculated killing effect on bacterial spores. An F_0 value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- Disease abbreviations in accordance with Annex II.

ANNEX VII Document Generated: 2023-11-13

minimu of 3,5 seconds									
Core tempera of 65,0 °C for a minimu of 42 seconds	ım							X	X
Core tempera of 60 °C for a minimu of 507 seconds	ım							X	X
Heat treatment to achieve desiccato maximum values of Aw of 0,93 and pH of 6	tion								
Heat treatme	ent	. 1:4.			X				

- a Safe commodity.
- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- g F₀ is the calculated killing effect on bacterial spores. An F₀ value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

ANNEX VII

Document Generated: 2023-11-13

to		
achieve		
a		
core		
temperature		
of		
65 °C		
for a		
period		
of		
time		
to		
achieve		
a		
minimum		
pasteurisation		
value		
of 40		
01 40		
NaturaK	X	
fermentation		
and		
maturation		
for		
bone-		
in		
meat:		
minimum		
9		
months,		
to		
achieve		
maximum		
values		
of		
Aw		
of		
0,93		
and		
pН		
of 6		
a Safe commodity.		

- a Safe commodity.
- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- **g** F_0 is the calculated killing effect on bacterial spores. An F_0 value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

ANNEX VII
Document Generated: 2023-11-13

NaturaK	X	X	
fermentation		11	
and			
maturation			
for			
de-			
boned			
meat:			
minimum			
9			
months,			
to			
achieve			
maximum			
values			
of			
Aw			
of			
0,93			
and			
pH			
pH of 6			
pH of 6 Natural	X	X	
pH of 6 Natural fermentation	X	X	
pH of 6 Natural fermentation for	X	X	
pH of 6 Natural fermentation	X	X	
pH of 6 Natural fermentation for loins:	X	X	
pH of 6 Natural fermentation for loins: minimum	X	X	
pH of 6 Natural fermentation for loins: minimum 140	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values of	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values of Aw	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values of Aw of	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values of Aw of 0,93	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values of Aw of 0,93 and	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values of Aw of 0,93 and pH	X	X	
pH of 6 Natural fermentation for loins: minimum 140 days to achieve maximum values of Aw of 0,93 and	X	X	

- Safe commodity.
- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- **g** F₀ is the calculated killing effect on bacterial spores. An F₀ value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

ANŃEX VII

Document Generated: 2023-11-13

	,			
Natural	X	X		
fermentation				
for				
hams:				
minimum				
190				
days				
to				
achieve				
maximum				
values				
of				
Aw				
of				
0,93				
0,93				
and				
pH				
of 6°				
Drying	X			
after				
salting				
Italian				
style				
bone-				
in				
hams:				
minimum				
313				
days ^e				
DryingX	X	X		
after				
salting				
Spanish				
style				
bone-				
in				
hams				
and				
loins ^e :				
a Safa commodity				

- a Safe commodity.
- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- ${f g}$ F₀ is the calculated killing effect on bacterial spores. An F₀ value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

ANNEX VII
Document Generated: 2023-11-13

— Iberian hams: minimum 252 days — Iberian shoulders:	
minimum 252 days — Iberian	
252 days — Iberian	
days — Iberian	
— Iberian	
minimum	
140	
days	
— Iberian	
loins:	
minimum	
126	
days	
— Serrano	
hams:	
minimum	
140	
days	
Maturation X	
of	
carcasses	
at a	
minimum town and the second section in the section in the section in the second section in the	
temperature	
of 2 °C	
for a	
minimum	
of 24	
hours	
following	
following slaughter	
Removal X X X	
of	
offal	
CASINGS	
a Safe commodity.	

- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- **g** F₀ is the calculated killing effect on bacterial spores. An F₀ value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

ANNEX VII

Document Generated: 2023-11-13

SaltingX with sodium chloride (NaCl) either dry or as saturated brine (Aw < 0,8), for a continuous period of 30 days or longer at an ambient temperature of 20 °C	SC ^d	X	X	X		
or above						
SaltingX with phosphate supplemented salt 86,5 % NaCl, 10,7 % Na ₂ HPO ₄ and 2,8 %		X	X	X		

- a Safe commodity.
- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- **g** F₀ is the calculated killing effect on bacterial spores. An F₀ value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

ANNEX VII

Document Generated: 2023-11-13

$Na_3P\Phi_4$							
either							
dry							
or as saturated							
brine							
(Aw							
< 0,8)							
for a							
continuous							
period							
of 30							
days or							
longer							
at an							
ambient							
temperature							
of 20 °C							
or or							
above							
Salting							
with							
sodium							
chloride							
(NaCl)							
minimum							
30							
days ^c							
Bleachingb							
Drying ^b							
MILK							
Heat X		SCa		SC ^a			
treatment							
(sterilizatio	n						
process) to							
a Safe comr	nodity.			1			
b Not for bovine, caprine and porcine casings.							
- 100 101 00 tille, otille, captille and potenic casings.							

- Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- F_0 is the calculated killing effect on bacterial spores. An F_0 value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- Disease abbreviations in accordance with Annex II.

ANŃEX VII

chilling.

Disease abbreviations in accordance with Annex II.

Document Generated: 2023-11-13

achi a min F ₀ valu of 3	imum e											
UH' (Ult high tem Mir 132 for a	tment ra perature) imum °C a imum	-					X					
UH' (Ult high tem	tment ra perature) imum °C a abble ling	:										
HTS (High tem short time	tment ST gh perature t	n)					X					
a												
b												
	Not for bovine, ovine, caprine and porcine casings.											
d	Safe commodity.											
	Only for porcine animals.											
	Only for poultry meat.											
	F_0 is the calculated killing effect on bacterial spores. An F_0 value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and											

(EU) 2016/429 of the... ANNEX VII

Document Generated: 2023-11-13 ation: There are currently no known outstanding effects for the

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

milk pH is lower than 7, minimum 72 °C for a minimum of 15 seconds Heat X treatment HTST (High temperature short								
treatment HTST (High temperature short								
time pasteurisation) if milk pH is 7 or higher, minimum 72 °C for a minimum of 15 seconds, applied twice								
Heat X treatment HTST (High temperature short								
Safe commodity.								
Not for bovine, ovine, caprine and porcine casings.								
 b Not for bovine, ovine, caprine and porcine casings. c Not for bovine, ovine, caprine and porcine casings. 								
Safe commodity.								
Only for porcine animals.								
e Only for porcine animals.								

 F_0 is the calculated killing effect on bacterial spores. An F_0 value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.

Disease abbreviations in accordance with Annex II.

ANNEX VII

Document Generated: 2023-11-13

time pasteurisation) combined with a physical treatment to achieve pH value below 6 for a minimum of 1 hour or to achieve a minimum of 72 °C, combined with										
Pasteur Sation Consisting in a single heat treatment with an effect at least equivalent to that a Safe commodity.	X	X								
b Not for bovine, ovine, caprine and porcine casings.										

- Not for bovine, ovine, caprine and porcine casings. c
- Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- F_0 is the calculated killing effect on bacterial spores. An F_0 value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- Disease abbreviations in accordance with Annex II.



- a Safe commodity.
- **b** Not for bovine, ovine, caprine and porcine casings.
- c Not for bovine, ovine, caprine and porcine casings.
- d Safe commodity.
- e Only for porcine animals.
- f Only for poultry meat.
- ${f g}$ F₀ is the calculated killing effect on bacterial spores. An F₀ value of 3 means that the coldest point in the product has been heated sufficiently to achieve the same killing effect as 121 °C (250 °F) in three minutes with instantaneous heating and chilling.
- h Disease abbreviations in accordance with Annex II.

Treatn	nent	HPAI	NCD
EGGS			
Heat tre	eatment:	X	
	Whole egg:		
		,0 °C –	
		8 sec.	
		mpletely	
		oked	
_	Whole egg b		
	— 60	°C –	
		8 sec.	
		mpletely	
		oked	
		,1 °C –	
		sec.	
	Liquid egg v		
		,6 °C –	
		0 sec.	
		,7 °C –	
		2 sec.	
	Plain or pure		
	yolk:	Ceg	
		°C –	
		8 sec.	
	10 % salted		
_			
		,2 °C – 8 sec.	
	Dried egg w		
		°C – 20	
	ho		
		,4 °C –	
	50,	,4 hours	

ANNEX VIII

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

	_	51,7 °C – 73,2 hours	
Heat tre	atment:		X
_	Whole 6	egg.	
	_	55 °C – 2	
		521 sec.	
		57 °C − 1	
		596 sec.	
		59 °C –	
		674 sec.	
		completely	
		cooked	
	Lianid e	egg white:	
	—	55 °C – 2	
		278 sec.	
		57 °C –	
		986 sec.	
		59 °C –	
		301 sec.	
	10 % sa	lted egg	
	yolk:	inca egg	
	yon.	55 °C −	
		176 sec.	
	Dried e	gg white:	
		57 °C –	
		54,0 hours	
		54,0 Hours	

ANNEX VIII

RISK-MITIGATING TREATMENTS FOR PRODUCTS NOT OF ANIMAL ORIGIN FROM THE PROTECTION ZONE(as referred to in Articles 36 and 52 of this Regulation)

Treatment	FMD ^a	RP	
Heat treatment, minimum temperature of 80 °C and for a minimum of 10 minutes, steam in a closed chamber	X	X	
Storage in package or bales under shelter at premises situated not closer than 2 km to the nearest outbreak and releasing from the premises do not take place before at least three months have elapsed following the completion of cleaning and	X	X	
a Disease abbreviations in accordance	with Annex II.		

ANNEX X

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

disinfection according to Article 15			
a	Disease abbreviations in accordance	with Annex II.	

ANNEX IX

MARKING OF FRESH MEAT FROM THE PROTECTION ZONE(as referred to in Articles 33 and 49 of this Regulation)

- 1. The mark to be applied to fresh meat of poultry originating in the protection zone and not intended to another Member State pursuant to Article 33(1)(b) must comply with the following:
- (a) shape and content:

Where 'XY' means the relevant country code provided for in point 6 of Part B of Section I of Annex II of Regulation (EC) No 853/2004 and '1234' means the approval number of the establishment referred to in point 7 of Part B of Section I of Annex II of Regulation (EC) No 853/2004;

- (b) dimensions:
 - 'XY' width of 8 mm,
 - '1234' width of 11 mm,
 - width outer diameter of not less than 30 mm,
 - line thickness of square of 3 mm.
- 2. The mark to be applied to fresh meat intended for treatment in a processing plant pursuant to Article 33(2)(a) shall consist in, either:
- (a) the identification mark provided for in Regulation (EC) No 853/2004 with an additional diagonal cross consisting of two straight lines intersecting at the centre of the stamp and enabling the information thereon to remain legible; or
- (b) a single oval stamp, 6,5 cm wide by 4,5 cm high, in which the following information must appear in perfectly legible characters:
 - on the upper part, the full name or ISO code of the Member State in capitals,
 - in the centre, the approval number of the slaughterhouse,
 - on the lower part, one of the following sets of initials CE, EC, EF, EG, EK, EY, EO, ES, EU, EB, WE or EZ,
 - two straight lines crossing at the centre of the stamp in such a way that the information is not obscured,
 - the letters must be at least 0,8 cm high and the figures at least 1 cm high.

ANNEX X

DURATION OF THE MEASURES IN THE PROTECTION ZONE(as referred to in Article 39 of this Regulation)

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

Category A diseases	Minimum period of duration of measures in the protection zone (Article 39(1))	Additional period of duration of surveillance measures in the protection zone (Article 39(3))
Foot and mouth disease	15 days	15 days
Infection with rinderpest virus	21 days	9 days
Infection with Rift Valley fever virus	30 days	15 days
Infection with lumpy skin disease virus	28 days	17 days
Infection with <i>Mycoplasma mycoides subsp. mycoides SC</i> (Contagious bovine pleuropneumonia)	45 days	Not applicable
Sheep pox and goat pox	21 days	9 days
Infection with peste des petits ruminants virus	21 days	9 days
Contagious caprine pleuropneumonia	45 days	Not applicable
African horse sickness	12 months	Not applicable
Infection with <i>Burkholderia</i> mallei (Glanders)	6 months	Not applicable
Classical swine fever	15 days	15 days
African swine fever	15 days	15 days
Highly pathogenic avian influenza	21 days	9 days
Infection with Newcastle disease virus	21 days	9 days

ANNEX XI

DURATION OF THE MEASURES IN THE SURVEILLANCE ZONE(as referred to in Articles 55 and 56 of this Regulation)

Category A diseases	Minimum period of duration of measures in the surveillance zone
Foot and mouth disease	30 days
Infection with rinderpest virus	30 days
Infection with Rift Valley fever virus	45 days
Infection with lumpy skin disease virus	45 days

Infection with <i>Mycoplasma mycoides</i> subsp. mycoides SC (Contagious bovine pleuropneumonia)	45 days
Sheep pox and goat pox	30 days
Infection with peste des petits ruminants virus	30 days
Contagious caprine pleuropneumonia	45 days
African horse sickness	12 months
Infection with <i>Burkholderia mallei</i> (Glanders)	Not applicable
Classical swine fever	30 days
African swine fever	30 days
Highly pathogenic avian influenza	30 days
Infection with Newcastle disease virus	30 days

ANNEX XII

SAMPLING PROCEDURES AND DIAGNOSTIC METHODS FOR CATEGORY A DISEASES IN AQUATIC ANIMALS

- 1. The following procedures apply to the clinical examination and collection of samples:
- (a) the clinical examination and the sampling for laboratory examinations must include:
 - (i) aquaculture animals of listed species showing clinical signs of the relevant category A disease; and
 - (ii) aquaculture animals likely to have recently died from the suspected/confirmed category A disease; and
 - (iii) aquaculture animals with an epidemiological link to a suspected or confirmed case of a category A disease;
- (b) the minimum number of samples to be collected is:

	Scenario			
Type of animals	Report of increased mortality	Introduction of infected animals	Post- mortem or clinical signs observed	Suspicion based on other circumstances
Molluscs (the whole animal)	30	30	_	150
Crustaceans	10		10	150
Fish	_	_	10	30

(c) the following additional criteria apply to the sampling of molluscs:

- (i) animals suspected to be infected must be selected for sampling. If listed species are present in the population of animals concerned by the suspicion, those must be selected for sampling;
- (ii) if weak, gaping or freshly dead but not decomposed molluscs are present, those must be selected first. If such molluscs are not present, the molluscs selected must include the oldest healthy molluscs;
- (iii) if the establishment uses more than one water source for mollusc production, molluscs representing all water sources must be included for sampling to ensure that all parts of the establishment are proportionally represented in the sample;
- (iv) when sampling from a group of mollusc farming establishments which apparently have identical epidemiological status, molluscs from a representative number of sampling points must be included in the sample.

The main factors to be considered when selecting those sampling points must be stocking density, water currents, the presence of listed species, both susceptible and vector species, bathymetry and management practices. Natural beds within or adjacent to the mollusc farming establishment(s) must be included in the sample;

- (d) the following additional criteria apply when sampling crustaceans:
 - (i) if weak or moribund crustaceans of listed species are present in the production units, those crustaceans must be selected first. If such animals are not present, the crustaceans selected must include crustaceans of different year classes, proportionally represented in the sample;
 - (ii) if more than one water source is used for crustacean production, crustaceans of listed species representing all water sources must be included in the sample to ensure that all parts of the establishment are proportionally represented in the sample;
 - (iii) when collection of samples from wild populations of listed species is required under Article 102(a) of this Regulation, the number and geographical distribution of the sampling points must be determined in a way that ensures a reasonable coverage of the area suspected to be infected.

The sampling points must be representative for the different ecosystems where the wild populations of susceptible species are located such as marine, estuary, river and lake systems;

- (e) the following additional criteria apply for sampling fish:
 - (i) if weak, abnormally behaving or freshly dead but not decomposed fish are present, those fish must be selected. If such animals are not present, the fish selected must include fish of listed species, belonging to different year classes, proportionally represented in the sample;
 - (ii) if more than one water source is used for fish production, listed species representing all water sources must be included for sampling to ensure that all parts of the establishment are proportionally represented in the sample;

- (iii) if rainbow trout (*Onchorynchus mykiss*) or European perch (*Perca fluviatilis*) are present, only fish of those species may be selected for sampling. If neither rainbow trout nor European perch are present, the sample must be representative of all other listed species present, following the criteria in points (a) to (d);
- (iv) when collection of samples from wild populations of listed species is required under Article 102(a) of this Regulation, the number and geographical distribution of the sampling points must be determined in a way that ensures a reasonable coverage of the area suspected to be infected.

The sampling points must also be representative of the different ecosystems where the wild populations of susceptible species are located such as marine, estuary, river and lake systems;

- (f) the selection of organs to be sampled, preparation, storage and shipment of the samples to the laboratory must be carried out in compliance with recommendations from the European Union reference laboratory for the relevant disease.
- 2. Samples must be examined in the laboratory using the diagnostic methods and procedures approved by the European Union reference laboratory for the relevant disease.

ANNEX XIII

MINIMUM PERIODS OF FALLOWING OF AFFECTED AQUACULTURE ESTABLISHMENTS

PERIODS FOR THE FALLOWING PROVIDED FOR IN ARTICLE 81 AND FOR THE SYNCHRONOUS FALLOWING PROVIDED FOR IN ARTICLE 96(4) AND (5) OF THIS REGULATION

Category A disease	Minimum period of fallowing of the affected establishment	Minimum period of synchronised fallowing of affected establishments in the same protection zone	Supplementary requirements
Infection with Mikrocytos mackini	6 months	4 weeks	Must include the coldest period of the year
Infection with Perkinsus marinus	6 months	4 weeks	Must include the warmest period of the year
Infection with Taura syndrome virus	6 weeks	4 weeks	Must include the warmest period of the year

ANNEX XIV

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

Infection with Yellow head syndrome virus	6 weeks	3 weeks	Must include the warmest period of the year
Epizootic haematopoietic necrosis	8 weeks	4 weeks	Must include the warmest period of the year

ANNEX XIV

CRITERIA FOR ESTABLISHING RESTRICTED ZONES AS REGARDS CATEGORY A DISEASES IN AQUATIC ANIMALS

- 1. Restricted zones as referred to in Article 85 must be defined on a case-by-case basis taking into account at least the following factors:
- (a) the accumulated number, the accumulated percentage and the distribution of the mortalities of molluscs/crustaceans/fish in the establishment or group of farming establishments infected with category A diseases;
- (b) relevant information regarding movements to and from the infected establishment(s);
- (c) the distance to and density of neighbouring establishments;
- (d) the presence of wild aquatic animals;
- (e) any knowledge concerning mortalities, suspected cases or outbreaks in wild aquatic animals which are, or could be related to the specific category A disease;
- (f) the proximity to processing establishments, and the species present at those establishments, especially as regards listed species;
- (g) farming practices applied in the affected and neighbouring establishments;
- (h) hydrodynamic conditions and other identified factors of epidemiological significance.
- 2. For the geographical demarcation of the protection and surveillance zones for category A diseases affecting molluscs and crustaceans, the following minimum requirements apply:
- (a) the protection zone must be established in the immediate vicinity of an establishment or group of farming establishments officially confirmed as infected with a category A disease and must correspond to an area determined according to appropriate hydrodynamic and epidemiological data;
- (b) the surveillance zone must be established outside the protection zone and must correspond to an area surrounding the protection zone, determined according to appropriate hydrodynamic or epidemiological data.
- 3. For the geographical demarcation of the protection and surveillance zones for category A diseases affecting fish, the following minimum requirements must apply:
- (a) the protection zone must be established around an establishment where *Epizootic hematopoietic necrosis* (EHN) has been confirmed. This zone shall correspond:

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

- (i) in coastal areas: to an area included in a circle with a radius of at least one tidal excursion or at least 5 km, whichever is larger, centred on the establishment in which EHN has been officially confirmed, or an equivalent area determined according to appropriate hydrodynamic or epidemiological data;
- (ii) in inland areas: to the entire water catchment area of the establishment in which EHN has been officially confirmed. The competent authority may limit the extension of the zone to parts of the water catchment area, or the area of the establishment, provided this does not compromise prevention of the spread of the disease;
- (b) the surveillance zone must be established by the competent authority outside the protection zone and must:
 - (i) in coastal areas: correspond to an area, surrounding the protection zone, of overlapping tidal excursion; or an area, surrounding the protection zone, and included in a circle of radius 10 km from the centre of the protection zone; or an equivalent area determined according to appropriate hydrodynamic or epidemiological data;
 - (ii) in inland areas: be an extended area outside the established protection zone.

ANNEX XV

SURVEILLANCE SCHEME AND DURATION OF CONTROL MEASURES IN THE SURVEILLANCE ZONE FOR CATEGORY ADISEASES IN AQUACULTURE ANIMALS(as referred to in Articles 98 and 101 of this Regulation)

1. Surveillance scheme

The establishments and groups of aquaculture establishments keeping listed species within a surveillance zone must undergo surveillance as provided for in Article 98 to check for infection with the relevant category A disease. The surveillance must include health visits, including sampling from production units. Those visits must be carried out by the competent authority in accordance with Tables 1 and 2.

The criteria set out in point 1 of Annex XII, as appropriate for the species, apply to sampling.

TABLE 1

Scheme for surveillance comprising health visits and samplings in establishments and groups of establishments for category A diseases in aquatic animals, except epizootic hematopoietic necrosis

Category A disease	Number of health visits per year	Number of laboratory examinations per year	Number of animals in the sample	Period of the year for sampling	Residency period of the sampled animals in the establishment
--------------------	---	---	---------------------------------	---------------------------------------	--

ANNEX XV

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

Infection with Mikrocytos mackini	1	1	150	When the prevalence of infection is known to be maximal or April–May, after 3–4 months period when seawater temperatures are less than 10 °C	4 months
Infection with Perkinsus marinus	1	1	150	When the prevalence of infection is known to be maximal or in the month of September, October or November	4 months
Infection with Taura syndrome virus	2	2	150	In the period of the year when water temperature is likely to reach its highest annual level	2 months
Infection with Yellow head syndrome virus	2	2	150	In the period of the year when water temperature is likely to reach its highest annual level	2 months

TABLE 2

Specific scheme for surveillance comprising health visits and samplings in establishments for epizootic haematopoietic necrosis (EHN) in aquatic animals⁰

1	1	•	,	
Type of				Number of fish in the sample
establishment				

- a The sampling of fish for laboratory examination must be carried out whenever the water temperature is between 11 and 20 °C. The water temperature requirement must also apply to health inspections. In establishments where the water temperature does not reach 11 °C during the year, sampling and health visits must be carried out when the water temperature is at its highest level.
- b Samples from broodstock must not include gonadal fluids, milt or ova as there is no evidence of EHN causing reproductive tract infection.

ANNEX XV

Document Generated: 2023-11-13

Changes to legislation: There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687. (See end of Document for details)

		Number of health inspections per year (2 years)	Number of samplings per year (2 years)	Number of growing fish	Number of brood stock fish ^b
(a)	Establish with broodsto		2	150 (first and second inspection)	150 (first or second inspection)
(b)	Establish with broodsto only		1	0	150 ^b (first or second inspection)
(c)	Establish without broodsto		2	150 (first and second inspection)	0

Maximum number of fish per pool: 10

- a The sampling of fish for laboratory examination must be carried out whenever the water temperature is between 11 and 20 °C. The water temperature requirement must also apply to health inspections. In establishments where the water temperature does not reach 11 °C during the year, sampling and health visits must be carried out when the water temperature is at its highest level.
- b Samples from broodstock must not include gonadal fluids, milt or ova as there is no evidence of EHN causing reproductive tract infection.

2. Duration of the control measures in the surveillance zone

Category A disease	Minimum periods of surveillance
Infection with Mikrocytos mackini	3 years
Infection with Perkinsus marinus	3 years
Infection with Taura syndrome virus	2 years
Infection with Yellow head syndrome virus	2 years
Epizootic haematopoietic necrosis	2 years

When the period of surveillance has elapsed and there has been no new detection of infection with the relevant category A disease, the measures in the surveillance zone must be lifted as provided for in Article 101 of this Regulation.

Changes to legislation:

There are currently no known outstanding effects for the Commission Delegated Regulation (EU) 2020/687.